

- Abbas A, Akram HM, Iqbal MS, Allah Y, Nadeem MA, Saeed M, Sahi KA (2004) Agro-economic relationship of component crops in a mash sesamum intercropping system. *Pakistan Journal of Biological Sciences* **7**(1), 69-72.
- Abbasi F, Naeem M, Khan HM (1981) Trace metal contents of undecorticated and decorticated dry legumes. *Pakistan Journal of Biochemistry* **14**(1), 33-38.
- Abdul J, Riaz A, Ehsan U, Nazir MS (2005) Agronomic approaches for productivity enhancement of upland rice-based intercropping systems under strip plantation. *Pakistan Journal of Agricultural Sciences* **42**(1/2), 14-17.
- Abe N, Sato H, Sakamura S (1987) Antifungal stress compounds from adzuki bean, *Vigna angularis*, treated with *Cephalosporium gregatum* type B. *Agricultural and Biological Chemistry* **51**(2), 349-354.
- Abe Y, Nakamura T, Inoue H (2000) Exploitation of Fabaceae plants by the Mexican bean beetle *Epilachna varivestis* (Coleoptera: Coccinellidae) in Japan. *Applied Entomology and Zoology* **35**(1), 81-85.
- Abrol IP, Palaniappan SP (1988) Green manure crops in irrigated and rainfed lowland rice-based cropping systems in South Asia. 71-82.
- Adachi T, Chiba I, Murata K, Sato H, Narikawa T (1980) The new adzuki bean variety Hokkaishiro. *Bulletin of Hokkaido Prefectural Agricultural Experiment Stations (Hokkaidoritsu Nogyo Shikenjo Shoho)*(43), 80-87.
- Adachi T, Shimada H, Murata K, Kikuchi A, Tabuchi K (1996) Studies on lethal seedlings of Azuki bean [*Vigna angularis*] hybrids. IV. Geographical distribution of lethal complementary gene in Kyushu and Korean local varieties. *Tohoku Journal of Crop Science*(39), 91-92.
- Adinarayananmurthy VV, Rao MVB, Satyanarayana A, Subramanyam D (1993) The crossability of *Vigna mungo* and *Vigna radiata* with *Vigna trilobata*. *International Journal of Tropical Agriculture* **11**(3), 209-213.
- Agarwal HR, Bhati DS (1989) Response to kidney bean (*Vigna aconitifolia* (Jacq.) Marechal) to application of nitrogen, phosphorus and *Rhizobium* inoculation. *Indian Journal of Agronomy* **34**(3), 383-385.
- Agarwal HR, Bhati DS (1995) Effect of weeding schedule on dewgram genotypes. *Farming Systems* **11**(3/4), 43-44.
- Ageeb OAA (1981) Agronomy and crop physiology. 1-18.
- Agrawal NK, Jain BL (1978) *Drechslera subpapendorfii* Mouchacca - a new record from India. *Current Science* **47**(16), 593-594.
- Aguilar HG, Tanigoshi LK, Lumpkin TA (1996) Evaluation of *Vigna angularis* (Willd) Ohwi and Ohashi accessions for resistance to *Tetranychus urticae* Koch (Acari:Tetranychidae). *Experimental & Applied Acarology* **20**(5), 237-247.
- Aher RK, Patil EN, Attarde DR, Solanke AV (1996) Evaluation of rainfed pearl millet based intercropping systems for scarcity zone of Dhule. *Journal of Maharashtra Agricultural Universities* **21**(2), 194-195.
- Ahmad M (1976) Effect of phosphine fumigation on the germination of edible legume seeds. *Journal of Stored Products Research* **12**(3), 211-212.
- Ahmad M, Harwood RF (1973) Studies on a whitefly-transmitted yellow mosaic of urd bean (*Phaseolus mungo*). *Plant Disease Reporter* **57**(9), 800-802.
- Ahmad Z, Bhatti MS, Anwar R (1986) Collection of *Vigna* in Punjab, Pakistan. *Plant Genetic Resources Newsletter*(67), 35-37.
- Ahmed AH, Mills PR (1985) Identification of peanut stunt virus in the Sudan. *Plant Disease* **69**(2), 173-174.
- Ahmed KS, Itino T, Ichikawa T (1999) Effects of plant oils on oviposition preference and larval survivorship of *Callosobruchus chinensis* (Coleoptera: Bruchidae) on azuki bean. *Applied Entomology and Zoology* **34**(4), 547-550.
- Ahn CS (1977) Interspecific hybridization among four species of the genus *Vigna* Savi. **8**(1), 7B.
- Ahn CS, Hartmann RW (1978) interspecific hybridization between rice bean *Vigna-umbellata* (thunb) Ohwi and Ohashi and adzuki bean *V angularis* (willd) Ohwi and Ohashi. *Journal of the American Society for Horticultural Science* **103**(4), 435-438.
- Ahuja AK, Kakkar VK, Gupta BK (2001) Nutritional evaluation of rice bean in buffalo calves. *Indian Journal of Animal Nutrition* **18**(2), 172-175.
- Ahuja DB (1999) Influence of intercropping and mixed cropping sesame with pearl millet, greengram and moth bean on the incidence of insect and mite pests. 183-186.
- Ahuja UR, Shekhawat PS, Dhanpat S, Saraswat PK, Jodha GS (2005) Yield gaps and constraints in the production of mung and moth bean in arid Rajasthan. **2**(2), 279-281.
- Akai J (1981) Studies on the epidemiology and control of sclerotinia disease of beans caused by *Sclerotinia sclerotiorum* (Lib.) De Bary. (36), 83pp.
- Akiyama T, Yamashita R (1993) Study on drying of Azuki beans. *Journal of the Japanese Society of Agricultural Machinery* **55**(6), 129-134.

- Ali SS, Askary TH (2001) Taxonomic status of phytonematodes associated with pulse crops. *Current Nematology* **12**(1/2), 75-84.
- Ali-Khan ST, McVetty P Potential of pulse crops in Manitoba. 29-33.
- Alkic' M, Čolo J, Tahmaz J (2005) Microflora from seed sprouts ready to eat. In. ' pp. 5-14. (Univerzitet u Sarajevu, Poljoprivredni Fakultet: Sarajevo)
- Amarowicz R, Estrella I, Hernández T, Troszyn'ska A (2008) [Antioxidant activity of extract of adzuki bean and its fractions](#). *Journal of Food Lipids* **15**(1), 119-136.
- Amarowicz R, Pegg RB (2006) [Content of proanthocyanidins in selected plant extracts as determined via n-butanol/HCl hydrolysis and a colorimetric assay or by HPLC - a short report](#). *Polish Journal of Food and Nutrition Sciences* **15/56**(3), 319-322.
- Amarowicz R, Troszyn'ska A, Barylko-Picielna N, Shahidi F (2004) [Polyphenolics extracts from legume seeds: correlations between total antioxidant activity, total phenolics content, tannins content and astringency](#). *Journal of Food Lipids* **11**(4), 278-286.
- Ambhore SS, Mundhe PR, Khobragade AM (2008) Influence of sorghum (*Sorghum biocolar* L.) legume associationship on fodder yield. *Journal of Soils and Crops* **18**(2), 379-381.
- Amrit B (2008) [Effect of fermentation on in vitro digestibilities and the level of antinutrients in moth bean \[\*Vigna aconitifolia\* \(Jacq.\) Marechal\]](#). *International Journal of Food Science & Technology* **43**(11), 2090-2094.
- Amrutheswara NG, Andani G, Chikkadevaiah (2005) Effect of sources of organic manure and P-solubilizer on nutrient uptake and yield in rice bean (*Vigna umbellata* Thunb.). *Mysore Journal of Agricultural Sciences* **39**(4), 433-436.
- Amtul W, Bhagwan S (2005) Transfer of technology and dissemination of value added information for agrarian prosperity in arid zone. **2**(2), 241-243.
- Amurtheswara NG, Andani G, Dhanapal GN (2005b) Effect of different organic manures on yield and economics of rice bean (*Vigna umbellata* Thunb.). *Mysore Journal of Agricultural Sciences* **39**(3), 331-334.
- Andrews AC, Manajuti D (1978) The cropping research programme in 1977-78. In. ' pp. 10-12)
- Angelis RCD, Terra ICM, Campos JVM (1986) Effect of dietary lactose on body development study in rats. *Arquivos de Gastroenterologia* **23**(3), 189-195.
- Anil V, Srivastava HP (1988) Studies on vesicular-arbuscular mycorrhiza of moth bean. 1. Assessment of root colonization by *Glomus fasciculatum*. *Indian Phytopathology* **41**(3), 424-427.
- Anju G, Jaiwal PK (1996b) In vitro evaluation of NaCl tolerance in wild and cultivated species of *Vigna*. *National Academy Science Letters* **19**(5/6), 101-106.
- Anju N, Boora P, Khetarpaul N (2000) Effect of domestic processing and cooking methods on the carbohydrate contents of newly released moth bean (*Phaseolus aconitifolius* Jacq.) cultivars. *Nutrition and Health* **14**(4), 265-269.
- Anju N, Boora P, Khetarpaul N (2001) [Effect of microwave cooking on the starch and protein digestibility of some newly released moth bean \(\*Phaseolus aconitifolius\* Jacq.\) cultivars](#). *Journal of Food Composition and Analysis* **14**(5), 541-546.
- Anju N, Boora P, Khetarpaul N (2008) Effect of domestic processing and cooking methods on some antinutritional factors of moth bean. *Journal of Dairying, Foods and Home Sciences* **27**(2), 114-119.
- Anonymous, Makkar HPS, Siddhuraju P, Becker K (2007) Chyrnotrypsin inhibitor. *Methods in Molecular Biology*, 7-9.
- Anoop N, Gupta AK (2003) Transgenic indica rice cv IR-50 over-expressing *Vigna aconitifolia* Δ1-pyrroline-5-carboxylate synthetase cDNA shows tolerance to high salt. *Journal of Plant Biochemistry and Biotechnology* **12**(2), 109-116.
- Anshuman N, Jena SN, Khanda CM, Behera B, Pati P (2009) Effect of genotypes and nitrogen levels on productivity, nutrient uptake and profitability of ricebean *Vigna umbellata* (Thunb.) Ohwi and Ohashi. *Environment and Ecology* **27**(2A), 892-894.
- Antonio H, Neumaier N (1986) Reaction of soil improving plant species to the root-knot nematode *Meloidogyne-Javanica*. *Nematologia Brasileira* **10**, 207-215.
- Antunes AMG, Campos LS (1995) Characterization of amino acids in three genera of the subtribe Phaseoliniae (Leguminosae-Phaseoleae). *Garcia de Orta. Série de Estudos Agronómicos* **19**(1/2), 43-50.
- Anuradha M, Maity BR, Samaddar KR (1994b) Induction of phytoalexin in ricebean (*Vigna umbellata*) by abiotic and biotic elicitors and its modulation by elevated temperature. *Journal of Mycopathological Research* **32**(1), 19-27.
- Aparna K, Naveeda K, Jamuna P (2000) Cooking quality and in vitro digestibility of legumes cooked in different media. *Journal of Food Science and Technology (Mysore)* **37**(2), 169-173.

- Arai K, Takeda H, Kashida Y (1985) Tritium concentration change in the crops by boiling .2. *Journal of Radiation Research* **26**(1), 72-72.
- Arcuri MB, Antunes OAC, Machado SP, Almeida CHF, Oestreicher EG (2004) Stability of immobilized D-hydantoinase from *Vigna angularis* and repeated cycles of highly enantioenriched production of N-carbamoyl-D-phenylglycines. *Amino Acids* **27**(1), 69-74.
- Ariga T, Koshiyama I, Fukushima D (1988) [Antioxidative properties of procyanidins B-1 and B-3 from azuki bean in aqueous systems](#). *Agricultural and Biological Chemistry* **52**(11), 2717-2722.
- Arinathan V, Mohan VR, Britto AJd (2003) [Chemical composition of certain tribal pulses in South India](#). *International Journal of Food Sciences and Nutrition* **54**(3), 209-217.
- Arivudainambi S (2007) Mating behaviour of blue butterfly, *Lampides boeticus* L. *Insect Environment* **13**(3), 115.
- Armstrong RD, McCosker KJ, Millar GR, Walsh K, Johnson S, Probert ME (1997) [Improved nitrogen supply to cereals in Central Queensland following short legume leys](#). *Australian Journal of Experimental Agriculture* **37**(3), 359-368.
- Arora R, Joshi UN, Gupta PP, Singh JV (2009) Effect of Yellow mosaic virus on pathogenesis related enzymes and chlorophyll content in mothbean (*Vigna aconitifolia*). *Acta Phytopathologica et Entomologica Hungarica* **44**(1), 49-60.
- Arora RK (1985) Diversity and collection of wild *Vigna* species in India. *Plant Genetic Resources Newsletter*(63), 26-33.
- Arora RK, Chandel KPS, Joshi BS, Pant KC (1980) Rice bean - tribal pulse of eastern-India. *Economic Botany* **34**(3), 260-263.
- Arora RK, Mehra KL (1978) Exploration in NE India (cereals & pulses). *Plant Genetic Resources Newsletter* (No. 34), 4-8.
- Arora SK, Das B, Paroda RS, Luthra YP (1975) Structural carbohydrates influencing the *in vitro* digestibility of moth (*Phaseolus aconitifolius* Jacq.). *Forage Research* **1**(2), 107-113.
- Arya ID, Sarita A, Rao DV, Shekhawat NS (1990) [Variation amongst protoplast-derived moth bean \*Vigna aconitifolia\* plants](#). *Euphytica* **47**(1), 33-38.
- Arya MPS, Raghuvir S (1995a) Effect of biological staking on the yield of ricebean (*Vigna umbellata*). *Bhartiya Krishi Anusandhan Patrika* **10**(3), 151-155.
- Arya MPS, Raghuvir S (1995b) Effect of biological staking on the yield of ricebean (*Vigna umbellata*). *Bhartiya Krishi Anusandhan Patrika* **10**(3), 151-155. [In Hindi]
- Arya MPS, Raghuvir S (1997) Growth and yield analysis under amaranth (*Amaranthus*)-based intercropping system. *Indian Journal of Agronomy* **42**(3), 432-435.
- Arya MPS, Raghuvir S, Agrawal AK (1997) Growth and yield of ricebean (*Vigna umbellata*) under different dates of sowing and seed rates. *Indian Journal of Agronomy* **42**(1), 127-130.
- Arya MPS, Singh R, Agrawal AK (1997) Growth and yield of ricebean (*Vigna umbellata*) under different dates of sowing and seed rates. *Indian Journal of Agronomy* **42**(1), 127-130.
- Arya MPS, Singh RV (1994b) Response of ricebean (*Vigna umbellata*) to the source and the levels of phosphorus. *Legume Research* **17**(1), 41-46.
- Arya MPS, Singh RV (1996b) Effect of sources and levels of phosphorus on the growth and nodulation behaviour of rice bean (*Vigna umbellata*). *Legume Research* **19**(3/4), 227-229.
- Arya MPS, Singh RV (1996c) Response of rice bean (*Vigna umbellata*) to planting geometry. *Indian Journal of Agronomy* **41**(1), 162-165.
- Asghar A, Malik MA, Nadeem MA, Tahir M, Rashid S (2002) Production potential of mash bean genotypes in response to phosphorus application. *International Journal of Agriculture and Biology* **4**(3), 355-356.
- Asian Vegetable Research and Development C (1987) Status of AVRDC germplasm.
- Asif I, Muhammad A, Nadeem A, Riaz A (2006) Growth and forage yield response of maize-legume mixed cropping to different sowing techniques. *Pakistan Journal of Agricultural Sciences* **43**(3/4), 126-130.
- Atar S, Patel CS, Mahajan V, Singh M, Sarma BK, Azad PK (1991) Maize-ricebean intercropping under mid-altitudes of Meghalaya. *Indian Journal of Hill Farming* **4**(2), 41-43.
- Avenido R, Motoda J, Hattori K (2001b) [Direct shoot regeneration from cotyledonary nodes as a marker for genomic groupings within the Asiatic \*Vigna\* \(subgenus Ceratotropis {Piper} Verdc.\) species](#). *Plant Growth Regulation* **35**(1), 59-67.
- Avenido RA, Hattori K (1999) [Differences in shoot regeneration response from cotyledonary node explants in Asiatic \*Vigna\* species support genomic grouping within subgenus Ceratotropis \(Piper\) Verdc.](#) *Plant Cell Tissue and Organ Culture* **58**(2), 99-110.

- Avenido RA, Hattori K (2000) [Benzyladenine-induced adventitious shoot regeneration from hypocotyls of adzukibean \(\*Vigna angularis\* {Willd.} Ohwi & Ohashi\)](#). *Plant Growth Regulation* **31**(3), 147-153.
- Avenido RA, Power JB, Davey MR (1993) Differential plant regeneration responses from protoplasts of mungbean [*Vigna radiata* (L.) Wilczek] and mothbean (*V. aconitifolia* Jacq Marechal). *Philippine Journal of Crop Science* **18**(3), 175-180.
- Avinash B, Nishi M, Joginder S, Sachendra B, Anil V (2005) Influence of indigenous AM fungi on nutrient uptake in different genotype of moth bean. *2*(2), 255-257.
- Awasthi OP, Singh IS, Bhargava R (2005) Allelopathic influence of aonla (*Emblica officinalis*) leaf extract on germination and seedling growth of seasonal crops. *Range Management and Agroforestry* **26**(2), 120-123.
- Awasthi OP, Singh IS, More TA (2009) Performance of intercrops during establishment phase of aonla (*Emblica officinalis*) orchard. *Indian Journal of Agricultural Sciences* **79**(8), 587-591.
- Babu R, Nanwal RK, Yadav BD, Satyajeet (2007) Biological studies in mothbean [*Vigna aconitifolia*] as influenced by varieties, row spacing and seed rate under rainfed conditions in semi-arid environment. *Annals of Agri Bio Research* **12**(2), 157-160.
- Babu R, Yadav BD, Nanwal RK (2008) Studies on nitrogen and phosphorus fertilization on chlorophyll content and yield in mothbean (*Vigna aconitifolia*) under rainfed conditions in semi-arid environment. *Annals of Biology* **24**(1), 29-31.
- Babu R, Yadav BD, Nanwal RK, Satyajeet (2007) NPK content and uptake studies in mothbean (*Vigna aconitifolia*) as influenced by varieties, row spacing and seed rate under rainfed conditions in semi-arid environment. *Haryana Journal of Agronomy* **23**(1/2), 87-90.
- Badea C, Basu SK (2009) The effect of low temperature on metabolism of membrane lipids in plants and associated gene expression. *Plant Omics* **2**(2), 78-84.
- Baek K, Kim H, Oh H, Yoon B, Kim J, Lee I (2004) Effects of crude oil, oil components, and bioremediation on plant growth. *Journal of Environmental Science and Health. Part A, Toxic/Hazardous Substances & Environmental Engineering* **39**(9), 2465-2472.
- Bagal-Kestwal D, Kestwal RM, Chiang B (2009) Fabrication of photometric dip-strip test systems for detection of  $\beta(1 \rightarrow 3)$ -d-glucan using crude  $\beta(1 \rightarrow 3)$ -d-glucanase from sprouts of *Vigna aconitifolia*. *Biosensors & Bioelectronics* **24**(8), 2566-2573.
- Baik B, Klamczynska B, Czuchajowska Z (1998) Particle size of unsweetened azuki paste as related to cultivar and cooking time. *Journal of Food Science* **63**(2), 322-326.
- Bairwa DK, Sharma JK, Kumawat KC (2006) Efficacy of insecticides, biopesticide and plant products against whitefly, *Bemisia tabaci* on mothbean, *Vigna aconitifolia*. *Indian Journal of Plant Protection* **34**(2), 210-212.
- Bairwa DK, Sharma JK, Kumawat KC (2007) Effect of intercropping on the incidence of sucking insect pests on moth bean [*Vigna aconitifolia* (Jacq.) Marechal]. *Annals of Arid Zone* **46**(2), 213-216.
- Baisakh B (1992) Interrelationship between yield and yield attributes in rice bean (*Vigna-umbellata*). *Indian Journal of Agricultural Sciences* **62**(9), 620-622.
- Bajpai MR, Gaur RG, Singh G (1975) A note on efficacy of vegetative covers and slopes on run off and fertility losses in a semi-arid tract of Rajasthan. *Annals of Arid Zone* **14**(4), 376-378.
- Baker LR, Chen NC, Park HG (1975) Effect of an immunosuppressant on an interspecific cross of the genus *Vigna*. *10*(3), 313.
- Balagtas GE, Ramirez DA (1991b) Genetic variation in Philippine collections of mungbean (*Vigna radiata* (L.) Wilczek), ricebean (*Vigna umbellata* L. Thumb and Ohwi) and cowpea (*Vigna unguiculata* L. Walp.). *Philippine Agriculturist* **74**(1), 103-119.
- Balaji P, Murthy MRN, Sreerama YN, Rao DR, Gowda LR (1997) Studies on simultaneous inhibition of trypsin and chymotrypsin by horsegram Bowman-Birk inhibitor. *Journal of Biosciences* **22**(5), 545-554.
- Baleshwar S, Agarwal PC, Indra R, Dinesh R, Khetarpal RK (2004) Pathogenic fungi detected in indigenous germplasm during 1999-2001. *Indian Journal of Plant Protection* **32**(1), 102-106.
- Bandyopadhyay SK (1988) Variation in host and bacteria in the nodulation of *Phaseolus* species. *Environment and Ecology* **6**(3), 542-546.
- Bangar ND, Amita L, Chavan BH (2009) Genetic divergence in mothbean. *Journal of Maharashtra Agricultural Universities* **34**(1), 10-13.
- Bangar ND, Lakra A, Chavan BH (2008) Correlation and path coefficient analysis in moth bean. *Journal of Maharashtra Agricultural Universities* **33**(2), 164-166.
- Bansal RP, Sen DN (1981) Differential germination behaviour in seeds of the Indian arid zone. *Folia Geobotanica et Phytotaxonomica* **16**(2), 317-330.

- Barik AK, Dhar S, Mukherjee AK (1994) Growth and herbage yield of maize and rice bean grown at pure and intercropping system. *Environment and Ecology* **12**(1), 47-50.
- Basu PS, Biswas C, Majhi R, Datta TK (1996) Serine proteinase from rice bean. *Indian Journal of Biochemistry & Biophysics* **33**(6), 491-497.
- Basu PS, Datta PK, Datta TK (1996) Possible mechanism for the inhibition of lectin-erythrocyte interaction in presence of endogenous lectin receptor. *Bioscience Reports* **16**(6), 453-458.
- Basudev B, Manoranjan M (1997) Effect of nitrogen fertilization and weed management on weed growth, nitrogen uptake and productivity of ricebean. *Annals of Agricultural Research* **18**(2), 214-217.
- Bates GW, Gaynor JJ, Shekhawat NS (1983) Fusion of plant protoplasts by electric fields. *Plant Physiology* **72**(4), 1110-1113.
- Bathar VM, Patel MM, Patel IC, Tikka SBS (2005a) Weed management in mothbean [*Vigna aconitifolia* (Jacq.) Marechal]. In ' pp. 177-178. (Scientific Publishers (India): Jodhpur)
- Bauchan GR (1982) Cytogenetic studies of the dry beans, species of the genera *Phaseolus* and *Vigna* (Leguminosae). *Dissertation Abstracts International, B* **43**(2), 326B.
- Beales N (2004) Review of the microbiological risks associated with sprouted seeds. *CCFRA Review No.41*, 72 pp.
- Bell RW, Edwards DG, Asher CJ (1990) [Growth and nodulation of tropical food legumes in dilute solution culture. Plant and Soil](#) **122**(2), 249-258.
- Benedetti BC, Jorge JT (1987a) Effect of the moisture content on the angle of repose of several grains. *Ciencia e Cultura (Sao Paulo)* **39**(2), 189-192.
- Benedetti BC, Jorge JT (1987b) Moisture content effects on bulk and true density and porosity for some grains. *Engenharia Agricola* **11**, 7-16.
- Beniwal RK (1999) Soil fertility management under pasture and silvipasture systems. 287-290.
- Benjakul S, Visessanguan W, An HJ (2001) [Properties of cysteine proteinase inhibitors from black gram and rice bean. Journal of Food Biochemistry](#) **25**(3), 211-227.
- Benken II, Budanova VI (1976) Trypsin inhibitors in seeds of *Phaseolus* spp. *Trudy po Prikladnoi Botanike, Genetike i Seleksii* **56**(3), 67-73.
- Berg Nvd, Aveling TAS, Venter SL (2002) The evaluation of six fungicides for reducing *Alternaria cassiae* on cowpea seed. *Crop Protection* **21**(6), 501-505.
- Berti F (1985a) Review of research carried out on *Phaseolus* in Zaïre, Rwanda and Burundi from 1945 to the present day. In 'Synthèse de travaux réalisés sur *Phaseolus* aux Zaïre, Rwanda et Burundi de 1945 à nos jours. *Synthèse De Travaux Réalisés Sur Phaseolus Aux Zaire, Rwanda Et Burundi De 1945 a Nos Jours (Synthesis of Work Conducted on Phaseolus in Zaire, Rwanda, and Burundi from 1945 to the Present).* V+54p. Centre D'information Sur Le Haricot Centro Internacional De Agricultura Tropical: Cali, Colombia. Paper, V+54P.)
- Bhadra SK, Hammatt N, Davey MR (1990) Callus induction from seedling protoplasts of *Vigna gracilis* and *V. trilobata*. *SABRAO Journal* **22**(1), 25-33.
- Bhadra SK, Hammatt N, Davey MR (1991a) Tissue and protoplast culture of rice bean (*Vigna umbellata* (Thunb.) Ohwi & Ohashi). *Tropical Agriculture* **68**(4), 345-348.
- Bhag M, Rana RS, Vandana J (1996a) Status of research on new crops in India. 198-204.
- Bhagat PD, Bhati TK, Tomar AS, Prasad RD (2005b) Transfer of technology on moth bean variety RMO-257 and its impact. **2**(1), 132-134. (Scientific Publishers (India): Jodhpur)
- Bhagwan S (1982) Quantitative and quality evaluation of some *Vigna* and *Phaseolus* species. **8**(4), 389.
- Bhagwan S (2006) Attitude of marginal farmers towards moth bean production technology. *Current Agriculture* **30**(1/2), 95-99.
- Bhagwan S, Chauhan TR (2006) Factors influencing the adoption of moth bean production technology in arid zone of Rajasthan. *Journal of Arid Legumes* **3**(1), 34-38.
- Bhagwan S, Gajja BL (2006) Constraints in adoption of new technology in arid zone: a case study of moth bean crop. *Current Agriculture* **30**(1/2), 125-127.
- Bhagwan S, Raj S, Chauhan KNK (2003) Adoption and constraints of moth bean production technology in the arid zone of Rajasthan. In ' pp. 510-514. (Scientific Publishers (India): Jodhpur)
- Bhagwan S, Rana OPS (1985) Variability in protein tryptophan and methionine contents in *Phaseolus* and *Vigna* spp. *Haryana Agricultural University Journal of Research* **15**(1), 15-20.
- Bhandari MM (1990) Correlation analysis in moth bean. *Transactions of Indian Society of Desert Technology*(15), 33-42.
- Bhandari MM (1991a) Association analysis in moth bean (*Vigna aconitifolia* Jacq. Marechal). *International Journal of Tropical Agriculture* **9**(1), 38-41.

- Bhandari MM (1991b) Euclidean cluster analysis in moth bean over two different environments. *International Journal of Tropical Agriculture* **9**(1), 21-24.
- Bhansali RR (2005a) Diseases of moth bean crop under *Prosopis cineraria* agroforestry system. **2**(2), 250-254.
- Bhansali RR (2005b) In quest of seed borne nature of yellow mosaic virus (YMV) in moth bean by using tissue culture technique. **2**(2), 354-356.
- Bhansali RR (2007) Viral biodiversity in arid legume crops. *Journal of Arid Legumes* **4**(2), 102-106.
- Bharat B, Gaur GK (1997) Effect of different feeds on growth of New Zealand white rabbits. *Indian Journal of Animal Research* **31**(1), 61-62.
- Bharat B, Gaur GK, Ahlawat SPS (1998) Genetic parameters of growth in New Zealand white rabbits adjusting different planes of feeds. *Indian Journal of Animal Sciences* **68**(2), 187-188.
- Bharat S, Yadava ND, Beniwal RK, Sanjay K (2001) Farm production losses in arid rainfed condition of Bikaner District. *Current Agriculture* **25**(1/2), 79-83.
- Bhardwaj CL, Paul YS, Singh BM (1985) Some new leguminous hosts of powdery mildew. *Indian Journal of Mycology and Plant Pathology* **15**(3), 292.
- Bhardwaj HL, Hankins A, Mebrahtu T, Mullins J, Rangappa M, Abaye O, Welbaum GE (1996) Alternative crops research in Virginia. 87-96.
- Bhargava MC, Choudhary RK, Yadav SR (2008) Biology and host preference of pulse beetle (*Callosobruchus chinensis* L.) on different pulses. *Journal of Maharashtra Agricultural Universities* **33**(1), 44-46.
- Bhargava PD, Bhatnagar SM (1981) Pulse production in Rajasthan. *Pulse Crops Newsletter* **1**(1), 9-11.
- Bhargava S, Chandra N (1989a) Effect of sodium chloride on callus cultures of moth bean *Vigna aconitifolia* (Jacq.) Marechal cv. IPCMO 926. *Indian Journal of Experimental Biology* **27**(1), 83-84.
- Bhargava S, Chandra N (1989b) Factors affecting regeneration from leaf explants of moth bean, *Vigna aconitifolia* (Jacq.) Marechal. *Indian Journal of Experimental Biology* **27**(1), 55-57.
- Bhargava S, Saxena SN, Gupta D (2004) In vitro flowering in pearl millet, cumin and mothbean. *National Journal of Plant Improvement* **6**(2), 79-81.
- Bhargava S, Upadhyaya S, Garg K, Chandra N (1983) Differentiation of shoot buds in hypocotyl explants and callus cultures of some legumes. 431-433.
- Bhargava SC, Smigocki AC (1994) [Transformation of tropical grain legumes using particle bombardment](#). *Current Science* **66**(6), 439-442.
- Bhargava SC, Chandra N (1989) In vitro regeneration of shoot apices of *Vigna aconitifolia* Jacq. Merechal. *Legume Research* **12**(4), 170-172.
- Bharodia RK, Talati GM (1984) Seasonal incidence of *Tetranychus neocaledonicus* Andre in Gujarat State. *Gujarat Agricultural University Research Journal* **10**(1), 64-65.
- Bhaskar RB, Singh UP, Ahmad ST (1990) Evaluation of promising moth bean cultivars for yield and quality parameters and resistance to yellow mosaic virus. *Range Management & Agroforestry* **11**(1), 41-43.
- Bhat NS, Manjunatha M, Raju GTT (1990) Evaluation of moth bean varieties for resistance to four insect pests. *Current Research - University of Agricultural Sciences (Bangalore)* **19**(2), 30-31.
- Bhati DS (1989) Effect of sowing date on mothbean. *Farming Systems* **5**(1-2), 52-54.
- Bhati DS, Mathur JR, Sharma RC (1987) 'Jwala', a virus-resistant mothbean. *Indian Farming* **37**(4), 7, 14.
- Bhati DS, Mathur JR, Sharma RC (1988) Response of moth bean (*Vigna aconitifolia* (Jacq.) Marechal) to graded levels of N and P. *Annals of Arid Zone* **27**(1), 63-64.
- Bhati TK, Rathore SS, Gaur NK (2005) Effect of improved varieties and nutrient management in kharif legumes under arid ecosystem in Institution Village Linkage Programme. **2**(2), 227-229.
- Bhatia JS, Ghosal AK (1992) Studies on fermentation in the camel (*Camelus dromedarius*). 271-274.
- Bhatt BP, Singh JK (2009) Allelopathic effects of tree species on crops in Eastern Himalaya, India. *Allelopathy Journal* **23**(2), 333-343.
- Bhattacharya N, Mukherjee AK (1998) Herbage growth of ricebean under different dates of sowing and levels of phosphorus. *Forage Research* **24**(2), 121-123.
- Bhatti IH, Riaz A, Nazir MS (2005) Agronomic traits of sesame as affected by grain legumes intercropping and planting patterns. *Pakistan Journal of Agricultural Sciences* **42**(1/2), 56-60.
- Bhavsar VV, Birani SP (1989) Variability, correlation and path analysis in moth bean. *Journal of Maharashtra Agricultural Universities* **14**(2), 148-150.
- Bhavsar VV, Birari SP (1991) Genetic divergence in moth bean, *Vigna aconitifolia* (Jacq.) Marechal. *Biovigyanam* **17**(2), 118-120.
- Bhudev D, Sharma JK, Kumawat KC (2005a) Bio-efficacy of insecticides against sucking insect pests of moth bean, *Vigna aconitifolia*. *Annals of Plant Protection Sciences* **13**(1), 91-93.

- Bhudev D, Sharma JK, Kumawat KC (2005b) Relative susceptibility of moth bean varieties *Vigna aconitifolia* to sucking pests. *Annals of Plant Protection Sciences* **13**(1), 246-247.
- Bhupinder S, Verma MM Introduction and genetic improvement of ricebean (*Vigna umbellata*) as a new pulse crop. 1987, p. 10
- Bhushan B, Gaur GK, Ahlawat SPS (1998) Genetic parameters of growth in New Zealand white rabbits adjusting different planes of feeds. *Indian Journal of Animal Sciences* **68**(2), 187-188.
- Bikash M, Anupam V (1996) Differentiation of natural variants of mungbean yellow mosaic geminivirus by host reactions and DNA-DNA hybridization. *International Journal of Tropical Plant Diseases* **14**(2), 189-202.
- Bishnoi S, Gautam DD (1991) Effect of dairy effluent on seed germination and seedling growth of some crop plants. *International Journal of Ecology and Environmental Sciences* **17**(1), 67-71.
- Bisht IS, Bhat KV, Lakhapaul S, Latha M, Jayan PK, Biswas BK, Singh AK (2005a) [Diversity and genetic resources of wild \*Vigna\* species in India](#). *Genetic Resources and Crop Evolution* **52**(1), 53-68.
- Biswas P, Chowdhury DM, Dana S, Mukherjee N (1983) Resistance of rice bean to *Rhizoctonia bataticola*.. *Indian Phytopathology* **36**(4), 744-745.
- Bora KK, Mathur SR, Makkhan L, Ganesh R (2003) Relative physiological and biochemical tolerance in moth bean cultivars to cadmium stress. *Current Agriculture* **27**(1/2), 81-84.
- Bora KK, Sankhla D, Sankhla N (1979) Pollutants and growth of Indian desert plants. I. Role of fluoride in early seedling growth and enzyme activity. *Transactions of the Indian Society of Desert Technology and University Centre of Desert Studies* **4**(1), 42-46.
- Bora KK, Upadhyaya A, Vaid K, Sankhla D, Sankhla N (1980) Pollutants and growth of Indian desert plants. II. Role of lead in early seedling growth and enzyme activity. *Transactions of Indian Society of Desert Technology and University Centre of Desert Studies* **5**(2), 39-44.
- Borah HK, Debcoudhury PK, Sheikh IA, Barman B (2001b) Genetic parameters correlations and path analysis among yield and yield characters in ricebean (*Vigna umbellata* (Thunb.) Ohwi and Ohasi). *Madras Agricultural Journal* **88**(10/12), 629-632.
- Borah UK (1994) Response of rainfed ricebean (*Vigna umbellata*) genotypes to varying spacings. *Indian Journal of Agronomy* **39**(4), 710-711.
- Branca C, Ricci D, Bassi M (1988) Epidermis integrity and epicotyl growth in azuki bean. *Journal of Plant Growth Regulation* **7**(2), 95-109.
- Bravo L, Siddhuraju P, Saura-Calixto F (1998) [Effect of various processing methods on the \*in vitro\* starch digestibility and resistant starch content of Indian pulses](#). *Journal of Agricultural and Food Chemistry* **46**(11), 4667-4674.
- Bravo L, Siddhuraju P, Saura-Calixto F (1999) [Composition of underexploited Indian pulses. Comparison with common legumes](#). *Food Chemistry* **64**(2), 185-192.
- Brewer IN, Horber E (1984) Evaluating resistance to *Callosobruchus chinensis* Linn. in different seed legumes. 435-443.
- Brockwell J, Andrews JA, et al. (1991) [Erratic nodulation and nitrogen fixation in field-grown pigeonpea \[\*Cajanus cajan\* \(L.\) Millsp.\]](#). *Australian Journal of Experimental Agriculture* **31**(5), 653-661.
- Budanova VI (1982a) New information in the systematics of the genera Phaseolus L. and Vigna Savi. *Trudy po Prikladnoi Botanike, Genetike i Selektii* **72**(1), 16-20.
- Budanova VI, Kozhushko NN, Galiullina EA (1980) Resistance of some Phaseolus species to drought. *Byulleten' Vsesoyuznogo Ordona Lenina i Ordona Druzhby Narodov Instituta Rastenievodstva Imeni N. I. Vavilova*(97), 44-48, 80.
- Budanova VI, Nikitina KV (1972) Effect of bacteria on the sowing quality of Phaseolus seeds. *Byulleten' Vsesoyuznogo Ordona Lenina Instituta Rastenievodstva imeni N.I. Vavilova*(No.23), 36-41.
- Budanova VI, Red'kina ZV (1981) Testa structure in seeds of some Phaseolus species. *Byulleten' Vsesoyuznogo Ordona Lenina i Ordona Druzhby Narodov Instituta Rastenievodstva Imeni N. I. Vavilova*(107), 29-34.
- Buendia-Claveria AM, Chamber M, Ruiz-Sainz JE (1989) A comparative study of the physiological characteristics, plasmid content and symbiotic properties of different *Rhizobium fredii* strains. *Systematic and Applied Microbiology* **12**(2), 203-209.
- Burhan HO (1978) Cotton and associated crops. Leguminous forage crops sequence experiment. 8-11.
- Cabi, No.April (2007) *Xanthomonas axonopodis* pv. *phaseoli*. [Distribution map]. In. ' pp. Map 401 (Edition 4))
- Cabrerasantiago MAI, Intengan CL, et al. (1986) [Protein-requirements of preschool-children consuming rice-milk, rice-toasted mung bean, and rice diets](#). *Qualitas Plantarum-Plant Foods for Human Nutrition* **36**(3), 167-178.

- Cakir S, Koksal F, Tapramaz R, Cakir O (1991) Electron-spin-resonance of free-radicals in some legumes, cereals and their aqueous-solutions under photolysis. *Radiation Physics and Chemistry* **38**(1), 17-21.
- Calkins RH, Huang KR (1979) Soybean production in Taiwan: a farm survey. (11(78-89)), 27pp.
- Campos JE, Whitaker JR, Yip TT, Hutchens TW, Blanco-Labra A (2004) Unusual structural characteristics and complete amino acid sequence of a protease inhibitor from *Phaseolus acutifolius* seeds. *Plant Physiology and Biochemistry* **42**(3), 209-214.
- Campos Ávila J, Fucikovsky Zak L (1981b) Study of some aspects of angular spot caused by *Isariopsis griseola* Sacc. on bean. *Revista Mexicana de Fitopatología* **1**(3), 32-35. [In Spanish]
- Carnovale E, Lugaro E, Marconi E (1991) [Protein quality and antinutritional factors in wild and cultivated species of Vigna spp.](#) *Plant Foods for Human Nutrition* **41**(1) 11-20.
- Carvalheira G, Guerra M (1998) The polytene chromosomes of tapetal cells in the anther of some *Vigna* Savi cultivars and species. *Cytobios* **94**(377), 161-168.
- Cerana R, Bonetti A, Marrè MT, Romani G, Lado P, Marrè E (1983) Effects of a brassinosteroid on growth and electrogenic proton extrusion in azuki bean epicotyls. *Physiologia Plantarum* **59**(1), 23-27.
- Chadha VK, Ranwa NS, Dadheech PK (1998) Synthesis and screening of biological activity of triazolo-thiadiazines and triazolo-thiadiazoles. *Journal of Phytological Research* **11**(2), 201-202.
- Chaghtai SM, Ali Q, Islam S (1978) Net primary productivity and efficiency of *Phaseolus aconitifolius* Jacq. (moth bean) in the cultivated fields of Peshawar, (N.W.F.P.) Pakistan. *Journal of Agricultural Research, Pakistan* **16**(2), 225-240.
- Chaitieng B, Kaga A, Tomooka N, Isemura T, Kuroda Y, Vaughan DA (2006) [Development of a black gram \[Vigna mungo \(L.\) Hepper\] linkage map and its comparison with an azuki bean \[Vigna angularis \(Willd.\) Ohwi and Ohashi\] linkage map.](#) *TAG Theoretical and Applied Genetics* **113**(7), 1261-1269.
- Chakrabarty PK, Shyam KR, Singh BM (1989) Host range studies with three isolates of *Colletotrichum lindemuthianum* causing anthracnose of French bean (*Phaseolus vulgaris* L.). *Indian Journal of Mycology and Plant Pathology* **19**(1), 14-17.
- Chakraborty M, Mukherjee AK (2001) Variation and character association in ricebean (*Vigna umbellata*). *Journal of Research, Birsa Agricultural University* **13**(1), 45-49.
- Chakraborty N, Mandal L (1981) Utilization of rice bean *phaseolus-calcaratus* seeds in chick ration. *Indian Journal of Poultry Science* **16**(4), 415-417.
- Chakraborty T, Mandal S (1999) Performance of different fodder crops under recommended and partially recommended fertilizer doses in lateritic soil of West Bengal. *Indian Agriculturist* **43**(1/2), 55-59.
- Champak B, Nirmala S, Pathak KML (2009) Growth characteristics, economics and hair mineral status of camel calves reared in different systems of management. *Indian Journal of Animal Sciences* **79**(9), 932-935.
- Champak B, Sahani MS (2001) Impact of camel production system on the sustainability of marginal farmers in hot arid villages of Thar desert. *Indian Journal of Animal Research* **35**(1), 10-14.
- Chand P (1975) Preliminary investigations on the antifeeding properties of triphenyltin acetate (Brestan-60) against *Diacrisia obliqua* Walker. *Indian Journal of Entomology* **37**(3), 316-317.
- Chand P (1979) Utilization of food by the Bihar hairy caterpillar *Diacrisia obliqua* Walker. *Indian Journal of Entomology* **41**(3), 237-239.
- Chandel KPS (1981) Wild *Vigna* species in the Himalayas. *Plant Genetic Resources Newsletter*(45), 17-19.
- Chandel KPS, Joshi BS, Arora RK, Pant KC (1978) Rice Bean - a new pulse with high potential. *Indian Farming* **28**(9), 19-22.
- Chandra S (1984) Field observations on plant association of solitarious adult desert locust (*Schistocerca gregaria* Forsk.) in western Rajasthan. *Plant Protection Bulletin, India* **36**(4), 23-28.
- Chang CA (1991) Virus diseases and their transmission of legume crops in Taiwan. 99-110.
- Chang CA (1993) Viruses of legume crops in Taiwan. *Plant Pathology Bulletin* **2**(3), 149-160.
- Chang NT (1988a) Population trends of *Megalurothrips usitatus* (Bagnall) (Thysanoptera: Thripidae) on adzuki bean and soybean examined by four sampling methods. *Plant Protection Bulletin, Taiwan* **30**(3), 289-302.
- Chang NT (1988b) The preference of thrips, *Megalurothrips usitatus* (Bagnall), for three leguminous plants. *Plant Protection Bulletin, Taiwan* **30**(1), 68-77.
- Chang NT (1989) Impacts of soil physical factors on the pupation of bean flower thrips, *Megalurothrips usitatus* (Bagnall) (Thysanoptera: Thripidae). *Plant Protection Bulletin, Taiwan* **31**(4), 377-386.
- Chang NT (1990a) *Ceranisus menes* (Walker) (Eulophidae: Hymenoptera), a new parasite of bean flower thrips, *Megalurothrips usitatus* (Bagnall) (Thripidae: Thysanoptera). *Plant Protection Bulletin (Taipei)* **32**(3), 237-238.

- Chang NT (1990b) Color preference of thrips (Thysanoptera: Thripidae) in the adzuki bean field. *Plant Protection Bulletin, Taiwan* **32**(4), 307-316.
- Chang NT (1992) Dispersion patterns of bean flower thrips, *Megalurothrips usitatus* (Bagnall), (Thysanoptera: Thripidae) on flowers of adzuki bean. *Plant Protection Bulletin (Taipei)* **34**(1), 41-53.
- Chapman KA, Delauney AJ, Kim JH, Verma DPS (1994) Structural organization of de novo purine biosynthesis enzymes in plants: 5-aminoimidazole ribonucleotide carboxylase and 5-aminoimidazole-4-N-succinocarboxamide ribonucleotide synthetase cDNAs from *Vigna aconitifolia*. *Plant Molecular Biology* **24**(2), 389-395.
- Charjan SKU, Tarar JL (1994) Seed quality of moth bean (*Vigna aconitifolia* (Jacq.) Marechal) as influenced by pulse beetle damage during storage. *Journal of Soils and Crops* **4**(1), 22-24.
- Charles-Edwards DA, Lawn RJ (1984) Light interception by grain legume row crops. *Plant, Cell and Environment* **7**(4), 247-251.
- Chatterjee A, Roy A, Ghosh SK (2008) Acquisition, transmission and host range of a begomovirus associated with yellow vein mosaic disease of mesta (*Hibiscus cannabinus* and *Hibiscus sabdariffa*). *Australasian Plant Pathology* **37**(5), 511-519.
- Chatterjee BN, Bhattacharyya KK (1986) 'Principles and practices of grain legume production.' (Oxford & IBH Publishing Co.: New Delhi) 492pp.
- Chatterjee BN, Bhattacharyya KK, Miah AA Growth analysis of rice bean. 1987, p. 249
- Chatterjee BN, Dana S (1977) Rice bean (*Vigna umbellata* (Thumb) Ohwi and Ohashi). *Tropical Grain Legume Bulletin*(No.10), 22-25.
- Chatterjee BN, Mukherjee AK (1979) A new rice bean for fodder and grain pulse. *Indian Farming* **29**(5), 29-31.
- Chau C, Cheung PCK (1999) Effects of the physico-chemical properties of three legume fibers on cholesterol absorption in hamsters. *Nutrition Research* **19**(2), 257-265.
- Chau C, Cheung PCK, Wong Y (1997a) Effects of cooking on content of amino acids and antinutrients in three Chinese indigenous legume seeds. *Journal of the Science of Food and Agriculture* **75**(4), 447-452.
- Chau C, Cheung PCK, Wong Y (1997b) Functional properties of protein concentrates from three Chinese indigenous legume seeds. *Journal of Agricultural and Food Chemistry* **45**(7), 2500-2503.
- Chau C, Cheung PCK, Wong Y (1998a) Hypocholesterolemic effects of protein concentrates from three Chinese indigenous legume seeds. *Journal of Agricultural and Food Chemistry* **46**(9), 3698-3701.
- Chau CF, Cheung PCK, Wong YS (1998b) Chemical composition of three underutilized legume seeds grown in China. *Food Chemistry* **61**(4), 505-509.
- Chaudhari GB, Patil JV, Barhate KK (2000a) Character association in rice bean (*Vigna umbellata* (Thumb.) Ohwi and Ohashi). *Legume Research* **23**(1), 25-28.
- Chaudhari GB, Patil JV, Barhate KK (2000c) Genetic variability in ricebean - *Vigna umbellata*. *Madras Agricultural Journal* **87**(4/6), 348-349.
- Chaudhari GB, Patil JV, Deshmukh RB (1999) Genetic divergence in germplasm of ricebean. *Indian Journal of Pulses Research* **12**(2), 251-253.
- Chaudhary HK (1997) Relative efficiency of line × tester and triple test cross designs for determining genetic architecture of yield and its contributing attributes in adzuki bean (*Vigna angularis* (Willd) Ohwi and Ohashi). *Acta Agronomica Hungarica* **45**(1), 47-55.
- Chaudhary HK, Tashi D, Gupta VP (1992) Genetic analysis for some physiological and seed quality traits in adzuki bean. *Crop Improvement* **19**(1), 48-52.
- Chaudhary HK, Tashi D, Gupta VP (1993a) Genetic architecture of yield traits in adzuki bean (*Vigna angularis* (Willd) Ohwi & Oashi). *Indian Journal of Genetics & Plant Breeding* **53**(3), 247-251.
- Chaudhary HK, Tashi D, Gupta VP (1993b) Inheritance of phenological and morphological traits in adzuki bean. *Crop Improvement* **20**(1), 55-58.
- Chaudhary HK, Tashi D, Gupta VP (1994) Combining ability for phenological and structural traits in adzukibean. *Indian Journal of Pulses Research* **7**(1), 7-10.
- Chaudhary MI, Adu-Gyamfi JJ, Saneoka H, Nguyen Tran N, Suwa R, Kanai S, El-Shemy HA, Lightfoot DA, Fujita K (2008) The effect of phosphorus deficiency on nutrient uptake, nitrogen fixation and photosynthetic rate in mashbean, mungbean and soybean. *Acta Physiologiae Plantarum* **30**(4), 537-544.
- Chaudhary MI, Fujita K (1998a) Comparison of phosphorus deficiency effects on the growth parameters of mashbean, mungbean, and soybean. *Soil Science and Plant Nutrition* **44**(1), 19-30.
- Chaudhary MI, Fujita K (1998b) Responses of dinitrogen fixation and photosynthetic activity in mashbean (*Vigna aconitifolia*) cultivars under low P conditions. *Soil Science and Plant Nutrition* **44**(1), 31-41.

- Chaudhur.Ap, Prasad B (1972) Flowering behavior and yield of rice-bean (*Phaseolus-calcaratus Roxb*) in relation to date of sowing. *Indian Journal of Agricultural Sciences* **42**(7), 627-&.
- Chavan UD, Ajim M, Chavan JK, Ryszard A (2009) Characteristics of starch from rice bean (*Vigna umbellata L.*) seeds - a short report. *Polish Journal of Food and Nutrition Sciences* **59**(1), 25-27.
- Chen BF, Don CT, Chen SS, Li JS, Ma Q (1993) Information on elite cultivars. *Crop Genetic Resources*(2), 48.
- Chen C, Pan L, Hu Y, Hu Z, Ding Y (2006a) Analysis of genetic variation of seed proteins in the genus *Vigna* and among its relatives cultivated in China. *Wuhan University Journal of Natural Sciences* **11**(3), 725-731.
- Chen G, Hsu M, Tan C, Sung H, Kuo CG, Fan M, Chen H, Chen S, Chen C (2005) [Cloning and characterization of a plant defensin VaD1 from azuki bean](#). *Journal of Agricultural and Food Chemistry* **53**(4), 982-988.
- Chen J, Wang S, Jing R, Mao X (2009) Cloning the PvP5CS gene from common bean (*Phaseolus vulgaris*) and its expression patterns under abiotic stresses. *Journal of Plant Physiology* **166**(1), 12-19.
- Chen LB (1993) Disease-resistant adzuki bean variety. *Crop Genetic Resources*(1), 48.
- Chen LB, Wei XH (1992) Evaluation on varietal resistance of *Phaseolus angularis* to *Cercospora canescens*. *Crop Genetic Resources*(1), 28-29.
- Chen NC, Baker LR, Honma S (1983b) [Interspecific crossability among four species of Vigna food legumes](#). *Euphytica* **32**(3), 925-937.
- Chen R, Tsuda S, et al. (2005) [Production of γ-linolenic acid in \*Lotus japonicus\* and \*Vigna angularis\* by expression of the Δ6-fatty-acid desaturase gene isolated from \*Mortierella alpina\*](#). *Plant Science* **169**(3), 599-605.
- Chen THH, Gusta LV, Tjahjadi C, Breene WM (1984) [Electrophoretic characterization of adzuki bean \(\*Vigna angularis\*\) seed proteins](#). *Journal of Agricultural and Food Chemistry* **32**(2), 396-399.
- Chen W, Grau CR, Adeea EA, Meng X (2000) [A molecular marker identifying subspecific populations of the soybean brown stem rot pathogen, \*Phialophora gregata\*](#). *Phytopathology* **90**(8), 875-883.
- Chen W, Gray LE, Grau CR (1996) Molecular differentiation of fungi associated with brown stem rot and detection of *Phialophora gregata* in resistant and susceptible soybean cultivars. *Phytopathology* **86**(10), 1140-1148.
- Chen W, Gray LE, Grau CR (1998) Characterization of a group I intron in the nuclear rDNA differentiating *Phialophora gregata* f.sp. *adzukicola* from *P. gregata* f.sp. *sojae*. *Mycoscience* **39**(3), 279-283.
- Chen WJ, Zhou Y (1991) A preliminary study on evapotranspiration of fields in Nanjing area. *Journal of Nanjing Institute of Meteorology* **14**(3), 325-332.
- Chen X, Srinivas P (2006) A new method to select the drought resistance adzuki bean germplasm. *Agricultural Science & Technology - Hunan* **7**(2), 14-17.
- Chen XZ, Yang J, Hu YC (1993) Preliminary study on the high yield culture techniques of small red bean (*Phaseolus angularis*). *Beijing Agricultural Sciences* **11**(5), 15-17.
- Chen Y, Chen C, Chen W (2003) Population parameters of the silverleaf whitefly (*Bemisia argentifolii* Bellows & Perring) on four legume cultivars. *Formosan Entomologist* **23**(4), 249-261.
- Chen YC (2003) Powdery mildew of vegetable soybean in Taiwan. *Plant Pathology Bulletin* **12**(3), 209-211.
- Cheng H, Shen M, Lu S, Hui E (2002) Application of Fuhuang ointment in treating epidemic parotitis. *Chinese Journal of Information on Traditional Chinese Medicine* **9**(11), 53-54.
- Cheng I, Yang R, Horng S (2003) Interactions between host preference and offspring survivorship of *Callosobruchus maculatus* (Fab.). *Formosan Entomologist* **23**(4), 291-303.
- Cheng J, Wei X (1999) Identification of Phytophthora blight of *Vigna unguiculata* subsp. *cylindrica* and its chemical control. *Acta Phytotaxonomica Sinica* **26**(2), 129-132.
- Cheng T, Pan Y, Guo Z (2009) Agrobacterium-induced hypersensitive necrotic reaction in azuki bean (*Vigna angularis*) epicotyls. *Crop, Environment & Bioinformatics* **6**(1), 15-26.
- Cherng J, Chiang W, Chiang L (2007) [Immunomodulatory activities of edible beans and related constituents from soybean](#). *Food Chemistry* **104**(2), 613-618.
- Cheung PCK, Chau C (1998) [Changes in the dietary fiber \(resistant starch and nonstarch polysaccharides\) content of cooked flours prepared from three Chinese indigenous legume seeds](#). *Journal of Agricultural and Food Chemistry* **46**(1), 262-265.
- Chhabra KS, Kooner BS (1987a) Susceptibility of different stages of Bihar hairy caterpillar, *Diacrisia obliqua* Walker to some insecticides. *Pesticides* **21**(6), 22-24.
- Chhetri DR, Mukherjee AK (2003) Influence of heavy metal toxicity on the metabolism of *Vigna umbellata* T. seeds. *Journal of Hill Research* **16**(1), 22-28.
- Chhetri DR, Supratim M, Safiruddin A (2004b) Physiological and biochemical responses of two rice bean (*Vigna umbellata* T.) cultivars to heavy metal stress. *Environment and Ecology* **22**(1), 27-33.
- Chi Y, Sakamaki Y, Kushigemachi K (2003) The seasonal abundance of the legume pod borer, *Maruca vitrata*, in Kagoshima, Japan. *Memoirs of the Faculty of Agriculture, Kagoshima University* **38**, 41-44.

- Chi Y, Sakamaki Y, Tsuda K, Kusigemati K (2004) Simple artificial diets for mass-reproduction of the legume pod borer, *Maruca vitrata* (Lepidoptera: Crambidae). *Japanese Journal of Applied Entomology and Zoology* **48**(4), 315-320.
- Chiang P, Chen S (2005) Effects of processing conditions on the quality of unsweetened adzuki bean paste. *Taiwanese Journal of Agricultural Chemistry and Food Science* **43**(2), 115-124.
- Chiang P, Chen S, Lin T, Ko W (2005) Effects on process factors for production and qualities of adzuki bean jelly. *Taiwanese Journal of Agricultural Chemistry and Food Science* **43**(4), 241-252.
- Chiba I (1982) Breeding for resistance to brown stem rot in adzuki beans. I. On varietal differences in resistance to BSR. *Bulletin, Hokkaido Prefectural Agricultural Experiment Station*(48), 56-63.
- Chiba I (1985) Breeding for resistance to brown stem rot (BSR) in adzuki beans II. Development of varietal differences. *Bulletin, Hokkaido Prefectural Agricultural Experiment Stations*(52), 79-84.
- Chiba I, Narikawa T, Murata K, Adachi T (1987) Breeding for resistance to brown stem rot (BSR) in adzuki beans. III. The inheritance of resistance to BSR and effect of its introduction. *Bulletin of Hokkaido Prefectural Agricultural Experiment Stations*(56), 1-7.
- Chic T, Tabuchi A, Wakabayashi K, Kamisaka S, Hoson T (2002) Action of xyloglucan hydrolase within the native cell wall architecture and its effect on cell wall extensibility in azuki bean epicotyls. *Plant and Cell Physiology* **43**(1), 21-26.
- Chiu YJ, Messina FJ (1994) Effect of experience on host preference in *Callosobruchus maculatus* (Coleoptera: Bruchidae): variability among populations. *Journal of Insect Behavior* **7**(4), 503-515.
- Cho N, Kang Y, Song C, Kang Y, Cho Y, Ko M (2003a) Effect of seeding dates on the growth characteristics, yield and feed value of whole crop Azuki bean in Jeju Island. *Journal of the Korean Society of Grassland Science* **23**(1), 7-12. [In Korean]
- Cho N, Kang Y, Song C, Kang Y, Cho Y, Ko M (2003b) Effects of phosphate application rate on the growth characteristics, yield and feed value of whole crop Azuki bean in Jeju Island. *Journal of the Korean Society of Grassland Science* **23**(1), 1-6. [In Korean]
- Choe KJ, Park PK, Cho NH, Cho KH, Mun YH (1989) Effect on the cleaning of soybean and adzuki bean by helices width of spiral-type separator. *Research Reports of the Rural Development Administration, Agricultural Engineering and Farm Products Utilization* **31**(1), 68-82. [In Korean]
- Choi W, Ryoo M (2002) Regulation of progeny sex by Anisopteromalus calandrae (Hymenoptera: Pteromalidae) in relation to host preference, host vulnerability and host Size. *Journal of Asia-Pacific Entomology* **5**(2), 193-200.
- Choi YM, Lee SH (1989) Identification of azuki bean mosaic virus on Phaseolus angularis. *Korean Journal of Plant Pathology* **5**(1), 49-53. [In Korean]
- Chopra DP, Mital SP (1979) New plants for semi-arid zone of West Rajasthan. *Indian Farming* **28**(11), 19-21.
- Choubey S, Bhagat RK (1998) Forage productivity in teosinte + ricebean intercropping system. *Journal of Research, Birsa Agricultural University* **10**(2), 167-171.
- Choubey S, Bhagat RK, Prasad NK, Srivastava VC (2000) Economic viability and energetics of teosinte (*Euchlaena maxicana*): ricebean (*Vigna umbellata*) associations under different fertility levels. *Indian Journal of Agricultural Sciences* **70**(1), 28-30.
- Choubey S, Bhagat RK, Prasad NK, Srivastava VC (2001) Soil fertility as influenced by teosinte (*Euchlaena maxicana*): ricebean (*Vigna umbellata*) intercropping system. *Journal of Research, Birsa Agricultural University* **13**(2), 131-136.
- Choubey S, Bhagat RK, Prasad NK, Srivastava VC (2002) Biological compatibility and herbage productivity in teosinte:ricebean intercrop association. *Journal of Research, Birsa Agricultural University* **14**(1), 29-31.
- Choubey S, Bhagat RK, Srivastava VC, Prasad NK (1999) Effect of planting pattern and nutrient management on yield and yield attributing characters of teosinte (*Euchlaena maxicana*) plus ricebean (*Vigna umbellata*) intercropping. *Indian Journal of Agronomy* **44**(3), 525-529.
- Choubey S, Prasad NK, Bhagat RK (1997) Suitable grass-legume combination for higher forage production. *Journal of Research, Birsa Agricultural University* **9**(1), 81-83.
- Chowdhury B (1985) Production of food and fuel in desert environment. **10**(2), 14-22.
- Chowdhury KA, Buth GM (1970) Seed coat structure and anatomy of Indian pulses. *Botanical Journal of the Linnean Society* **63**(Supplement), 169-179.
- Chowdhury RK, Chowdhury JB (1983a) Compatibility between *Vigna radiata* (L.) Wilczek and *Vigna umbellata* (Thunb) Ohwi and Ohashi. *Genetica Agraria* **37**(3/4), 257-266.
- Chowdhury RK, Singh BP (1982) Intercropping of pulses -- a good bet. *Intensive Agriculture* **20**(4), 15-17.

- Chrispeels MJ, Sa MFGd, Higgins TJV (1998) Genetic engineering with  $\alpha$ -amylase inhibitors makes seeds resistant to bruchids. *8*(2), 257-263.
- Christou P (1997) [Biotechnology applied to grain legumes](#). *Field Crops Research* **53**(1/3), 83-97.
- Chun YS, Ryoo MI (1992) Oviposition response of azuki bean weevil (Coleoptera: Bruchidae) to the densities of azuki bean and influence of intraspecific competition on the response. *Korean Journal of Entomology* **22**(2), 73-80.
- Chung WH, Ono Y, Kakishima M (2003) Life cycle of *Uromyces appendiculatus* var. *azukicola* on *Vigna angularis*. *Mycoscience* **44**(6), 425-430.
- Clem RL, Hall TJ (1994) [Persistence and productivity of tropical pasture legumes on three cracking clay soils \(Vertisols\) in north-eastern Queensland](#). *Australian Journal of Experimental Agriculture* **34**(2), 161-171.
- Cohn J, Stokkermans T, Kolli VK, Day RB, Dunlap J, Carlson R, Hughes D, Peters NK, Stacey G (1999) Aberrant nodulation response of *Vigna umbellata* to a *Bradyrhizobium japonicum* NodZ mutant and nodulation signals. *Molecular Plant-Microbe Interactions* **12**(9), 766-773.
- Corvera A, Prome D, Prome JC, Martinez-Romero E, Romero D (1999) The nolL gene from Rhizobium etli determines nodulation efficiency by mediating the acetylation of the fucosyl residue in the nodulation factor. *Molecular Plant-Microbe Interactions* **12**(3), 236-246.
- Cosgrove DJ (1981) Rapid suppression of growth by blue light. Occurrence, time course, and general characteristics. *Plant Physiology* **67**(3), 584-590.
- D'Souza E, Bourke RM, Akus WL (1986) Intensification of subsistence agriculture on the Nembi plateau Papua New Guinea 3. Sweet potato cultivar trials crop rotation trials and crop introductions. *Papua New Guinea Journal of Agriculture Forestry and Fisheries* **34**(1-4), 41-48.
- Dabi RK, Gour HN (1988) Field screening of mothbean (*Vigna aconitifolia*) for susceptibility to insect pests and diseases. *Indian Journal of Agricultural Sciences* **58**(11), 843-844.
- Dadarwal KR, Grover R, Tauro P (1982) Uptake hydrogenase in Rhizobium and nodule leghemoglobin in cow pea miscellany hosts. *Archives of Microbiology* **133**(4), 303-306.
- Dames FJ (1972) A study on the consumption of tapilan, mungo, cowpea, corn and soybean as soilage for swine. *Araneta Research Journal* **19**(3), 206-222.
- Dana S, Karmakar PG (1990a) species relation in Vigna subgenus Ceratotropis and its implication in breeding. *Janick, J. (Ed.). Plant Breeding Reviews, Vol. 8. Xi+388p. Timber Press, Inc.: Portland, Oregon, USA. Illus. Maps*, 19-42.
- Danimihardja S, Lester RN (1974) Comparative studies of some Asiatic species of Phaseolus using electrophoretic analyses of seed proteins. *Annales Bogoriensis* **5**(4), 159-171.
- Das A (2002) Effect of rice bean (*Vigna umbellata* Thunb) supplementation on utilization of mixed jungle grass in goats. *Indian Journal of Animal Nutrition* **19**(1), 47-50.
- Das B, Arora SK, Luthra YP (1974) A note on the effect of phosphorus and potash fertilization on chemical composition and in vitro digestibility of moth forage (*Phaseolus aconitifolius* Jacq.). *Indian Journal of Animal Sciences* **44**(11), 896-897.
- Das KC, Sarkar S, Mukherjee BN, Palchowdhry S, Bhowmik MK (1992) Status of certain macrominerals in cattle in relation to soil and plant of Midnapore district of West Bengal. *Indian Journal of Animal Sciences* **62**(9), 907-909.
- Das M, Chattopadhyay P, Roquib MA (1997b) Correlation and path coefficient analysis of grain yield and its components in rice bean *Vigna umbellata* (Thunb.) Ohwi and Ohashi. *Environment and Ecology* **15**(3), 641-645.
- Das M, Chattopadhyay P, Roquib MA (1997d) Genetic variability in grain yield and its components in rice bean *Vigna umbellata* (Thunb.) Ohwi and Ohashi. *Environment and Ecology* **15**(3), 625-628.
- Das ND, Dana S (1980) Linkage in rice bean. *Indian Journal of Genetics and Plant Breeding* **40**(1), 105-116.
- Das ND, Dana S (1981a) Genetics of some qualitative characters in rice bean *Vigna-umbellata*. *Indian Journal of Genetics and Plant Breeding* **41**(3), 401-405.
- Das ND, Dana S (1981b) Inheritance of seed yield components in rice bean *Phaseolus-calcaratus*. *Indian Journal of Genetics and Plant Breeding* **41**(2), 264-267.
- Das ND, Dana S (1987) [Natural outcrossing in rice bean](#). *Plant Breeding* **98**(1), 68-71.
- Das ND, Dana S (1990) Gene effects in four metric traits of rice bean. *Indian Journal of Genetics & Plant Breeding* **50**(1), 27-32.
- Das PK, Chatterjee BN (1976) Leaf area index, light regime and growth of forage crop. *Forage Research* **2**(2), 165-171.

- Das PK, Chatterjee BN (1977) Competition in grass-legume mixed forage. *Indian Journal of Agricultural Sciences* **47**(1), 47-51.
- Das SK, Bardoloi RK (2008) Productive performances of grower broiler rabbit on roughage feeding. *Indian Veterinary Journal* **85**(4), 101-102.
- Das SK, Das A, Bordoloi RK (2004) Performance of indigenous rabbits of Meghalaya fed on different roughages. *Livestock International* **8**(8), 13-15.
- Das Y, Thakur SR, Singh KP, Sharma SK (2004) Association among growth and yield parameters in ricebean (*Vigna umbellata*) genotypes. *Indian Journal of Plant Physiology* **9**(1), 94-97.
- Dasilveira PM, Dasilva SC, Dasilva OF, Damaceno MA (1994) Agricultural system under irrigation. *Pesquisa Agropecuaria Brasileira* **29**(8), 1243-1252.
- Daspattanayak PS, Basak RK (1995) Efficiency of partially acidulated Purulia rock phosphate prepared with nitric acid as phosphorus source for legume forage crops. *Environment and Ecology* **13**(1), 233-234.
- Datta DK, Mukherjee AK, Samaddar KR (1986a) Biotic and abiotic induction of a fungitoxic compound in rice bean (*Vigna umbellata* (Thunb.) Ohwi and Ohashi). *Journal of Phytopathology* **117**(3), 244-251.
- Datta DK, Samaddar KR (1989) Induction of phytoalexin in rice bean by infection, metabolic inhibitors and irradiation. *Indian Journal of Mycological Research* **27**(1), 7-13.
- Datta KS, Hasija RC, Jai D (1990) Germination and early seedling growth of some kharif crops as affected by salinity. *Haryana Agricultural University Journal of Research* **20**(3), 172-181.
- Datta PK, Basu PS, Datta TK (1988a) Purification of human erythrocytes specific lectins from rice bean, Phaseolus calcaratus syn. *Vigna umbellata*, by high-performance liquid chromatography. *Journal of Chromatography* **431**(1), 37-44.
- Daulay HS, Chakravarty AK, Bhati GN (1972) Intercropping of rainy-season legumes in Cenchrus ciliaris L. and Lasiurus sindicus Henr. pastures in the arid zone of Rajasthan. *Indian Journal of Agricultural Sciences* **42**(2), 148-151.
- Daulay HS, Henry A (1988) Farming system for managing marginal rainfed arid lands. 283-292.
- Daulay HS, Henry A, Bhati TK (2000) Screening of suitable genotypes of cluster-bean, moth bean and green gram in legume based intercropping with pearl millet. *Current Agriculture* **24**(1/2), 85-88.
- Daulay HS, Henry A, Bhati TK, Mathur BK (2006) Compatibility of promising varieties of moth bean and cowpea for inter/mixed cropping in pearl millet in arid regions of western Rajasthan. *Journal of Arid Legumes* **3**(2), 44-48.
- Dave S, Yadav BK, Tarafdar JC (2008) Phytate phosphorus and mineral changes during soaking, boiling and germination of legumes and pearl millet. *Journal of Food Science and Technology (Mysore)* **45**(4), 344-348.
- Davis TD, Gehlot HS, Williams CF, Sankhla N (1987) Comparative shoot growth-retarding activities of paclobutrazol and XE-1019. 121-124.
- Daya R (1999) Monitoring of crop failure pattern for crop planning in kharif season under rainfed conditions. *International Journal of Tropical Agriculture* **17**(1/4), 159-164.
- De Angelis RC, Vecchia MG, Toledo JT, Campos JVM (1980) Peri natal effect of the protein quality in rats. *Arquivos de Gastroenterologia* **17**(1), 40.
- De DN, Susmita K (1997) Genetic resource of Indian phaseoli: distribution, validity of species and endemism. *Indian Journal of Genetics & Plant Breeding* **57**(1), 53-65.
- De LC, Dhiman KR (2001) Effect of leaf manures, potassium and GA3 on growth, flowering and longevity of tuberose. *Journal of Ornamental Horticulture (New Series)* **4**(1), 50-52.
- de Oliveira IMV, Fujimori E, da Silva L (2000) Rice and bean diets increase hepatic and plasmatic activity of gammaglutamyltranspeptidase in growing rats. *Archivos Latinoamericanos De Nutricion* **50**(4), 346-352.
- De R (1981) Agronomic management of crops and cropping systems for arid and semiarid lands. 511-531.
- Deb MP, Joshi PA (2007) Microbiological analysis of sprouts and effect of spices on microbial load. *Journal of Food Science and Technology (Mysore)* **44**(5), 545-547.
- Deepinder K, Kapoor AC (1990a) Some antinutritional factors in rice bean (*Vigna umbellata*): effects of domestic processing and cooking methods. *Food Chemistry* **37**(3), 171-179.
- Deepinder K, Kapoor AC (1990b) Starch and protein digestibility of rice bean (*Vigna umbellata*): effects of domestic processing and cooking methods. *Food Chemistry* **38**(4), 263-272.
- Deepinder K, Kapoor AC (1992) Nutrient composition and antinutritional factors of rice bean (*Vigna umbellata*). *Food Chemistry* **43**(2), 119-124.
- Del Rosario RR, Lozano Y, Pamorasamit S, Noel MG (1981) The trypsin inhibitor activity of legume seeds. *Philippine Agriculturist* **63**(4), 339-344.

- Delauney AJ, Hu CAA, Kishor PBK, Verma DPS (1993) Cloning of ornithine  $\delta$ -aminotransferase cDNA from *Vigna aconitifolia* by trans-complementation in Escherichia coli and regulation of proline biosynthesis. *Journal of Biological Chemistry* **268**(25), 18673-18678.
- Deng D (2002) Jin Hongyuan's recipe for the treatment of peliosis nephritis. *Chinese Journal of Information on Traditional Chinese Medicine* **9**(10), 65.
- Deng Y, Wang J, Zhao Z, Liu H (2003b) Comparative biology of two geographical populations of the southern cowpea weevil, *Callosobruchus chinensis* (L.) (Coleoptera: Bruchidae), in Southwest China. *Entomologia Sinica* **10**(4), 257-264.
- Deokar AB, Bharud RW, Umrani NK (1983) Screening of moth bean genotypes for yellow mosaic disease resistance and yield. *Journal of Maharashtra Agricultural Universities* **8**(2), 198-199.
- Deokar AT, Birari SP, Bhavsar VV (1991) Genetic diversity in respect of grain yield and its components in moth bean (*Vigna aconitifolia* Jacq. Marechal). *Annals of Agricultural Research* **12**(1), 83-87.
- Desai ID, Garcia Tavares ML, Dutrade Oliveira BS, Douglas A, Duarte FAM, Dutrade Oliveira JE (1980) Food-habits and nutritional-status of agricultural migrant workers in southern Brazil. *American Journal of Clinical Nutrition* **33**(3), 702-714.
- Desai ID, Swann MA, Garcia Tavares ML, Dutra De Oliveira BS, Duarte FAM, Dutra De Oliveira JE (1980a) Vitamin E status of agricultural migrant workers in southern Brazil. *Federation Proceedings* **39**(3), Abstract 2756.
- Desai ID, Swann MA, Tavares MLG, De Oliveira BSD, Duarte FAM, De Oliveira JED (1980b) Vitamin E status of agricultural migrant workers in southern Brazil. *American Journal of Clinical Nutrition* **33**(12), 2669-2673.
- Deshpande AN, Kumbhar DD, Daftardar SY (1988) Rock phosphate with green manure and farmyard manure as phosphorus source for rice-bean crop sequence on lateritic soil. *Journal of Maharashtra Agricultural Universities* **13**(1), 10-11.
- Deshpande HW, Wankhede DB, Ghatge PU, Rode AB (2005) Studies on sensory qualities of cake supplemented with native mothbean, horsegram and cowpea flours. *2*(2), 351-353.
- Deshpande HW, Wankhede DB, Syed HM, Shekrapure JL, Bhadarge HH (2003) Studies on sensory and nutritional qualities of breads supplemented with native and germinated horse gram and cowpea flours. In '!' pp. 324-327. (Scientific Publishers (India): Jodhpur)
- Deshpande PK, Bhasin RK (1974) Embryological studies in *Phaseolus aconitifolius* Jacq. Obs. [*Vigna aconitifolia*]. *Botanical Gazette* **135**(2), 104-113.
- Deshwal HL, Sharma JK (1996) Relative susceptibility of different varieties of mothbean against white fly, *Bemisia tabaci* Genn. *Annals of Biology (Ludhiana)* **12**(2), 252-254.
- Desiraju S (1998) Effect of aluminium on growth, lipid peroxidation, superoxide dismutase and peroxidase activities in rice bean and french bean seedlings. *Indian Journal of Plant Physiology* **3**(3), 240-242.
- Desiraju S, Rathore VS (2000) Influence of manganese toxicity on photosynthesis in ricebean (*Vigna umbellata*) seedlings. *Photosynthetica* **38**(3), 449-453.
- Deswal DP, Chand U (1997b) Standardization of the tetrazolium test for viability estimation in ricebean (*Vigna umbellata* (Thunb.) Ohwi & Ohashi) seeds. *Seed Science and Technology* **25**(3), 409-417.
- Deswal DP, Chand U, Dahiya BS (2002a) Standardization of germination test in ricebean (*Vigna umbellata* (Thunb) Ohwi and Ohashi). *Forage Research* **28**(2), 106-109.
- Deswal DP, Chand U, Dahiya BS, Kharb RPS (2002b) Prediction of field emergence through seed vigour and viability tests in ricebean. *Forage Research* **27**(4), 255-261.
- Devi TR, Prodhan HS, Brajendra N, Rohinikumar M (2002a) Effectiveness and efficiency of gamma rays in inducing chlorophyll mutations in M2 generation of three rice bean genotypes (*Vigna umbellata* (L.) Thunb. Ohwi & Ohashi). *Environment and Ecology* **20**(4), 890-893.
- Devi TR, Singh NB (2006) Mutagenic induction of variability and selection in M2 generation of selected ricebean (*Vigna umbellata* (Thunb.) Ohwi and Ohashi) cultivars of Manipur. *Legume Research* **29**(2), 150-153.
- Dhaliwal JS, Tiwana MS (1988) Differential response of cowpea and rice bean varieties to cowpea yellow mosaic virus and insect pests as influenced by different dates of sowing. *Journal of Research Punjab Agricultural University* **25**(3), 446-450.
- Dikshit HK, Jhang T, Singh NK, Koundal KR, Bansal KC, Chandra N, Tickoo JL, Sharma TR (2007) Genetic differentiation of *Vigna* species by RAPD, URP and SSR markers. *Biologia Plantarum* **51**, 451-457.
- Dikshit N, Aghora TS (2004) Diversity in cowpea and some *Vigna* species from Orissa, India. *Journal of Economic and Taxonomic Botany* **28**(1), 1-5.
- Dileep S, Pushpa S, Vimal S, Sunil K (2006) Protease from germinating moth bean seeds. *Indian Journal of Plant Physiology* **11**(1), 100-103.

- Dinca V, Popa G (1984) Some aspects of interspecific hybridization in Phaseolus. *Probleme de Genetica Teoretica si Aplicata* **16**(3), 201-208. [In Romanian]
- Djiwanti SR, Sugawara K, Kondo N, Kobayashi K, Ogoshi A (1999) Effect of soybean cyst nematode infestation on the incidence of brown stem rot in adzuki bean cultivars. *Soil Microorganisms* **53**(1), 19-25.
- Dobhal VK, Gautam NK (1994a) Phenotypic stability in adzuki bean (*Vigna angularis*). *Indian Journal of Agricultural Sciences* **64**(3), 179-181.
- Dobhal VK, Gautam NK (1994b) Stability analysis for yield and component characters in rice-bean (*vigna-umbellata*). *Indian Journal of Agricultural Sciences* **64**(4), 237-239.
- Dobhal VK, Rana JC (1997) Genetic variability and association analysis in adzuki bean (*Vigna angularis*). *Indian Journal of Agricultural Sciences* **67**(4), 171-172.
- Doi K, Kaga A, Tomooka N, Vaughan DA (2002) Molecular phylogeny of genus Vigna subgenus Ceratotropis based on rDNA ITS and atpB-rbcL intergenic spacer of cpDNA sequences. *Genetica* **114**(2), 129-145.
- Doiphode DA, Mungikar AM (2007) Rhizosphere and rhizoplane mycoflora under the influence of deproteinized leaf juice of lucerne. In ' ' pp. 75-88. (Daya Publishing House: Delhi)
- Dong Y, Cao Y (2003) Quality characteristics of germplasm resources of food crops and their utilization. *Scientia Agricultura Sinica* **36**(1), 111-114.
- Dubey SC, Mishra B (1995b) Three new diseases in rice bean (*Vigna umbellata*). *Journal of Research, Birsa Agricultural University* **7**(1), 75-78.
- Dushyent G, Bohra A (1999) Control of Macrophomina phaseolina by root extracts. *Journal of Eco-Physiology* **2**(1), 15-17.
- Dutta M, Phogat BS, Dhillon BS (2007) Genetic improvement and utilization of major under-utilised crops in India. In ' ' (Eds S Ochatt and SM Jain) pp. 251-298)
- Dutta M, Prasad R, Tripathi KC (1993) Increase in seed size of hill-grown legumes. *Indian Journal of Agricultural Sciences* **63**(12), 836-838.
- Dutta R, Deka J, Sarma BN Feasibility of intercropping fodder legumes in direct seeded upland rice. 1996a, pp. 147-149
- Dutta R, Deka J, Sarma BN (1996b) Nutrient uptake and residual soil fertility under upland rice based intercropping systems. *Journal of the Agricultural Science Society of North East India* **9**(1), 87-89.
- Dutta TR, Gupta JN, Gupta SR (1975) Pre- and post-emergence control of Parthenium hysterophorus Linn. *Science and Culture* **42**(3), 179-181.
- Dwivedi GK (1994) Genetic divergence in rice bean for Al-stress in acid soil. *Legume Research* **17**(3/4), 207-212.
- Dwivedi GK (1996a) Tolerance of some crops to soil acidity and response to liming. *Journal of the Indian Society of Soil Science* **44**(4), 736-741.
- Dwivedi GK (1996b) Tolerance of some pulses in acid soil. *Legume Research* **19**(1), 40-46.
- Dwivedi GK (1997) Nodulation behaviour of ricebean (*Vigna umbellata*). *Legume Research* **20**(3/4), 179-184.
- Dwivedi NK (2007) Collection and conservation of genetic diversity in legume crops from arid and semi-arid regions. *Journal of Arid Legumes* **4**(1), 35-39.
- Dwivedi NK, Bhandari DC, Neelam B, Dabas BS, Chandel KPS (1999) Conservation of genetic diversity of arid legumes. 49-56.
- Dwivedi NK, Yadav DP, Pareek BK (2005) Diversity in guar and mothbean in arid and semi-arid regions in India. **2**(2), 325-327.
- Dynia JF, Camargo OA (1998) Effects of liming, green manuring, and phosphate addition on electrochemical attributes of an oxisol from Central Brazil. *Communications in Soil Science and Plant Analysis* **29**(5-6), 755-762.
- Eapen S, George L (1990) Ontogeny of somatic embryos of *Vigna acutifolia*, *Vigna mungo* and *Vigna radiata*. *Annals of Botany* **66**, 219-226.
- Eapen S, Gill R (1986) Regeneration of plants from cultured root explants of mothbean (*Vigna acutifolia* L. Jacq. Marechal). *Theoretical and Applied Genetics* **72**(3), 384-387.
- Eapen S, Köhler F, Gerdemann M, Schieder O (1987) Cultivar dependence of transformation rates in moth bean after co-cultivation of protoplasts with *Agrobacterium tumefaciens*. *Theoretical and Applied Genetics* **75**(1), 207-210.
- Eapen S, Ravinder G, Rao PS (1986) Tissue culture studies in mothbean. *Current Science, India* **55**(15), 707-709.
- Egawa Y (1988b) Phylogenetic differentiation between three Asian *Vigna* species, *V. radiata*, *V. mungo* and *V. umbellata*. *Bulletin of the National Institute of Agrobiological Resources, Japan*(4), 189-200.
- Egawa Y (1991b) Phylogenetic relationships in Asian *Vigna* species. 87-94.

- Egawa Y, Nakagahra M, Fernandez GCJ (1988b) Cytogenetical analysis of tetraploid *Vigna glabrescens* by interspecific hybridization involving diploid Asian *Vigna* species. 200-204.
- El-Shemy HA, Mutasim K, Wakasa K, Ishimoto M (2002) Reproducible transformation in two grain legumes - soybean and azuki bean - using different systems. *7*(2B), 709-719.
- Ellis RH, Lawn RJ, et al. (1994a) [Towards the reliable prediction of time to flowering in 6 annual crops .4. Cultivated and wild mung bean](#). *Experimental Agriculture* **30**(1), 31-43.
- Eusebio PS (1991) Effect of dehulling on the nutritive-value of some leguminous seeds as protein-sources for tiger prawn, penaeus-monodon, juveniles. *Aquaculture* **99**(3-4), 297-308.
- Evans IM, Boulter D (1974) [Chemical methods suitable for screening for protein content and quality in cowpea \(\*Vigna unguiculata\*\) meals](#). *Journal of the Science of Food and Agriculture* **25**(3), 311-322.
- Fageria NK, Baligar VC, Wright RJ (1991) [Influence of phosphate rock sources and rates on rice and common bean production in an oxisol](#). *Plant and Soil* **134**(1), 137-144.
- Fan C, Lee C (2001) Purification of D-hydantoinase from adzuki bean and its immobilization for N-carbamoyl-D-phenylglycine production. *Biochemical Engineering Journal* **8**(2), 157-164.
- Farnham MW, Griffith SM, Miller SS, Vance CP (1990) Aspartate aminotransferase in alfalfa root nodules. III. Genotypic and tissue expression of aspartate aminotransferase in alfalfa and other species. *Plant Physiology* **94**(4), 1634-1640.
- Faroda AS (1972) Effect of seeding rate and row spacings on fodder production of moth bean (*Phaseolus aconitifolius*). *Annals of Arid Zone* **11**(3/4), 183-186.
- Faroda AS, Singh RC (1980) Grow pulses as a catch crop. *Indian Farming* **30**(2), 15-16.
- Feng ZZ, Kobayashi K (2009) Assessing the impacts of current and future concentrations of surface ozone on crop yield with meta-analysis. *Atmospheric Environment* **43**(8), 1510-1519.
- Fery RL (1980) Genetics of *Vigna*. *Horticultural Reviews* **2**, 311-394.
- Fery RL (2002a) New opportunities in *Vigna* Trends in new crops and new uses. Proceedings of the Fifth National Symposium, Atlanta, Georgia, USA, 10-13 November, 2001. In ' (Eds J Janick and A Whipkey) pp. 424-428) (ASHS Press: Alexandria)
- Firake NN, Bangal GB, Gutal GB (1994) Effects of contour planting of matakai with subabul on soil moisture and fodder yield. *Journal of Maharashtra Agricultural Universities* **19**(1), 141-142.
- Flores M CM, Madriz I PM, Warnock de Parra R, Trujillo de Leal A (2005b) Plant height and yield components of six genotypes of the *Vigna* genus in two locations of Venezuela. *Revista de la Facultad de Agronomía, Universidad del Zulia* **22**(4), 351-364.
- Fonseca Marques MF, Bora PS (2000) Chemical composition and analysis of aminoacids from beans. *Ciencia y Tecnología Alimentaria* **2**(5), 248-252. [In Spanish]
- Ford CW (1982b) Accumulation of O-methyl-inositol in water-stressed *Vigna* species. *Phytochemistry* **21**(5), 1149-1151.
- Fox CW (1993) A quantitative genetic analysis of oviposition preference and larval performance on two hosts in the bruchid beetle, *Callosobruchus maculatus*. *Evolution* **47**(1), 166-175.
- Fox CW, Tatar M (1994) Oviposition substrate affects adult mortality, independent of reproduction, in the seed beetle *Callosobruchus maculatus*. *Ecological Entomology* **19**(2), 108-110.
- Frenken G, Hornetz B, Jaetzold R, Willems W (1993) Actual landuse advice in marginal areas of SE-Kenya by atmosphere-ocean-telecommunication. *Tropenlandwirt* **94**(April), 3-12.
- Fujihara S, Abe H, Minakawa Y, Akao S, Yoneyama T (1994) Polyamines in nodules from various plant-microbe symbiotic associations. *Plant and Cell Physiology* **35**(8), 1127-1134.
- Fujihara S, Abe H, Yoneyama T (1995) A new polyamine 4-aminobutylcadaverine: occurrence and biosynthesis in root nodules of adzuki bean plant *Vigna angularis*. *Journal of Biological Chemistry* **270**(17), 9932-9938.
- Fujihara S, Terakado J, Nishibori N (2006) [Accumulation of an aromatic amine, β-phenethylamine, in root nodules of adzuki bean \*Vigna angularis\*](#). *Plant and Soil* **280**(1/2), 229-237.
- Fujihara S, Terakado J, Takenaka M, Yoneyama T (2002) [Specific occurrence of β-phenethylamine in root nodules formed from \*Bradyrhizobium-legume\* symbiosis](#). *Plant and Soil* **238**(1), 123-132.
- Fujii K (1983) Resource dependent stability in an experimental laboratory resource-herbivore-carnivore system. *Researches on Population Ecology*(Supplement 3), 15-26.
- Fujii K (2009) Cause of egg mortality in azuki bean weevils (*Callosobruchus chinensis*): physical or chemical? *Population Ecology* **51**(1), 99-104.
- Fujii K, Miyazaki S (1987a) Infestation resistance of wild legumes (*Vigna sublobata*) to azuki bean weevil, *Callosobruchus chinensis* (L.) (Coleoptera: Bruchidae) and its relationship with cytogenetic classification. *Applied Entomology and Zoology* **22**(2), 229-230.

- Fujii Y (2000) Allelopathy in Vicia, Mucuna and related leguminous species. 191-204.
- Fujii Y, Kageyama D, Hoshizaki S, Ishikawa H, Sasaki T (2001) Transfection of Wolbachia in Lepidoptera: the feminizer of the adzuki bean borer Ostrinia scapulalis causes male killing in the Mediterranean flour moth *Ephestia kuhniella*. *Proceedings of the Royal Society of London. Series B, Biological Sciences* **268**(1469), 855-859.
- Fujine O, Akino S, Kondo N, Naito S (2003) Toxic activities of bud cell culture filtrates of *Fusarium oxysporum* f.sp. *adzukicola* to calluses of adzuki bean cultivars. *Memoirs of the Graduate School of Agriculture, Hokkaido University* **25**(2), 195-202.
- Fujita I, Nishimune A, Watanabe K (1978) Studies on the application of slow-release nitrogen fertilizers to the main crops in the brown andosol of Tokachi district. 1. Effect of slow-release nitrogen fertilizers on the main crops at a standard nitrogen supply. *Research Bulletin of the Hokkaido National Agricultural Experiment Station*(120), 51-74.
- Fujita K, Ofosubudu KG, Ogata S (1992) [Biological nitrogen-fixation in mixed legume-cereal cropping systems](#). *Plant and Soil* **141**(1-2), 155-175.
- Fujita S (2007) Studies on the breeding of Adzuki bean cultivars resistant to Adzuki bean brown stem rot (BSR) and *Phytophthora* stem rot (PSR). (115), 55 pp.
- Fujita S, Kondo N, Shimada H, Murata K, Naito S (2007) Re-evaluation and selection of adzuki beans to breed cultivars resistant to new race of *Phialophora gregata* f. sp. *adzukicola*, the causal agent of adzuki bean brown stem rot (BSR). *Breeding Research* **9**(3), 87-95.
- Fujita S, Murata K, et al. (2002) A new adzuki bean variety "Syumari" with soil-borne disease resistance and excellent processing quality. *Bulletin of Hokkaido Prefectural Agricultural Experiment Stations*(82), 31-40.
- Fujita S, Shimada H, Aoyama S, Murata K, Chiba I, Matsukawa I (2005) A new white Adzuki bean "kita-hotaru" with excellent processing quality and soil-borne disease resistance. *Bulletin of Hokkaido Prefectural Agricultural Experiment Stations*(88), 13-24.
- Fujita S, Shimada H, Murata K, Aoyama S, Chiba I, Matsukawa I, Minami M (2003) A new Dainagon-brand adzuki bean "Toyomi-dainagon" with excellent seed quality and soil-borne disease resistance. *Bulletin of Hokkaido Prefectural Agricultural Experiment Stations*(84), 25-36.
- Fukuda T, Maruyama N, Salleh MRM, Mikami B, Utsumi S (2008) [Characterization and crystallography of recombinant 7S globulins of Adzuki bean and structure-function relationships with 7S globulins of various crops](#). *Journal of Agricultural and Food Chemistry* **56**(11), 4145-4153.
- Fukuda T, Prak K, Fujioka M, Maruyama N, Utsumi S (2007) [Physicochemical properties of native adzuki bean \(\*Vigna angularis\*\) 7S globulin and the molecular cloning of its cDNA isoforms](#). *Journal of Agricultural and Food Chemistry* **55**(9), 3667-3674.
- Fukushima M, Ohashi T, Kojima M, Ohba K, Shimizu H, Sonoyama K, Nakano M (2001) Low density lipoprotein receptor mRNA in rat liver is affected by resistant starch of beans. *Lipids* **36**(2), 129-134.
- Fuller DQ (2007) [Contrasting patterns in crop domestication and domestication rates: recent archaeobotanical insights from the Old World](#). *Annals of Botany* **100**(5), 903-924.
- Gadekar P, Dhumale DB, Raut RS (1990) Interrelationship between yield and its components in rice bean (*Vigna umbellata*). *Indian Journal of Agricultural Sciences* **60**(8), 547-549.
- Gaikwad CB, Patil AJ, Gare BN, Patil JD (1994) Contingent crop planning for delayed onset of monsoon in scarcity area of Maharashtra. *Journal of Maharashtra Agricultural Universities* **19**(2), 226-228.
- Gajja BL, Parihar RP (2003) Economic efficiency of legumes in arid zone of Rajasthan. *Current Agriculture* **27**(1/2), 35-39.
- Gampawar AS, Zinjarde RM, Todekar MG (1992) Comparative study of yield and quality of fodder of forage legumes. *Journal of Soils and Crops* **2**(1), 104-105.
- Gangwar B, Jayan PK (1986) New crop introductions in Andamans. *Indian Farming* **35**(11), 18-21.
- Garg BK (2003) Physiological aspects of abiotic stress tolerance in arid legumes. In '' pp. 347-354. (Scientific Publishers (India): Jodhpur)
- Garg BK (2004) Physiology of drought tolerance of arid zone crops. I. Clusterbean and moth bean. *Current Agriculture* **28**(1/2), 13-20.
- Garg BK (2007) Role of thiourea in improving the productivity of arid legumes. *Journal of Arid Legumes* **4**(1), 1-5.
- Garg BK, Burman U (2007) Influence of thiourea on arid legumes at farmer's fields. *Journal of Arid Legumes* **4**(2), 149-151.
- Garg BK, Burman U, Kathju S (2004) Effect of water stress on moth bean [*Vigna aconitifolia* (Jacq.) Marechal] genotypes. *Indian Journal of Plant Physiology* **9**(1), 29-35.

- Garg BK, Burman U, Kathju S (2005) Comparative water relations, photosynthesis and nitrogen metabolism of arid legumes under water stress. *Journal of Plant Biology* **32**(2), 83-93.
- Garg BK, Gupta IC (1997) Physiology of salt tolerance of arid zone crops. III. Mung bean and moth bean. *Current Agriculture* **21**(1/2), 35-48.
- Garg BK, Kathju S, Burman U (2001) Influence of water stress on water relations, photosynthetic parameters and nitrogen metabolism of moth bean genotypes. *Biologia Plantarum* **44**(2), 289-292.
- Garg BK, Shyam K, Vyas SP, Lahiri AN (1996) Relative effects of sodium salts on growth, yield and metabolism of arid legumes. *Plant Physiology & Biochemistry (New Delhi)* **23**(2), 148-152.
- Garg BK, Uday B (2006) Yield improvement of clusterbean and moth bean by thiourea application under arid conditions. *Journal of Arid Legumes* **3**(1), 51-54.
- Garg BK, Uday B, Shyam K (2004) The influence of phosphorus nutrition on the physiological response of moth bean genotypes to drought. *Journal of Plant Nutrition and Soil Science* **167**(4), 503-508.
- Garg DK, Kakani RK, Deora VS (2004a) Drought tolerance studies in mothbean [*Vigna aconitifolia* (Jacq.) Marechal]. *Journal of Arid Legumes* **1**(1), 20-23.
- Garg DK, Kakani RK, Deora VS (2004b) Genetic variation in drought response of mothbean [*Vigna aconitifolia* (Jacq.) Marechal] genotypes in target environment. *Journal of Arid Legumes* **1**(1), 44-47.
- Garg DK, Kakani RK, Sharma RC, Gupta PC (2003) Genetic variability and character association in moth bean under hyper arid region. In ' pp. 93-97. (Scientific Publishers (India): Jodhpur)
- Ge KL, Wang YS, Yuan XM, Huang PM (1991) Method of regenerating plantlets from mesophyll protoplasts of *Phaseolus angularis*. (US 40977087).
- Ge KL, Wang YZ, Yuan XM, Huang PM, Yang JS, Nie ZP, Testa D, Lee N (1989) Plantlet regeneration from protoplasts isolated from mesophyll cells of adsuki bean (*Phaseolus angularis*, Wight). *Plant Science (Limerick)* **63**(2), 209-216.
- Gehlot HS, Davis TD, Sankhla A, Sankhla D, Upadhyaya A, Sankhla N (1987) Influence of triazole growth regulators and GA3 on in vitro differentiation from moth bean callus. 103-107.
- Gehlot HS, Sunita M, Anila P (2003) Role of specific ions on growth and metabolic patterns in moth bean (C3) and pearl millet (C4) of Indian desert. *Indian Journal of Plant Physiology* **8**(2), 145-149.
- Gehlot HS, Upadhyaya A, Davis TD, Sankhla N (1989) Growth and organogenesis in moth bean callus as affected by paclobutrazol. *Plant and Cell Physiology* **30**(6), 933-936.
- Ghanshyam, Pareek BL (2009) Effect of sulphur and thiourea on growth and yield of mothbean [*Vigna aconitifolia* (Jacq.) Marechal] in arid Western Rajasthan. *Annals of Biology* **25**(1), 17-19.
- Ghatge PU, Wankhede DB, Rodge AB, Deshpande HW (2005) Studies on utilization of mothbean, horsegram and cowpea malt flour in baked products (cookies). **2**(2), 345-347.
- Ghobrial GI (1980) Role of direct-seeded rice in Sudan Gezira crop rotation. *International Rice Research Newsletter* **5**(5), 22.
- Ghosal TK, Dutta S, Senapati SK, Deb DC (2004) Role of phenol contents in legume seeds and its effect on the biology of *Callosobruchus chinensis*. *Annals of Plant Protection Sciences* **12**(2), 442-444.
- Ghosh P, Patra AP, Dhar R, Nayek SS (2006) Growth and productivity performance of different rainy season legumes in the Gangetic plains of West Bengal. *Journal of Crop and Weed* **2**(1), 58-62.
- Ghosh P, Patra AP, Haque R, Nayek SS (2007) Balance sheet of soil nutrients (NPK) as influenced by rainy season legumes and their recycling in wheat based crop sequences. *Indian Agriculturist* **51**(3/4), 111-116.
- Ghosh P, Patra AP, Nayek SS (2005) Studies on performance of different rainy season legumes with respect to their nodulation, NPK content, uptake and residual soil fertility in the Entisol of West Bengal. *Journal of Crop and Weed* **1**(2), 87-91.
- Giga DP, Smith RH (1985) Oviposition markers in *Callosobruchus maculatus* F. and *Callosobruchus rhodesianus* Pic. (Coleoptera, Bruchidae): asymmetry of interspecific responses. *Agriculture, Ecosystems and Environment* **12**(3), 229-233.
- Gill AS, Patil SD, Yadav CL (1984) Intercropping studies in *Cenchrus ciliaris*. *Agricultural Science Digest, India* **4**(4), 206-207.
- Gill AS, Yadav CL (1989) Performance of moth varieties for forage yields under rainfed conditions. *Agricultural Science Digest (Karnal)* **9**(2), 105-106.
- Gill HS, Sharma SR (1980) Blight resistant pulses for hilly areas. *Indian Farming* **30**(2), 23.
- Gill PS, Kanwar S (1980) Performance of cowpea and moth for forage yield under different irrigation levels and dates of planting. *Haryana Agricultural University Journal of Research* **10**(4), 532-535.
- Gill R, Eapen S, Rao PS (1989) Regeneration and differentiation of protoplasts in grain legumes. In ' pp. 99-100. (Kluwer Academic Publishers: Dordrecht)

- Gill R, Mishra KP, Rao PS (1987) [Stimulation of shoot regeneration of \*Vigna aconitifolia\* by electrical control.](#) *Annals of Botany* **60**(4), 399-403.
- Gill RK, Sarjeet S, Brar GS (2008) Genetic diversity studies in ricebean (*Vigna umbellata*) germplasm. *Crop Improvement* **35**(1), 41-44.
- Gillespie DR (1986) A simple rearing method for fungus gnats *Corynoptera* sp. (Diptera: Sciaridae) with notes on life history. *Journal of the Entomological Society of British Columbia* **83**, 45-48.
- Ginzberg I, Stein H, Kapulnik Y, Szabados L, Strizhov N, Schell J, Konez C, Zilberstein A (1998) [Isolation and characterization of two different cDNAs of Δ1-pyrroline-5-carboxylate synthase in alfalfa, transcriptionally induced upon salt stress.](#) *Plant Molecular Biology* **38**(5), 755-764.
- Girase PP, Sonawane PD, Wadile SC (2007) Effect of pearl millet (*Pennisetum glaucum*) based intercropping system on yield and economics of pearl millet on shallow soils under rainfed conditions. *International Journal of Agricultural Sciences* **3**(2), 192-193.
- Giri G, De R (1981) [Short-season fodder legume effects on the grain yield and nitrogen economy of barley under dryland conditions.](#) *Journal of Agricultural Science, UK* **96**(2), 457-461.
- Gleeson D, Lelu-Walter MA, Parkinson M (2005) Overproduction of proline in transgenic hybrid larch (*Larix × leptoleuropaea* (Dengler)) cultures renders them tolerant to cold, salt and frost. *Molecular Breeding* **15**(1), 21-29.
- Godbole DA, Kunachgi MN, Potdar UA, Krishnamurthy KV, Mascarenhas AF (1984) Studies on drought resistant legume: the moth bean, *Vigna aconitifolia* (Jacq.) Marechal. II. Morphogenetic studies. *Plant Cell Reports* **3**(2), 75-78.
- Goel S, Raina SN, Ogihara Y (2002b) Molecular evolution and phylogenetic implications of internal transcribed spacer sequences of nuclear ribosomal DNA in the Phaseolus-Vigna complex. *Molecular Phylogenetics and Evolution* **22**(1), 1-19.
- Gokhale VG, Srivastava BK (1975) Ovipositional behaviour of *Callosobruchus maculatus* (Fabricius) (Coleoptera: Bruchidae)-I. Distribution of eggs and relative ovipositional preference on several leguminous seeds. *Indian Journal of Entomology* **37**(2), 122-128.
- Gomathinayagam P, Ramaswamy NM (1994) Seed, protein pattern of cowpea (*Vigna unguiculata* L) and its distant species. *Journal of Plant Biochemistry and Biotechnology* **3**(2), 149-151.
- Gong H, Xie H, Ma R, Cao M, Wan P, Jin W, Pu S (2008) Genetic diversity assessment of azuki bean by SSR markers. *Journal of Agricultural Biotechnology* **16**(5), 872-880.
- Gonzalvo S, Nieves D, Ly J, Macías M, Carón M, Martínez V (2001) Estimates of nutritive value of Venezuelan feed resources destined for monogastric animals. *Algunos aspectos del valor nutritivo de alimentos venezolanos destinados a animales monogástricos.* **13**(2), 1-7.
- Gopinathan MC, Babu CR (1985) [Structural diversity and its adaptive significance in seeds of \*Vigna-minima\* \(Roxb.\) Ohwi and Ohashi and its allies \(Leguminosae-Papilionoideae\).](#) *Annals of Botany* **56**(6), 723-732.
- Gopinathan MC, Babu CR (1986a) Meiotic studies of the F1 hybrid between rice bean (*Vigna umbellata*) and its wild relative *V. minima*. *Genetica, Netherlands* **71**(2), 115-117.
- Gopinathan MC, Babu CR (1988a) Evaluation utilization and conservation of *Vigna-minima* the wild relative of rice bean. *Acta Universitatis Upsaliensis Symbolae Botanicae Upsalienses* **28**(3), 92-102.
- Gopinathan MC, Babu CR, Chatterjee SR, Abrol YP (1987a) [Nutritional potential of \*Vigna minima\* \(Roxb.\) Ohwi and Ohashi. 1. Seed protein content and amino acid composition.](#) *Qualitas Plantarum: Plant Foods for Human Nutrition* **36**(4), 345-355.
- Gopinathan MC, Babu CR, Chatterjee SR, Abrol YP (1987b) [Nutritional potential of \*Vigna minima\* \(Roxb.\) Ohwi and Ohashi. 2. Seed protein fractions and their amino acid composition.](#) *Qualitas Plantarum: Plant Foods for Human Nutrition* **36**(4), 357-365.
- Gopinathan MC, Babu CR, Shivanna KR (1986a) [Interspecific hybridization between rice bean \(\*Vigna umbellata\*\) and its wild relative \(\*V. minima\*\): fertility-sterility relationships.](#) *Euphytica* **35**(3), 1017-1022.
- Gossrau R (1991) Catalytic histochemistry of acid and neutral hydrolases in plant seedlings. *Histochemical Journal* **23**, 483-489.
- Goto K, Takahashi M, Nishiiri K, Abe K (1985) Effect of overhead flooding on growth and yield of soyabeans and azuki beans. *Research Bulletin of the Hokkaido National Agricultural Experiment Station*(141), 127-145.
- Gour HN (1989) Yellow mosaic virus affects the ion leakage from *Vigna aconitifolia*. *Folia Microbiologica* **34**(2), 132-135.
- Gowda BLV, Puttaswamy (1977) *Phaseolus aconitifolius* Jacq., a new host of *Alcides collaris* Pascoe (Curculionidae: Coleoptera). *Current Research* **6**(4), 68.

- Gowda CLL, Upadhyaya HD, Ghaffar MA (2007) Importance of under-utilized indigenous legumes in Asia-Pacific region. *Proceedings of the 1st International Conference on Indigenous Vegetables and Legumes Prospectus for Fighting Poverty, Hunger and Malnutrition*(752), 95-98.
- Grant G, Edwards JE, Puszta A (1995)  [\$\alpha\$ -Amylase inhibitor levels in seeds generally available in Europe](#). *Journal of the Science of Food and Agriculture* **67**(2), 235-238.
- Grant G, More LJ, McKenzie NH, Dorward PM, Buchan WC, Telek L, Puszta A (1995) [Nutritional and hemagglutination properties of several tropical seeds](#). *Journal of Agricultural Science* **124**, 437-445.
- Gray LE, Hepburn AG (1992) Mitochondrial DNA restriction patterns of *Phialophora gregata* isolates from soybean and adzuki bean. *Phytopathology* **82**(2), 211-215.
- Guggari AK, Biradar BM, Kalaghatali SB (2003) Studies on productivity potential and economics of castor based intercropping systems under dryland conditions. *Research on Crops* **4**(3), 331-335.
- Gulati A, Jaiwal PK (1996) In vitro evaluation of NaCl tolerance in wild and cultivated species of Vigna. *National Academy Science Letters-India* **19**(5-6), 101-106.
- Gulati IJ, Yadav SR, Singh SP (2005b) Soil test based fertilizer recommendation for mothbean in Torripsamments of Rajasthan. **2**(1), 135-138. (Scientific Publishers (India): Jodhpur)
- Gulnaz M, Shahnaz D (2003) Biological control of root-infecting fungi with *Trichoderma harzianum*. *Pakistan Journal of Botany* **35**(5), 971-975.
- Gunn CR (1973) Recent nomenclatural changes in *Phaseolus* L. and *Vigna* Savi. *Crop Science* **13**(4), 496.
- Gupta AP, Chahal RS, Shukla UC (1979) Performance of various intercrops in pigeon pea. *Haryana Agricultural University Journal of Research* **9**(2), 158-159.
- Gupta BM, Dabi RK, Gupta HC, Sharma SK (1979) Effect of host plants on the larval development of *Diacrisia obliqua* Walker. *Journal of Entomological Research* **3**(2), 224-226.
- Gupta BM, Yadava CPS (1985) Effect of BHC dust on germination of groundnut and mothbean. *Indian Journal of Agricultural Sciences* **55**(12), 754-755.
- Gupta BN, Singh RB, Chatterjee D (1981) CHEMICAL-COMPOSITION AND NUTRITIVE-VALUE OF RICE BEAN (*PHASEOLUS-CALCARATUS ROXB*) HAY. *Indian Veterinary Journal* **58**(9), 727-730.
- Gupta CR, Awasthi OP, Shukla N (1999) Intercropping in cashew orchard under rainfed condition - a model for Bastar plateau zone of Madhya Pradesh. *Cashew* **13**(2), 18-22.
- Gupta CS, Samaddar KR (1997) Interaction between soil organisms and Rhizobium. *Journal of Mycopathological Research* **35**(2), 137-142.
- Gupta GN, Singh G, Kachwaha GR (1998) Performance of *Prosopis cineraria* and associated crops under varying spacing regimes in the arid zone of India. *Agroforestry Systems* **40**(2), 149-157.
- Gupta HK, Yadav BPS, Gupta JJ, Bujarbarua KM (1993) Utilization of leguminous roughages in rabbit rations. *Indian Journal of Animal Sciences* **63**(4), 481-483.
- Gupta HK, Yadav BPS, Gupta JJ, Bujarbarua KM (1998) Influence of feeding green ricebean (*Vigna umbellata*, Thunb) on growth and feed conversion efficiency in meat rabbits in the Eastern Himalayan Region of India. *Proceedings of the 6th World Rabbit Congress, Vols 1-3*, A195-A197.
- Gupta HS (1987) [A rapid staining technique for staging of microspores in rice](#) (*Oryza-sativa-L*) and rice bean (*Vigna-umbellata*). *Current Science* **56**(20), 1072-1073.
- Gupta JJ (2002) Nonconventional feed resources for poultry production in Arunachal Pradesh. In ' Ed. KA Singh) pp. 139-146
- Gupta JJ, Bujarbarua KM (2006) Organic production of leguminous fodders and their feeding value in rabbits. *Indian Journal of Animal Sciences* **76**(2), 151-153.
- Gupta JJ, Yadav BPS, Gupta HK (1992b) Rice bean (*Vigna umbellata*) as poultry feed. *Indian Journal of Animal Nutrition* **9**(1), 59-62.
- Gupta JP, Gupta GN (1983) Effect of grass mulching on growth and yield of legumes. *Agricultural Water Management* **6**(4), 375-383.
- Gupta NK, Rao KVP, Rathore MS (1989) Performance of forage crops in Manipur. *Indian Journal of Hill Farming* **2**(1), 63-64.
- Gupta P, Mukherjee D (1981) Biochemical studies during seedling growth of *Phaseolus aconitifolius* L. *Seed Research* **9**(2), 137-143.
- Gupta RC, Dixit BS, Banerji R (1995) Oils of some non-conventional legume seeds. *Journal of the Oil Technologists' Association of India* **27**(4), 239, 241.
- Gupta S, Kozak M, Sahay G, Durrai AA, Mitra J, Verma MR, Pattanayak A, Thongbam PD, Das A (2009) [Genetic parameters of selection and stability and identification of divergent parents for hybridization in rice bean \(\*Vigna umbellata\* Thunb. \(Ohwi and Ohashi\)\) in India](#). *Journal of Agricultural Science* **147**(5), 581-588.

- Gupta SK, Gopalakrishna T (2009) [Genetic diversity analysis in blackgram \(\*Vigna mungo\* \(L.\) Hepper\) using AFLP and transferable microsatellite markers from azuki bean \(\*Vigna angularis\* \(Willd.\) Ohwi & Ohashi\).](#)  
*Genome* **52**(2), 120-129.
- Gupta SN, Verma PK, Tyagi CS (2002) Variability among late maturing vis-a-vis extra early maturing genotypes of ricebean. *Annals of Biology* **18**(1), 99-100.
- Gupta SR, Gupta JN Survey, collection and maintenance of germplasm. 30-31.
- Gupta US (1984) Production potential of grain legumes. In. ' pp. 319-365)
- Gupta VP, Plaha P, Rathore PK (2002) [Partly fertile interspecific hybrid between a black gram × green gram derivative and an adzuki bean.](#) *Plant Breeding* **121**(2), 182-183.
- Gupta VP, Sharma VK, Sharma JD, Rathore PK (1994) Genetic divergence in adzuki bean. *Crop Improvement* **21**(1/2), 49-53.
- Gutierrez Bd, Pulido F J (1978) Life-cycle and habits of *Anticarsia gemmatalis*, a pest of soyabean in the Cauca Valley. *Ciclo de vida y habitos de Anticarsia gemmatalis plaga de la soya en el Valle del Cauca.* **4**(1/2), 3-9.
- Gyanendar S, Major S, Dhiman KR (1994) Rice bean: a promising pulse for Sikkim. *Indian Farming* **44**(1), 30.
- Gyanendra S, Chaudhary BS, Singh SP (1998) Stability analysis of some agro-morphological characters in ricebean. *Annals of Agricultural Research* **19**(4), 411-414.
- Gyanendra S, Major S (1996) Combining ability for yield and its components in rice bean. *Indian Journal of Genetics & Plant Breeding* **56**(4), 520-525.
- Gyanendra S, Major S, Dhiman KR (1992) Correlation and path analysis in ricebean under mid altitude conditions. *Crop Improvement* **19**(2), 152-154.
- Gyanendra S, Singh SP, Chaudhary BS (1998) Genetic divergence analysis in ricebean. *Indian Journal of Genetics & Plant Breeding* **58**(1), 101-105.
- Gyanendra S, Singh SP, Chaudhary BS (2001) Genetic variability for yield and its components in rice bean. *Annals of Agricultural Research* **22**(4), 476-480.
- Hachiya K (1989) Distribution of the flower thrips, *Frankliniella intonsa* (Trybom) among flowers, leaves, vegetative points and others on adzuki bean [*Vigna angularis*]. *Annual Report of the Society of Plant Protection of North Japan*(40), 112-114.
- Hachiya K (1990a) Effect of injury by the flower thrips, *Frankliniella intonsa* Trybom, on pod formation and yield of adzuki bean *Phaseolus angularis* L. [*Vigna angularis*]. *Japanese Journal of Applied Entomology and Zoology* **34**(4), 326-329.
- Hachiya K (1990b) Occurrence of thrips on adzuki bean. *Bulletin of Hokkaido Prefectural Agricultural Experiment Stations*(61), 1-6.
- Hacker JB, Williams RJ, Pengelly BC (1996) A characterisation study of the genus *Vigna* with regard to potential as forage. (22), 32 pp.
- Hadiuzzaman S, Miah MAK (1989) Invitro organogenesis in rice bean (*Vigna-umbellata* L.). *Bangladesh Journal of Botany* **18**(2), 157-162.
- Hamada S, Ezaki S, Hayashi K, Toko K, Yamafuji K (1992) Electric current precedes emergence of a lateral root in higher plants. *Plant Physiology* **100**(2), 614-619.
- Hamana K, Niitsu M, Samejima K (1996) Further polyamine analyses of leguminous seeds and seedlings: the occurrence of novel linear, tertiary branched and quaternary branched pentaamines. *Canadian Journal of Botany* **74**(11), 1766-1772.
- Hammer K, Won DC, Han UX (1989) Report on a mission for the collection of plant genetic resources in the Democratic People's Republic of Korea, 1988. *Kulturpflanze* **37**, 391-399.
- Han ED, An JH (1990) Studies on ecology and control of the azuki bean weevil *Callosobruchus-chinensis* l. In Korea. *Research Reports of the Rural Development Administration (Suweon)* **32**(1 Crop Prot), 24-59.
- Han K, Fukushima M, Kato T, Kojima M, Ohba K, Shimada K, Sekikawa M, Nakano M (2003a) Enzyme-resistant fractions of beans lowered serum cholesterol and increased sterol excretions and hepatic mRNA levels in rats. *Lipids* **38**(9), 919-924.
- Han K, Fukushima M, Ohba K, Shimada K, Sekikawa M, Chiji H, Lee C, Nakano M (2004) Hepatoprotective effects of the water extract from adzuki bean hulls on acetaminophen-induced damage in rat liver. *Journal of Nutritional Science and Vitaminology* **50**(5), 380-383.
- Han K, Fukushima M, Shimizu K, Kojima M, Ohba K, Tanaka A, Shimada K, Sekikawa M, Nakano M (2003b) Resistant starches of beans reduce the serum cholesterol concentration in rats. *Journal of Nutritional Science and Vitaminology* **49**(4), 281-286.

- Han K, Iijuka M, Shimada K, Sekikawa M, Kuramochi K, Ohba K, Ruvini L, Chiji H, Fukushima M (2005) Adzuki resistant starch lowered serum cholesterol and hepatic 3-hydroxy-3-methylglutaryl-CoA mRNA levels and increased hepatic LDL-receptor and cholesterol 7 $\alpha$ -hydroxylase mRNA levels in rats fed a cholesterol diet. *British Journal of Nutrition* **94**(6), 902-908.
- Han K, Lee D (1995) Identification and etiological characteristics of anthracnose fungi isolated from soyabean, adzuki beans and mung beans. *Korean Journal of Plant Pathology* **11**(1), 30-38. [In Korean]
- Han OK, Kaga A, Isemura T, Wang XW, Tomooka N, Vaughan DA (2005) [A genetic linkage map for azuki bean \[\*Vigna angularis\* \(Willd.\) Ohwi & Ohashi\]](#). *TAG Theoretical and Applied Genetics* **111**(7), 1278-1287.
- Han T, Wu C, Mentreddy RS, Zhao J, Xu X, Gai J (2005) Post-flowering photoperiod effects on reproductive development and agronomic traits of long-day and short-day crops. *Journal of Agronomy and Crop Science* **191**(4), 255-262.
- Hanafy MS, Rahman SM, Khalafalla MM, El-Shemy HA, Nakamoto Y, Ishimoto M, Wakasa K (2006) [Accumulation of free tryptophan in azuki bean \(\*Vigna angularis\*\) induced by expression of a gene \(OASA1D\) for a modified  \$\alpha\$ -subunit of rice anthranilate synthase](#). *Plant Science* **171**(6), 670-676.
- Hang AN, McClary DC, Gilliland GC, Lumpkin TA (1993) Plant configuration and population effects on yield of azuki bean in Washington State. 588-590.
- Haq N (1983) New food legume crops for the tropics. In: ' pp. 144-160)
- Hara H (1993) Rivellia basilaris (Wiedemann) (Diptera, Platystomatidae) and its allied species in East Asia I. *Japanese Journal of Entomology* **61**(4), 819-831.
- Harada G, Kondo N (2009) Adzuki bean leaf infection by Phytophthora vignae f. sp. adzukicola and resistance evaluation using detached leaves inoculated with zoospores. *Journal of General Plant Pathology* **75**(1), 52-55.
- Harano T, Fujisawa M, Miyatake T (2006) Effect of oviposition substrate on female remating in Callosobruchus chinensis (Coleoptera: Bruchidae). *Applied Entomology and Zoology* **41**(4), 569-572.
- Harano T, Yasui Y, Miyatake T (2006) Direct effects of polyandry on female fitness in Callosobruchus chinensis. *Animal Behaviour* **71**(3), 539-548.
- Hardman LL, Vance CP, Groat RG (1980) Growth, nitrogen fixation and nodule development in adzuki bean (*Vigna angularis*). 82.
- Harpal S, Mishra D, Kushwaha HL (2005a) Development of tractor drawn three furrows (six rows) multi-crop seed-cum-fertilizer drill for arid legumes. **2**(2), 372-374.
- Harpal S, Mishra D, Nahar NM, Kushwaha HL (2005b) Energy use pattern in cultivation of legumes in arid region. **2**(2), 365-371.
- Harris D, Fry GJ, Miller ST (1994) Microtopography and agriculture in semi-arid Botswana. 2. Moisture availability, fertility and crop performance. *Agricultural Water Management* **26**(1+2), 133-148.
- Harrison HF, Jr. (1988) Metribuzin tolerance within the genus Vigna. *Weed Technology* **2**(1), 59-63.
- Hartman GL, Huang YH, Li S (2004) [Phytotoxicity of \*Fusarium solani\* culture filtrates from soybeans and other hosts assayed by stem cuttings](#). *Australasian Plant Pathology* **33**(1), 9-15.
- Harvey PR, Warren RA, Wakelin S (2008) The Pythium-Fusarium root disease complex - an emerging constraint to irrigated maize in southern New South Wales. **48**(3), 367-374.
- Hasan N, Jain RK (1984) Screening mothbean (*Vigna aconitifolius* Jacq) germplasm for resistance to root-knot nematode (*Meloidogyne incognita*). *Agricultural Science Digest, India* **4**(4), 198-199.
- Hashimoto T (1996) Effect of continuous cropping on autolysis period of darkness seedling test. *Research Bulletin of the Hokkaido National Agricultural Experiment Station*(164), 1-6.
- Hataya T, Saito M, Sumino A, Abe H, Tsuchiya S (2001) Effect of the mixture of kasugamycin and copper oxychloride on bacterial disease of adzuki bean caused by *Pseudomonas syringae* pv. *syringae* van Hall. Society of Plant Protection of North Japan, Akita.
- Hattori T, Ohta Y (1985) Induction of phenylalanine ammonia-lyase activation and isoflavone glucoside accumulation in suspension-cultured cells of red bean, *Vigna angularis*, by phytoalexin elicitors, vanadate, and elevation of medium pH. *Plant and Cell Physiology* **26**(6), 1101-1110.
- Hazra CR, Pradeep B (1993) Effect of legume intercropping in rainfed pearl millet on forage yield, microclimate and soil fertility. *Range Management & Agroforestry* **14**(2), 125-130.
- He N, Sawada S, Kato K, Kojima M (2006) [The growth of leaf surface in the initial growth stage of azuki bean in cool weather regions](#). *Japanese Journal of Crop Science* **75**(2), 197-203.
- He TY (1992) Extra-large adzuki bean germplasm resource Xian Feng Xiao Dou. *Crop Genetic Resources*(3), 48.
- He WX (1987) Identification of nutritional quality in grain legumes. *Zuowu Pinzhong Ziyuan*(1), 13-17.
- He WX (1991) Rapid determination of amylose in grain legumes. *Crop Genetic Resources*(2), 33-35.

- Hegde U, Tilak VD (1981) Anatomy of the seedling in the Leguminosae-II. *Indian Journal of Botany* **4**(2), 96-102.
- Heintel S, Wanek W, Richter A (2002) Carbon isotope signatures of leaf carbon fractions: a new approach for studying short-term water deficits of plants. In 'C&S Papers Series No.11/P.' pp. 243-244. (International Atomic Energy Agency (IAEA): Vienna)
- Henry A (2001) Genotype × environment interaction for seed yield in mothbean germplasm. *Current Agriculture* **25**(1/2), 85-89.
- Henry A, Daulay HS (1983a) Genotype X environment interactions for seed yield in mothbean. *Indian Journal of Agricultural Sciences* **53**(6), 469-470.
- Henry A, Daulay HS (1983b) Genotype X environment interactions of induced mutants in mothbean. *Madras Agricultural Journal* **70**(11), 705-708.
- Henry A, Daulay HS (1983c) Performance of induced mutants in moth bean. *Indian Journal of Genetics and Plant Breeding* **43**(3), 342-344.
- Henry A, Daulay HS Induced mutants in mothbean and clusterbean. 1987, Ludhiana, pp. 109-110
- Henry A, Daulay HS (1990) Induced dark brown mutant in mothbean. *Madras Agricultural Journal* **77**(9-12), 501-504.
- Henry A, Kackar NL (2002a) Adaptability and relative performance of moth bean varieties. *Current Agriculture* **26**(1/2), 67-69.
- Henry A, Kackar NL (2002b) Phenotypic stability in moth bean. *Current Agriculture* **26**(1/2), 71-73.
- Henry A, Krishna GVSR (1986) Genetic divergence in mothbean. *Indian Journal of Agricultural Sciences* **56**(4), 298-299.
- Henry A, Kumar D National Symposium on "Arid legumes for sustainable agriculture and trade", Central Arid Zone Research Institute, Jodhpur, India, 5-7 November 2004. Volume 1. 2005b, Jodhpur, p. iv + 200 pp. (Scientific Publishers (India): Jodhpur)
- Henry A, Kumar D National Symposium on "Arid legumes for sustainable agriculture and trade", Central Arid Zone Research Institute, Jodhpur, India, 5-7 November 2004. Volume 2. 2005c, Jodhpur, pp. iv + 201-424. (Scientific Publishers (India): Jodhpur)
- Heo N, Kang M, Ha K, Kim H, Choi J (1997) Identification of virus from azuki bean plant. *Korean Journal of Crop Science* **42**(2), 160-165. [In Korean]
- Hernandez X E, Ramos R A, Martinez A MA (1979) Ethnobotany. 113-138. [In Spanish]
- Hertel K, Roberts K (2003) 'Insect and mite control in field crops 2003.' (NSW Agriculture: Orange) vi + 90 pp.
- Hertel K, Roberts K (2005) 'Insect and mite control in field crops 2005.' (NSW Department of Primary Industries: Orange) vi + 86 pp.
- Hertel K, Woodlands K (2001) 'Insect and mite control in field crops 2001.' (NSW Agriculture: Orange) vi. + 82 pp.
- Higashida S, Kitagawa I (2002) Application of subsoiler filled with bark manure to wet Andosols in the Tokachi district. *Bulletin of Hokkaido Prefectural Agricultural Experiment Stations*(83), 51-54.
- Hira CK, Kanwar JK, Gupta N, Kochhar A (1988) Cooking quality and nutritional-evaluation of the rice-bean (*Vigna-umbellata*). *Journal of Food Science and Technology-Mysore* **25**(3), 133-136.
- Hirano H, Kagawa H, Okubo K (1992) Characterization of proteins released from legume seeds in hot water. *Phytochemistry* **31**(3), 731-735.
- Hirasawa E, Yoshida T, Yoshida I (1994) Effects of wounding, ethylene and abscisic acid on the development of α-amylase activity in detached cotyledons of *Pisum sativum*. *Journal of Plant Physiology* **143**(3), 359-362.
- Hitokoto H, Morozumi S, Wauke T, Sakai S, Kurata H (1981) Fungal contamination and mycotoxin-producing potential of dried beans. *Mycopathologia* **73**(1), 33-38.
- Hiwale SS, Kakade OK, Gohil DI, Bagle BG, Dhandhar DG (2007) Allelopathic influence of hortisilvicultural tree species on arable crops under semi arid conditions. *Indian Journal of Agroforestry* **9**(1), 23-27.
- Hiwatari M, Tsunoda T, Yashima Y, Maekawa M, Niizeki M, Kita F (1989) Applied cropping system and yields of five crops grown in the field for farm practice from 1967 to 1988. *Research Bulletin of the University Farm, Hokkaido University* **26**, 65-70.
- Hodgson AS, Holland JF, Rayner P (1989) Effects of field slope and duration of furrow irrigation on growth and yield of six grain-legumes on a waterlogging-prone vertisol. *Field Crops Research* **22**(3), 165-180.
- Hollington PA, Joshi KD, Mueller RAE, Andersen P, Yadavendra JP, Kumar N, Neog SB, Bajracharya J, Shrestha PK Ricebean: food security through popularisation of an underutilised grain legume in India and Nepal. 2008, pp. 460-461
- Holloway GJ (1986) The potency and effect of phytotoxins within yellow split-pea (*Pisum sativum*) and adzuki bean (*Vigna angularis*) on survival and reproductive potential of *Sitophilus oryzae* (L.) (Coleoptera: Curculionidae). *Bulletin of Entomological Research* **76**(2), 287-295.

- Holloway GJ, Mackness MI (1988) The evolutionary adaptation of enzyme systems in *Sitophilus oryzae* fed on toxic legumes. *Entomologia Experimentalis et Applicata* **48**(2), 165-172.
- Holmes JHG, Dixon RM, Smith C, Entonu S, Chau H, Sanpote S, Acharyya B (1993) Resistance of trypsin inhibitors to fermentation by rumen microflora. 183-186.
- Honda H, Ohsawa K (1990) Chemical ecology for stored products insects. *Journal of Pesticide Science* **15**(2), 263-270.
- Honda H, Yamamoto I, Sakai A (1989) Analysis of the oviposition preference from the egg-distribution in *Callosobruchus chinensis* L. *Journal of Agricultural Science, Tokyo Nogyo Daigaku* **33**(3), 239-245.
- Hong EH, Kim SD, Hwang YH, Lee YH, Moon YH, Park RK (1986) A new good quality and high-yielding adzuki bean variety, Chungjupat. *Research Reports, Rural Development Administration, Korea Republic, Crops* **28**(1), 148-150. [In Korean]
- Hong EH, Kim SD, Lee YH, Hwang YH, Moon YH, Ham YS (1984) Chungwonpat, a new high yielding variety of *Vigna angularis* tolerant of late planting. *Research Reports, Office of Rural Development, S. Korea, Crop* **26**(2), 82-84. [In Korean]
- Hong EH, Lee SY, Lee YH, Moon YH, Kim SD (1993) Early maturity, good seed quality, new adzuki bean variety "Chungbupat". *RDA Journal of Agricultural Science, Upland & Industrial Crops* **35**(2), 179-183. [In Korean]
- Hong Z, Lakkineni K, Zhang Z, Verma DPS (2000) Removal of feedback inhibition of  $\Delta 1$ -pyrroline-5-carboxylate synthetase results in increased proline accumulation and protection of plants from osmotic stress. *Plant Physiology* **122**(4), 1129-1136.
- Hong ZL, Miao GH, Verma DPS (1993) p34cdc2 protein kinase homolog from mothbean (*Vigna aconitifolia*). *Plant Physiology* **101**(4), 1399-1400.
- Hori Y, Murakoso T, Fukumura M, Toriiizuka K, Ida Y (2009) Constituents and antioxidative activity of a hot-water extract of adzuki (*Vigna angularis*) beans. *Journal of Japanese Society of Nutrition and Food Science* **62**(1), 3-11.
- Hori Y, Sato S, Hatai A (2006) Antibacterial activity of plant extracts from azuki beans (*Vigna angularis*) in vitro. *Phytotherapy Research* **20**(2), 162-164.
- Horino T, Fukuoka T, Hagio T (1992) Differences in the nitrogen and mineral contents of seeds of cereals, buckwheat and pulses. *Japanese Journal of Crop Science* **61**(1), 28-33.
- Horng S, Lin H, Wu W, Godfray HCJ (1999) Behavioral processes and egg-laying decisions of the bean weevil, *Callosobruchus maculatus*. *Researches on Population Ecology* **41**(3), 283-290.
- Horvat F, Stainier F (1980b) Study of the exine in the Phaseolus-Vigna complex and related genera. IV. *Pollen et Spores* **22**(2), 139-172. [In French]
- Horáčková V (1986) Resistance of potato varieties to Fusarium rot in relation to duration of growth period, starch content and culinary quality. *Vedecké Práce Výzkumného a Šlechtitelského Ústavu Bramborářského v Havlíčkově Brodě* **10**, 41-50. [In Czech]
- Hoson T (1990a) Effect of auxin on autolysis of cell walls in azuki bean epicotyls. *Plant and Cell Physiology* **31**(2), 281-287.
- Hoson T (1990b) Regulation of cell wall metabolism by auxin. *Chemical Regulation of Plants* **25**(1), 40-56.
- Hoson T, Kamisaka S, Masuda Y (1996) Suppression of gravitropic response of primary roots by submergence. *Planta* **199**(1), 100-104.
- Hoson T, Masuda Y (1991) Effect of xyloglucan nonasaccharide on cell elongation induced by 2,4-dichlorophenoxyacetic acid and indole-3-acetic acid. *Plant and Cell Physiology* Hoson T, Masuda Y, Sone Y, Misaki A (1991) Xyloglucan antibodies inhibit auxin-induced elongation and cell wall loosening of azuki bean epicotyls but not of oat coleoptiles. *Plant Physiology* **96**(2), 551-557.
- Hoson T, Sone Y, Misaki A, Masuda Y (1993) Role of xyloglucan breakdown in epidermal cell walls for auxin-induced elongation of azuki bean epicotyl segments. *Physiologia Plantarum* **87**(2), 142-147.
- Hoson T, Tabuchi A, Masuda Y (1995) Mechanism of oxyloglucan breakdown in cell walls of azuki bean epicotyls. *Journal of Plant Physiology* **147**(2), 219-224.
- Hoson T, Wakabayashi K, Masuda Y (1995) Inhibition of the breakdown of xyloglucans in azuki bean epicotyls by concanavalin A. *Plant and Cell Physiology* **36**(5), 897-902.
- Hsieh CH (1990) An econometric analysis of the rational expectations of farm product prices under risk: the case of Adzuki beans in Taiwan. *Journal of Agricultural Economics (Nongye Jingyi)*(48), 35-45.
- Hsieh HM, Pomeranz Y, Swanson BG (1992) Composition, cooking time, and maturation of azuki (*Vigna angularis*) and common beans (*Phaseolus vulgaris*). *Cereal Chemistry* **69**(3), 244-248.

- Hsieh HM, Swanson BG, Lumpkin TA (2000) Abrasion, grinding, cooking and the composition and physical characteristics of azuki Koshi an. *Journal of Food Processing and Preservation* **24**(2), 87-106.
- Hsieh LC, Chang YC, Chien CC (1988) Studies on the ecology of the powdery mildew of beans. *Journal of Agricultural Research of China* **37**(1), 76-85.
- Hsu C, Chen Y (1994) The new adzuki bean cultivar Kaohsiung No. 6. *Taiwan Agriculture* **30**(5), 90-100.
- Hsu SH (1983) Studies on dew. Dew on winter crops of dry land in southwest Taiwan. *National Science Council Monthly* **11**(6), 527-536.
- Hu CAA, Delauney AJ, Verma DPS (1992) A bifunctional enzyme ( $\Delta 1$ -pyrroline-5-carboxylate synthetase) catalyzes the first two steps in proline biosynthesis in plants. *Proceedings of the National Academy of Sciences of the United States of America* **89**(19), 9354-9358.
- Hu T, Tsai LT, Fu YK, Shen ML (1985) Utilization of gamma rays for insect pest control of bagged mungbean and adzuki bean. *Plant Protection Bulletin, Taiwan* **27**(1), 63-68.
- Hu W, Lan Y, Horng S (1995) Effects of bean size on larval competition and oviposition preference of *Callosobruchus maculatus*. *Chinese Journal of Entomology* **15**(4), 321-332.
- Hu X, Zhang Y, Hu C, Tao M, Chen S (1998a) A comparison of methods for drying seeds: vacuum freeze-drier versus silica gel. *Seed Science Research* **8**(Supplement No. 1), 29-33.
- Hu XR, Zhang YL, Hu CL, Tao M, Chen SP (1998b) A comparison of methods for drying seeds: vacuum freeze-drier versus silica gel. *Seed Science Research* **8**, 29-33.
- Huang C, Horng S (2001) Superparasitism of seeds affects the life history strategy of the adult bean weevil, *Callosobruchus maculatus* (Fab.). *Formosan Entomologist* **21**(3), 287-297.
- Huang C, Yang R, Lee H, Horng S (2005) Beyond fecundity and longevity: trade-offs between reproduction and survival mediated by behavioural responses of the seed beetle, *Callosobruchus maculatus*. *Physiological Entomology* **30**(4), 381-387.
- Huang LQ, Qi ZT (1994) A new Penicillium from *Phaseolus angularis*. *Acta Mycologica Sinica* **13**(4), 264-266.
- Huang PM, Ge KL (1989) Plant regeneration by callus derived from mesophyll protoplast of *Phaseolus angularis*. *Acta Agriculturae Shanghai* **5**(1), 31-36.
- Huang R, Huang T (2002) The physicochemical properties of adzuki bean. *Taiwanese Journal of Agricultural Chemistry and Food Science* **40**(4), 280-287.
- Huang R, Huang T (2003) Effects of storage on the physicochemical properties of adzuki bean and its ann (red bean jam). *Taiwanese Journal of Agricultural Chemistry and Food Science* **41**(1), 28-36. [In Thai]
- Huber W (1979) The role of abscisic acid and cytokinins in plants under stress conditions. *Berichte der Deutschen Botanischen Gesellschaft* **92**(1), 193-207. [In German]
- Huber W, Kreutmeier F, Sankhla N (1977) Eco-physiological studies on Indian arid zone plants. 6. Effect of sodium chloride and abscisic acid on amino-acid and protein metabolism in leaves of *Phaseolus aconitifolius*. *Zeitschrift fur Pflanzenphysiologie* **81**(3), 234-247.
- Hung AT (1986) The multiple cropping system in Kaohsiung district of Taiwan. 41-56.
- Huss DL (1978) Importance of range development in dryland systems of farming for integration of crops and livestock husbandries in the Near East. 188-192.
- Hussain T, Imura O (1989) Effects of gamma radiation in survival and reproduction of *Callosobruchus chinensis* (L.) (Coleoptera: Bruchidae). *Applied Entomology and Zoology* **24**(3), 273-280.
- Hussain T, Imura O, Qureshi ZA (1994) Effect of gamma radiation on postembryonic development following irradiation of *Callosobruchus chinensis* eggs. *Pakistan Journal of Zoology* **26**(1), 7-9.
- Hussain T, Tontisirin K, Chaowanakarnkit L (1983) Protein digestibility of weaning foods prepared from rice minced meat and rice-mungbean combination in infants using a short-term nitrogen-balance method. *Journal of Nutritional Science and Vitaminology* **29**(4), 497-508.
- Ibrahim AES (1995) Summer forage yield of irrigated Sudan grass (*Sorghum sudane[n]se* (Piper) Stapf) and phillipesera (*Vigna trilobata* Verd.) in monoculture and mixtures. *Tropical Science* **35**(1), 1-8.
- Igarashi Y, Yoshioka Y, Sanada Y, Yamaguchi-Shinozaki K, Wada K, Shinozaki K (1997) Characterization of the gene for  $\Delta 1$ -pyrroline-5-carboxylate synthetase and correlation between the expression of the gene and salt tolerance in *Oryza sativa* L. *Plant Molecular Biology* **33**(5), 857-865.
- Igbedioh SO, Shaire S, Aderiye BJI (1995) Effects of processing on total phenols and proximate composition of pigeonpea (*Cajanus cajan*) and climbing bean (*Vigna umbellata*). *Journal of Food Science and Technology-Mysore* **32**(6), 497-500.
- Ignacimuthu S (2000) Ribosomal DNA repeat unit polymorphism in some pulses. *Indian Journal of Experimental Biology* **38**(2), 196-198.

- Ignacimuthu S, Schumann K, Zink D, Nagl W (1997) [Restriction fragment length polymorphisms of the rRNA genes in some pulses](#). *Current Science* **72**(9), 624-626.
- Iida T, Kawane M, Okubo K (1999) Reactive oxygen-scavenging compound isolated from Adzuki beans. In '!' pp. 261-270. (AOCS Press: Champaign)
- Iida T, Yoshiki Y, Kahara T, Okubo K, Ohru H (1997) A saponin conjugated with 2,3-dihydro-2,5-dihydroxy-6-methyl-4H-pyran-4-one from *Vigna angularis* [seeds]. *Phytochemistry* **45**(7), 1507-1509.
- Iida T, Yoshiki Y, Okubo K, Ohru H, Kinjo J, Nohara T (1999) Triterpenoid saponins from *Vigna angularis*. *Phytochemistry* **51**(8), 1055-1058.
- Iizuka N (1990) Studies on virus diseases of adzuki bean (*Vigna angularis* Wight) in Japan. *Bulletin of the Tohoku National Agricultural Experiment Station*(82), 77-113.
- Ikehashi H (1985) New summer crop cultivars registered by the Ministry of Agriculture, Forestry and Fisheries in 1985. Azuki bean. *Japanese Journal of Breeding* **35**(4), 458-459.
- Ikushima T, Shimmen T (2005) [Mechano-sensitive orientation of cortical microtubules during gravitropism in azuki bean epicotyls](#). *Journal of Plant Research* **118**(1), 19-26.
- Ikushima T, Soga K, Hoson T, Shimmen T (2008) Role of xyloglucan in gravitropic bending of azuki bean epicotyl. *Physiologia Plantarum* **132**(4), 552-565.
- Imagawa K, Toko K, Ezaki S, Hayashi K, Yamafuji K (1991) Electric potentials during gravitropism in bean epicotyls. *Plant Physiology* **97**(1), 193-196.
- Imai T, Chinushi T, Hoshino S, Shimada N (1990) On the white spots surrounding the hilum of azuki bean (*Vigna angularis*). *Journal of the Niigata Agricultural Experiment Station*(37), 63-64.
- Imura O (1991) A comparative study of the feeding habits of *Tribolium freemani* Hinton and *Tribolium castaneum* (Herbst) (Coleoptera: Tenebrionidae). *Applied Entomology and Zoology* **26**(2), 173-182.
- Inagawa H, Nishizawa T, Tsukioka D, Suda T, Chiba Y, Okutomi T, Morikawa A, Soma GI, Mizuno D (1992) Homeostasis as regulated by activated macrophage. II. LPS of plant origin other than wheat flour and their concomitant bacteria. *Chemical and Pharmaceutical Bulletin* **40**(4), 994-997.
- Inoue K, Yokota H, Yamada Y (1988) Effect of Ca in the medium on root growth under low pH conditions. *Soil Science and Plant Nutrition* **34**(3), 359-374.
- Inouhe M, Ito R, Ito S, Sasada N, Tohoyama H, Joho M (2000) Azuki bean cells are hypersensitive to cadmium and do not synthesize phytochelatins. *Plant Physiology* **123**(3), 1029-1036.
- Inouhe M, Yamamoto R (1991) Effects of 2-deoxygalactose on auxin-induced growth and levels of UDP-sugars in higher plants. *Plant and Cell Physiology* **32**(3), 433-438.
- Inouye N, Maeda T, Mitsuhashi K (1982) Cucumber mosaic virus isolated from *Calanthe discolor*. *Nogaku Kenkyu* **60**(1), 1-11.
- International Board for Plant Genetic R, No (1988) Report on an IBPGR mission in northern Thailand for the collection of *Vigna umbellata* var. gracilis.. In. ')
- Ipe IM, Bhati YKS (1977) On the biology of *Melanagromyza sojae* (Zehnt.) (Diptera, Agromyzidae) and on its incidence on five pulse crops in the Agra district, India. *Zur Biologie von Melanagromyza sojae (Zehnt.) (Diptera, Agromyzidae) und zu ihrem Auftreten in funf Hulsenfruchtarten bei Agra, Indien.* **50**(7), 104-106.
- Iriawati, Miyake H, Taniguchi T, Takeoka Y (1996a) [Effects of iodoacetamide on the fine structure of azuki bean protoplast and its fusion product](#). *Japanese Journal of Crop Science* **65**(3), 532-539.
- Iriawati, Miyake H, Taniguchi T, Takeoka Y (1996b) [Ultrastructural studies on the fusion product between protoplasts of mung bean and adzuki bean](#). *Japanese Journal of Crop Science* **65**(2), 330-339.
- Isemura T, Kaga A, Konishi S, Ando T, Tomooka N, Han O, Vaughan DA (2007) [Genome dissection of traits related to domestication in azuki bean \(\*Vigna angularis\*\) and comparison with other warm-season legumes](#). *Annals of Botany* **100**(5), 1053-1071.
- Ishige F, Mori H, Yamazaki K, Imaseki H (1993a) Cloning of a complementary DNA that encodes an acidic chitinase which is induced by ethylene and expression of the corresponding gene. *Plant and Cell Physiology* **34**(1), 103-111.
- Ishige F, Mori H, Yamazaki K, Imaseki H (1993b) Identification of a basic glycoprotein induced by ethylene in primary leaves of azuki bean [*Vigna angularis*] as a cationic peroxidase. *Plant Physiology* **101**(1), 193-199.
- Ishige F, Yamazaki K, Mori H, Imaseki H (1991) The effects of ethylene on the coordinated synthesis of multiple proteins: accumulation of an acidic chitinase and a basic glycoprotein induced by ethylene in leaves of azuki bean, *Vigna angularis*. *Plant and Cell Physiology* **32**(5), 681-690.
- Ishihara S, Amano K (1980) Studies on the mechanization of harvesting field beans-development and improvement of the pod-gathering device for the bean special combine. *Research Bulletin of the Hokkaido National Agricultural Experiment Station*(126), 135-147.

- Ishii Y, Shimada M (2008) Competitive exclusion between contest and scramble strategists in *Callosobruchus* seed-beetle modeling. *Population Ecology* **50**(2), 197-205.
- Ishimoto M (1996) Identification of bruchid resistance factor in the common bean and enhancement of bruchid resistance in the azuki bean. *Gamma Field Symposia*(35), 39-50.
- Ishimoto M, Kitamura K (1989) Growth inhibitory effects of an  $\alpha$ -amylase inhibitor from the kidney bean, *Phaseolus vulgaris* (L.) on three species of bruchids (Coleoptera: Bruchidae). *Applied Entomology and Zoology* **24**(3), 281-286.
- Ishimoto M, Sato T, Chrispeels MJ, Kitamura K (1996) Bruchid resistance of transgenic azuki bean expressing seed  $\alpha$ -amylase inhibitor of common bean. *Entomologia Experimentalis et Applicata* **79**(3), 309-315.
- Ishwar S, Rathore MS, Chandawat MS (2003) Effect of thiourea on dew bean grown in low fertile undulating soil of the Thar Desert. In '!' pp. 233-235. (Scientific Publishers (India): Jodhpur)
- Islam A, Duara PK, Barua PK (2002) Genetic variability in a set of rice bean genotypes assessed over sowing dates. *Journal of the Agricultural Science Society of North-East India* **15**(1), 61-66.
- Ismail S, Wankhede DB, Syed HM, Kulkarni AS, Fayazuddin M (2003) The effect of germination on the changes in protein, free amino acid and in-vitro protein digestibility of moth bean and horse gram. In '!' pp. 315-317. (Scientific Publishers (India): Jodhpur)
- Isobe K, Tsuboki Y (1998a) Relationship between the amount of root exudate and the infection rate of arbuscular mycorrhizal fungi in gramineous and leguminous crops. *Plant Production Science* **1**(1), 37-38.
- Isobe K, Tsuboki Y (1998b) The relationship between growth promotion by arbuscular mycorrhizal fungi and root morphology and phosphorus absorption in gramineous and leguminous crops. *Japanese Journal of Crop Science* **67**(3), 347-352.
- Itagaki M, Soejima H, Ishii K, Sugiyama T, Hayashi Y (2003) Biological activities and structure-activity relationship of substitution compounds of N-[2-(3-indolyl)ethyl]succinamic acid and N-[2-(1-naphthyl)ethyl]succinamic acid, derived from a new category of root-promoting substances, N-(phenethyl)succinamic acid analogs. *255*(1), 67-75.
- Ito S, Kojima M, Ohnishi M, Fujino Y (1985) Triterpenic lipids in azuki bean seeds. *Journal of the Agricultural Chemical Society of Japan* **59**(9), 895-900.
- Itoh T (2002) Localization of cellulose synthases in terminal complexes - demonstration by freeze-fracture techniques coupled with immunogold-labeling. *Regulation of Plant Growth & Development* **37**(1), 44-50.
- Itoh T, Furuichi Y (2005) Hot-water extracts from adzuki beans (*Vigna angularis*) stimulate not only melanogenesis in cultured mouse B16 melanoma cells but also pigmentation of hair color in C3H mice. *Bioscience, Biotechnology and Biochemistry* **69**(5), 873-882.
- Itoh T, Furuichi Y (2009) Lowering serum cholesterol level by feeding a 40% ethanol-eluted fraction from HP-20 resin treated with hot water extract of adzuki beans (*Vigna angularis*) to rats fed a high-fat cholesterol diet. *Nutrition* **25**(3), 318-321.
- Itoh T, Itoh Y, Hibasami H, Katsuzaki H, Imai K, Furuichi Y, Komiya T (2005a) Isolation of a substance inducing apoptosis of cultured human gastric cancer cells from a hot-water extract of adzuki (*Vigna angularis*). *Journal of Japanese Society of Nutrition and Food Science* **58**(5), 281-287.
- Itoh T, Itoh Y, Hibasami H, Katsuzaki H, Imai K, Furuichi Y, Komiya T (2005b) Vignoside, a novel new sesquiterpene glucoside obtained from a hot-water extract of adzuki bean (*Vigna angularis*). *Nippon Shokuhin Kagaku Kogaku Kaishi = Journal of the Japanese Society for Food Science and Technology* **52**(7), 319-323.
- Itoh T, Itoh Y, Mizutani M, Fujishiro K, Furuichi Y, Komiya T, Hibasami H (2002) A hot water extract of adzuki (*Vigna angularis*) induces apoptosis in cultured human stomach cancer cells. *Nippon Shokuhin Kagaku Kogaku Kaishi = Journal of the Japanese Society for Food Science and Technology* **49**(5), 339-344.
- Itoh T, Itoh Y, Mizutani M, Fujishiro K, Furuichi Y, Komiya T, Hibasami H (2004) Hot-water extracts from adzuki beans (*Vigna angularis*) suppress not only the proliferation of KATO III cells in culture but also benzo(a)pyrene-induced tumorigenesis in mouse forestomach. *Journal of Nutritional Science and Vitaminology* **50**(4), 295-299.
- Itoh T, Kita N, Kurokawa Y, Kobayashi M, Horio F, Furuichi Y (2004) Suppressive effect of a hot water extract of adzuki beans (*Vigna angularis*) on hyperglycemia after sucrose loading in mice and diabetic rats. *Bioscience, Biotechnology and Biochemistry* **68**(12), 2421-2426.
- Itoh T, Kobayashi M, Horio F, Furuichi Y (2009) Hypoglycemic effect of hot-water extract of adzuki (*Vigna angularis*) in spontaneously diabetic KK-Ay mice. *Nutrition* **25**(2), 134-141.

- Itoh T, Umekawa H, Furuichi Y (2005) Potential ability of hot water adzuki (*Vigna angularis*) extracts to inhibit the adhesion, invasion, and metastasis of murine B16 melanoma cells. *Bioscience, Biotechnology and Biochemistry* **69**(3), 448-454.
- Iwanami Y (1982) Inhibition of phototropism and gravitropism in cuttings of bean (*Phaseolus angularis*) by treatment with myrmic acid and related compounds. *Botanical Gazette* **143**(4), 483-485.
- Iwasaki A (1998) Major injury period of adzuki bean seeds by tarnished plant bug, *Lygus rugulipennis* Poppius. *Annual Report of the Society of Plant Protection of North Japan*(49), 146-149.
- Iwasaki A (2001) Effect of insecticides on the European tarnished plant bug, *Lygus rugulipennis* Poppius (Heteroptera, Miridae). 3. Control of damage of adzuki beans, *Vigna angularis*, in the field. Society of Plant Protection of North Japan, Akita.
- Iwasaki A, Furukawa K (1996) Injury to adzuki bean seeds caused by tarnished plant bug, *Lygus rugulipennis* Poppius (Heteroptera: Miridae), in Hokkaido. *Annual Report of the Society of Plant Protection of North Japan*(47), 118-121.
- Jaaska V (1996) Isoenzyme diversity and phylogenetic affinities among the *Phaseolus* beans (Fabaceae). *Plant Systematics and Evolution* **200**(3/4), 233-252.
- Jaaska V, Jaaske V (1990b) Isoenzyme variation in Asian beans. *Botanica Acta* **103**(3), 281-290.
- Jadhav PS, Mulik NG, Chavan PD (1997) Allelopathic effects of *Ipomoea carnea* spp. *fistulosa* on growth of wheat, rice, sorghum and kidneybean. *Allelopathy Journal* **4**(2), 345-348.
- Jadhav SJ, Sonone AH, Aher RP, Dahat DV (2001b) Variability and character association in rice bean {*Vigna umbellata* (Thunb) Ohwi and Ohashi}. *Legume Research* **24**(2), 105-108.
- Jagnow G (1971) Observations on root nodules of fodder legumes and on the influence of urea and superphosphate fertilizers on the development of root nodules on groundnuts in the irrigated area of Gezira (Sudan). *Zeitschrift fur Acker- und Pflanzenbau* **134**(2), 125-134. [In German]
- Jai Rao K, Thirumalachar DK (1977) New record of alternate host plants of groundnut leaf miner *Stomopteryx subsecivella* Zeller (Syn.: *S. nerteria* Meyrick) (Lepidoptera: Gelechiidae). *Current Science* **46**(3), 91-92.
- Jain GL (1972) Secondary and micro-nutrients in relation to crops in arid zone of India. II. Effect of spray application of S, Fe, Zn and Mo on growth, post harvest characters and yield of moth (*Phaseolus aconitifolius* jacq.). *Indian Journal of Agronomy* **17**(4), 271-278.
- Jain GL, Raheja PC (1980) Secondary and micronutrients in relation to crops in arid zone of India-VII. Nutrient uptake in moth as influenced by spray application of S, Fe, Zn and Mo. *Indian Journal of Agronomy* **25**(4), 685-690.
- Jain HK Genetic improvement and production prospects of food legumes. 33-42.
- Jain HK, Mehra KL (1980a) Evolution, adaptation, relationships, and uses of the species of *Vigna* cultivated in India. 459-468.
- Jain J, Chopra VL (1988) Genotypic differences in response to regeneration of in vitro cultures of moth bean *Vigna aconitifolia* (Jacq.) Marechal. *Indian Journal of Experimental Biology* **26**(9), 654-656.
- Jain KC, Bajpai MR (1971) Economics of the common mixture of mixed cropping in the semi-arid tract of central Rajasthan. *Annals of Arid Zone* **10**(4), 251-254.
- Jain NC, Babu CR (1982) Seed coat polymorphism in *Vigna calcarata* and its evolutionary significance. *Seed Science and Technology* **10**(3), 451-456.
- Jain SC (1985) Insecticidal control of yellow mosaic virus disease of moth bean transmitted by whitefly (*Bemisia tabaci* Genn.). *Indian Botanical Reporter* **4**(1), 87-88.
- Jain SC (1992) Evaluation of moth bean (*Vigna aconitifolia* (Jacq.) Marechal) lines against yellow mosaic virus. *Madras Agricultural Journal* **79**(2), 118-119.
- Jain SC, Singh RC, Sharma RC, Mathur JR (1982) Seed mycoflora of moth bean (*Vigna aconitifolia*): its pathogenicity and control. *Indian Journal of Mycology and Plant Pathology* **12**(2), 137-141.
- Jakhar SS, Chauhan MS, Duhan JC (1998) Management of root rot of cotton (*Gossypium* species) caused by Rhizoctonia species. *Plant Disease Research* **13**(2), 115-118.
- Jakhmola RC, Roy AK (1995) Feeding energy supplements to adult male camels during winter. *Indian Journal of Animal Sciences* **65**(1), 80-83.
- Jamadar MM, Balikai RA, Nayakar NY (1998) Reaction of KBMB-1, a mothbean variety to diseases under dryland conditions. *Karnataka Journal of Agricultural Sciences* **11**(1), 246-247.
- Jat BL, Yadav GL, Singh M (2003) Dalbergia sissoo based agroforestry model for arid regions. *Environment and Ecology* **21**(1), 31-33.
- Java M, Mishra KN (2008) Effect of different strains of Rhizobium on nitrogen fixation and other growth characteristics of two leguminous species. *Plant Archives* **8**(1), 307-310.

- Jaya M, Mishra KN (2007) Effect of SO<sub>2</sub> and NO<sub>2</sub> generated acidification on nodulation of Alysicarpus monilifer and Phaseolus trilobus (Linn) in alluvials of U.P., India. *National Academy Science Letters* **30**(5/6), 165-168.
- Jayachandran M, Veerabadran V (1996) Nitrogen economy through intercropping green manure in semi dry rice. *Madras Agricultural Journal* **83**(7), 466-468.
- Jenjiha JK, Kalia NR (2007a) Studies on genetic variability in rice bean [*Vigna umbellata* (Thunb) Ohwi and Ohashi] under monocropping and intercropping system. *Agricultural Science Digest* **27**(3), 162-165.
- Jenjiha JK, Kalia NR (2007b) Studies on genetic variability in rice bean [*Vigna umbellata* (Thunb) Ohwi and Ohashi] under monocropping and intercropping system. *Agricultural Science Digest* **27**(3), 162-165.
- Jeswani LM, Baldev B (1990) 'Advances in pulse production technology.' (ICAR: New Delhi) viii + 190□pp.
- Jhansi K (2004) Management of shootfly and stem borer incidence in fodder sorghum by intercropping with leguminous fodder crops. *Indian Journal of Plant Protection* **32**(2), 139.
- Jhorar BS, Dhillon RS, Hooda MS (2006) Sand dune stabilization and management of sandy soils through agro-forestry models. In ' pp. 311-315. (Westville Publishing House: New Delhi)
- Jia S, Wang B, Zhang L, Guan X (2007) Studies on pollen submicroscopic morphology and numerical taxonomy of common bean (*Phaseolus vulgaris* L.) and its related species. *Acta Agriculturae Zhejiangensis* **19**(2), 79-83.
- Jiang LQ, Din BZ, Jin WL, Fen ST (1986) Investigations on flowering and fruiting in *Phaseolus angularis*. *Journal of Agricultural Sciences, Jiangsu Academy of Agricultural Sciences* **2**(2), 41-44.
- Jin W (1995) An initial study on ecoclimatic division of adzuki bean resources in China. *Beijing Agricultural Sciences* **13**(6), 1-5.
- Jin W, Bai L, Wen Z, Pu S, Zhao B (2006) Analysis on genetic system of 100-seed weight in adzuki bean. *Acta Agronomica Sinica* **32**(9), 1410-1412.
- Jin W, Hirofumi Y, Matiko I, Kentaro Y (2003a) The sequence variations of intron-3 of the  $\alpha$ -amylase gene in adzuki bean. *Agricultural Sciences in China* **2**(10), 1076-1082.
- Jin W, Hirofumi Y, Matiko I, Kentaro Y (2003b) The variation of the intron-3 at the  $\alpha$ -amylase gene in adzuki bean. *Scientia Agricultura Sinica* **36**(5), 480-486.
- Jin W, Pu S, Zhao B, Su L (2006a) genetic variation of 100-seed-weight and starch granule size in adzuki bean germplasm. *Acta Agriculturae Boreali-Sinica* **21**(6), 41-44.
- Jin W, Pu S, Zhao B, Wang L, Wu G, Su L (2006b) Genetic variation of quality characters of seed in adzuki bean germplasm resources. *Acta Agronomica Sinica* **32**(8), 1223-1230.
- Jin W, Pu S, Zhao B, Wu G (2005) Genetic variation of seed colour and sweetened bean purée producing rate of adzuki bean germplasm resources. *Acta Agriculturae Boreali-Sinica* **20**(6), 28-33.
- Jin W, Wen Z, Pu S, Zhao B (2004) Genetic diversity of adzuki bean (*Vigna angularis*) germplasm resources based on RAPD markers. *Acta Agronomica Sinica* **30**(7), 686-691.
- Jin WL (1989) Characteristics and breeding potential of some northern varieties of adzuki bean in Nanjing. *Zuowu Pinzhong Ziyuan*(3), 10-12.
- Jin WL, Yamaguchi H, Futsuhara Y (1993) Study of interspecific differences between cultivated and wild adzuki bean species. I. Differences in and analysis of correlations between the main characters of adzuki bean seeds and seedlings. *Crop Genetic Resources*(1), 4-6.
- Jindal JK, Patel PN (1980) Variability in xanthomonads of grain legumes .1. Pathogenic behavior and taxonomy. *Phytopathologische Zeitschrift-Journal of Phytopathology* **99**(4), 332-356.
- Jindal SK (1988) Intra-specific divergence in moth bean for grain yield and its components. *Annals of Arid Zone* **27**(3-4), 267-270.
- Jindal SK, Vir S (1983) Adaptability in moth bean (*Vigna aconitifolia* (Jacq.) Marechal) for seed yield in rainfed conditions. *Annals of Arid Zone* **22**(1), 83-86.
- Jolli RB, Ekbote SD (2005) Effect of seed treatment and packaging material on bruchid control and viability of moth bean seeds. *Karnataka Journal of Agricultural Sciences* **18**(1), 174-176.
- Jolli RB, Karabhantanal SS, Jayaprakash TC (2005) Influence of pesticides and storage methods in the management of pulse beetle (*Callosobruchus* spp.) and their effects on seed viability in mothbean. *Journal of Entomological Research* **29**(2), 159-162.
- Joomjantha S, Wanapat M (2008a) Effect of intercropping on biomass yield and chemical composition of cassava. *Livestock Research for Rural Development* **20**(Suppl.), unpaginated.
- Joomjantha S, Wanapat M (2008b) Effect of supplementation with tropical protein-rich feed resources on rumen ecology, microbial protein synthesis and digestibility in swamp buffaloes. *Livestock Research for Rural Development* **20**(Suppl.), unpaginated.

- Joseph LS, Bouwkamp JC (1978) [Karyomorphology of several species of \*Phaseolus\* and \*Vigna\*](#). *Cytologia* **43**(3/4), 595-600.
- Joshi KD, Bhandari B, Gautam R, Bajracharya J, Hollington PA Ricebean: a multipurpose underutilised legume. 2008a, pp. 234-248
- Joshi NL (1989) [Efficient use of energy inputs in rainfed pearl millet](#). *Agriculture, Ecosystems and Environment* **25**(2-3), 91-102.
- Joshi NL (1997) Response of pearl millet-based cropping systems to nitrogen in the arid region. 5-23.
- Joshi NL (1999) Pearl millet-based cropping systems to achieve sustainability in the arid region. 147-154.
- Joshi NL, Panjab S (1981) Towards higher bajra yields in arid Rajasthan. *Indian Farming* **31**(4), 3-4, 17.
- Joshi S, Dwivedi P, Ohar DN (1989) Chemical examination of *Phaseolus trilobus* Ait. seeds. *Herba Hungarica* **28**(1-2), 67-69.
- Joshi S, Upadhyaya A, Davis TD, Sankhla AK, Sankhla D, Sankhla N (1988) Interaction between abscisic acid and GA3 in in vitro regeneration from moth bean callus. 35-38.
- Junaid M, Rathi YPS, Khan AA (1991b) Seed borne mycoflora of rice bean (*Vigna umbellata* L.) and the effect of seed treatment on seed germinability. *Bioved* **2**(1), 67-70.
- Jung J, Seo B, Cho J, Kwon Y, Kim G (2009) Occurrence of lepidopteran insect pests and injury aspects in adzuki bean fields. *Korean Journal of Applied Entomology* **48**(1), 29-35. [In Korean]
- Jyoti S, Swarnkar PL (1996) Morphogenetic and biochemical variations under different spectral lights in leaf cultures of *Vigna aconitifolia*. *Journal of Phytological Research* **9**(1), 89-93.
- Jyoti S, Swarnkar PL (1998a) Isozymic variations in *Vigna aconitifolia* under salinity stress. *Journal of Phytological Research* **11**(2), 179-181.
- Jyoti S, Swarnkar PL (1998b) Organ specific isoperoxidases in *Vigna aconitifolia* under stress condition. *Indian Journal of Plant Physiology* **3**(4), 296-299.
- Jyotishi A, Deshmukh MR, Duhoon SS (2004) Profitable niger, *Guizotia abyssinica* (L.f) Coss. based intercropping studies with new crop combinations for different regions in India. *Journal of Oilseeds Research* **21**(1), 198-199.
- Jyotsna S, Satija CK (1996) In vitro hybridization in incompatible crosses of *Vigna* species. *Crop Improvement* **23**(1), 29-32.
- Kadam SS, Ghorpade VM, Adsule RN, Salunkhe DK (1986) [Trypsin inhibitor in moth bean: thermal stability and changes during germination and cooking](#). *Qualitas Plantarum Plant Foods for Human Nutrition* **36**(1), 43-46.
- Kaga A, Hosaka K, Kimura T, Misoo S, Kamijima O (1993) Application of random amplified polymorphic DNA (RAPD) analysis for adzuki bean and its related genera. *Science Reports of Faculty of Agriculture, Kobe University* **20**(2), 171-176.
- Kaga A, Isemura T, Tomooka N, Vaughan DA (2008) The genetics of domestication of the azuki bean (*Vigna angularis*). *Genetics* **178**(2), 1013-1036.
- Kaga A, Ishii T, Tsukimoto K, Tokoro E, Kamijima O (2000a) [Comparative molecular mapping in Ceratotropis species using an interspecific cross between azuki bean \(\*Vigna angularis\*\) and rice bean \(\*V-umbellata\*\)](#). *Theoretical and Applied Genetics* **100**(2), 207-213.
- Kaga A, Ohnishi M, Ishii T, Kamijima O (1996) [A genetic linkage map of azuki bean constructed with molecular and morphological markers using an interspecific population \(\*Vigna angularis\* × \*V. nakashimae\*\)](#). *Theoretical and Applied Genetics* **93**(5/6), 658-663.
- Kaga A, Tomooka N, Egawa Y, Hosaka K, Kamijima O (1996b) [Species relationships in the subgenus Ceratotropis \(genus Vigna\) as revealed by RAPD analysis](#). *Euphytica* **88**(1), 17-24.
- Kagawa H, Yamauchi F, Hirano H (1987) Soybean basic 7 S globulin represents a protein widely distributed in legume species. *FEBS Letters* **226**(1), 145-149.
- Kageyama K, Ui T (1983) Host range and distribution of *Pythium myriotylum* and unidentified *Pythium* sp. contributed to the monoculture injury of bean and soybean plants. *Annals of the Phytopathological Society of Japan* **49**(2), 148-152.
- Kahlon TS, Smith GE, Shao Q (2005) [In vitro binding of bile acids by kidney bean \(\*Phaseolus vulgaris\*\), black gram \(\*Vigna mungo\*\), bengal gram \(\*Cicer arietinum\*\) and moth bean \(\*Phaseolus aconitifolius\*\)](#). *Food Chemistry* **90**(1/2), 241-246.
- Kailash C, Singh M, Rathore MS, Shekhawat NS (2009) Somatic embryogenesis and in vitro plant regeneration in moth bean [*Vigna aconitifolia* (Jacq.) Marechal]: a recalcitrant grain legume. *Plant Biotechnology Reports* **3**(3), 205-211.

- Kaiser WJ, Wyatt SD, Klein RE (1991) Epidemiology and seed transmission of two tobacco streak virus pathotypes associated with seed increases of legume germ plasm in eastern Washington. *Plant Disease* **75**(3), 258-264.
- Kajiwara H, Hirano H (1992) Analysis of adzuki bean seed protein by partial amino acid sequencing of polypeptide separated on two-dimensional gel. *Japanese Journal of Breeding* **42**(4), 727-735.
- Kajiwara H, Tomooka N (1998) Comparative analysis of genus *Vigna* seeds using antiserum against a synthesized multiple antigenic peptide. *Electrophoresis* **19**(18), 3110-3113.
- Kakani RK, Garg DK, Sharma RC, Shiv N (2003) Evaluation of germplasm lines of moth bean. In '' pp. 107-111. (Scientific Publishers (India): Jodhpur)
- Kakani RK, Sharma RC, Garg DK (2003) Genetic divergence, variability and character association in moth bean. In '' pp. 98-103. (Scientific Publishers (India): Jodhpur)
- Kakepota TM, Naqvi SMSH, Rustamani MA, Munshi GM (1984) Relative resistances of different pulses to *Callosobruchus chinensis* Lin. and its population fluctuation. *Proceedings of the Entomological Society of Karachi*(14/15), 101-106.
- Kaku T, Tabuchi A, Wakabayashi K, Hoson T (2004) Xyloglucan oligosaccharides cause cell wall loosening by enhancing xyloglucan endotransglucosylase/hydrolase activity in azuki bean epicotyls. *Plant and Cell Physiology* **45**(1), 77-82.
- Kalia HR, Dawa T (1974) Studies on the nature of hard seeds in some important pulse crops. *Himachal Journal of Agricultural Research* **2**(1), 42-46.
- Kamble S, Misra HS, Mahajan SK, Eapen S (2003) A protocol for efficient biotic transformation of mothbean *Vigna aconitifolia* L. Jacq. Marechal. *Plant Molecular Biology Reporter* **21**(4), 457a-457j.
- Kamlesh S, Neelam K, Saroj B (2001a) Effect of dehulling soaking and germination on nutrient composition of ricebean and fababean. *Haryana Agricultural University Journal of Research* **31**(1/2), 35-40.
- Kamlesh S, Neelam K, Saroj B (2001b) Processing of newly released ricebean and fababean cultivars: changes in total and available calcium, iron and phosphorus. *International Journal of Food Sciences and Nutrition* **52**(5), 413-418.
- Kamlesh S, Neelam K, Saroj B (2002a) Antinutrients and protein digestibility of fababean and ricebean as affected by soaking, dehulling and germination. *Journal of Food Science and Technology (Mysore)* **39**(4), 418-422.
- Kamlesh S, Neelam K, Saroj B (2002b) Variability in physico-chemical properties and nutrient composition of newly released ricebean and fababean cultivars. *Journal of Food Composition and Analysis* **15**(2), 159-167.
- Kanamori M, Ikeuchi T, Ibuki F, Kotaru M, Kuang Kung K (1982) Amino acid composition of protein fractions extracted from *Phaseolus* beans and the field bean (*Vicia faba* L.). *Journal of Food Science* **47**(6), 1991-1994.
- Kanawade LR, Bhosale BW, Kadam MS (1990) Effects of moisture content on certain selected physical properties of pulse seeds. *Journal of Maharashtra Agricultural Universities* **15**(1), 60-62.
- Kandpal BK, Kumawat RN, Mertia RS (2006) Agronomic evaluation of promising genotypes of moth bean under hyper arid conditions of Jaisalmer. *Journal of Arid Legumes* **3**(1), 1-10.
- Kandpal BK, Mertia RS, Kumawat RN (2006) Evaluation of *Vigna aconitifolia* genotypes for their suitability to aeolian plains of extreme arid region. *Journal of Arid Legumes* **3**(1), 30-33.
- Kaneta M, Hikichi H, Endo S, Sugiyama N (1980) Identification of flavones in sixteen Leguminosae species. *Agricultural and Biological Chemistry* **44**(6), 1407-1408.
- Kaneta T, Kakimoto T, Shibaoka H (1993) Actinomycin D inhibits the GA3-induced elongation of azuki bean epicotyls and the reorientation of cortical microtubules. *Plant and Cell Physiology* **34**(7), 1125-1132.
- Kaneta T, Kakimoto T, Shibaoka H (1997) Gibberellin A3 causes a decrease in the accumulation of mRNA for ACC oxidase and in the activity of the enzyme in azuki bean (*Vigna angularis*) epicotyls. *Plant and Cell Physiology* **38**(10), 1135-1141.
- Kang TH, Cha YH, Chang YH (1980) Studies on inorganic matter in feedstuffs. III. Contents of inorganic matter in domestic feedstuffs. *Research Reports of the Office of Rural Development, Livestock & Veterinary* **22**, 46-56. [In Korean]
- Kang Y, Cho N, Yook W, Kang M (1998) Forage yield and quality of summer grain legumes and forage grasses in Cheju Island. *Korean Journal of Crop Science* **43**(4), 245-249.
- Kao CW (1991) Recent achievements in the biological control of plant diseases in ROC on Taiwan. 181-185.
- Kapila R (1988) Ecological status of host range in zabrotes-subfasciatus bohemian. *Indian Journal of Entomology* **50**(4), 503-512.
- Kar BC, Behera B, Satpathy SC, Mishra PK (1993b) Crop-weed competition in ricebean (*Vigna umbellata*). *Indian Journal of Agronomy* **38**(3), 511-512.

- Karadge BA, Kulkarni H (1992) Mineral nutrition of moth bean (*Phaseolus aconitifolius* Jacq.) under saline conditions. *Biovigyanam* **18**(2), 89-93.
- Karamac' M, Kosin'ska A, Rybareczyk A, Amarowicz R (2007) Extraction and chromatographic separation of tannin fractions from tannin-rich plant material. *Polish Journal of Food and Nutrition Sciences* **57**(4), 471-474.
- Karasawa T (2004) Arbuscular mycorrhizal associations and interactions in temperate cropping systems. (179), 1-71.
- Karasawa T, Arihara J, Kasahara Y (2000) Effects of previous crops on arbuscular mycorrhizal formation and growth of maize under various soil moisture conditions. *Soil Science and Plant Nutrition* **46**(1), 53-60.
- Karmakar PG, Biswas MR, Dana S (1989a) [Meiosis in allotetraploid of \*Vigna mungo\* X \*V. umbellata\* cross. Cytologia](#) **54**(1), 79-82.
- Karsono H, Saono S (1984) Effect of Rhizobium inoculation on the growth of *Vigna umbellata*.. In 'Pengaruh inokulan Rhizobium pada pertumbuhan *Vigna umbellata*.' pp. 137-138)
- Karuppuchamy P, Mohanasundaram M (1987) Bioecology and control of chilli muranai mite, *Polyphagotarsonemus latus* (Banks) (Taronemidae: Acari). *Indian Journal of Plant Protection* **15**(1), 1-4.
- Karwasra RS, Anil K, Sharma SK (2008) Effect of biofertilizers and chemical fertilizers on growth and yield of ricebean (*Vigna umbellata* L.). *Haryana Journal of Agronomy* **24**(1/2), 7-8.
- Kashiwaba K, Matsuda T, Hara H, Chonan N (1993) Fine structure related to reserves accumulation in azuki bean cotyledon (*Vigna angularis* L.). *Report of the Tohoku Branch, the Crop Science Society of Japan*(36), 29-32.
- Kashiwaba K, Tomooka N, Kaga A, Han OK, Vaughan DA (2003) Characterization of resistance to three bruchid species (*Callosobruchus* spp., Coleoptera, Bruchidae) in cultivated rice bean (*Vigna umbellata*). *Journal of Economic Entomology* **96**(1), 207-213.
- Katayama J, Sano K (1989) Damage analysis of red bean plants caused by the common cutworm, *Spodoptera litura* (Lepidoptera: Noctuidae). *Japanese Journal of Applied Entomology and Zoology* **33**(2), 57-62.
- Katayama TC (1979) Some morphological characters of pulses collected in the Indian sub-continent. *Memoirs of the Faculty of Agriculture, Kagoshima University* **15**, 29-48.
- Kathju S, Garg BK, Vyas SP, Lahiri AN (2003) Sustainable production of moth bean through genotype management under arid environments. *Journal of Arid Environments* **53**(1), 137-143.
- Kato A, Inouhe M (2008) [Mannose accommodation of \*Vigna angularis\* cells on solid agar medium involves its possible conversion to sucrose mediated by enhanced phosphomannose isomerase activity. Journal of Plant Research](#) **121**(3), 339-349.
- Kato A, Tohoyama H, Joho M, Inouhe M (2007) [Different effects of galactose and mannose on cell proliferation and intracellular soluble sugar levels in \*Vigna angularis\* suspension cultures. Journal of Plant Research](#) **120**(6), 713-719.
- Kato J (2000) Studies on characteristics for food processing of adzuki beans and common beans, and factors for their variation. (95), 91 pp.
- Kato J, Meguro T (1998) Two-dimensional indication of seed coat colours of adzuki beans [*Vigna angularis*] using lightness and chromaticity as coordinates. *Japanese Journal of Soil Science and Plant Nutrition* **69**(2), 190-194.
- Kato J, Meguro T, Suzuki MM, Deeth HC (2000) Variations in the seed coat colour of adzuki beans in the aspects of varieties, harvest years and growing locations, using two-dimensional colour mapping. *Plant Production Science* **3**(1), 61-66.
- Kato J, Souma C (2005) Nondestructive estimation method for polyphenol content and antioxidative activity of Adzuki bean. *Japanese Journal of Soil Science and Plant Nutrition* **76**(2), 205-208.
- Kato R, Nagayama E, Suzuki T, Uchida K, Shimomura TH, Harada Y (1993) Promotion of plant cell division by an epidermal growth factor. *Plant and Cell Physiology* **34**(6), 789-793.
- Kato S, Yamaguchi H, Shimamoto Y, Mikami T (2000) The chloroplast genomes of azuki bean and its close relatives: a deletion mutation found in weed azuki bean. *Hereditas (Landskrona)* **132**(1), 43-48.
- Kaur D, Kapoor AC (1990a) [Some antinutritional factors in rice bean \(\*Vigna-umbellata\*\) - effects of domestic processing and cooking methods. Food Chemistry](#) **37**(3), 171-179.
- Kaur D, Kapoor AC (1990b) [Starch and protein digestibility of rice bean \(\*Vigna-umbellata\*\) - effects of domestic processing and cooking methods. Food Chemistry](#) **38**(4), 263-272.
- Kaur D, Kapoor AC (1992) [Nutrient composition and antinutritional factors of rice bean \(\*Vigna-umbellata\*\). Food Chemistry](#) **43**(2), 119-124.
- Kaur M, Kawatra BL (1999) Molar ratios to predict zinc availability from domestically processed ricebean (*Vigna umbellata*). *Journal of Food Science and Technology-Mysore* **36**(6), 548-550.

- Kaur M, Kawatra BL (2000) Effect of domestic processing on flatus producing factors in ricebean (*Vigna umbellata*). *Nahrung-Food* **44**(6), 447-450.
- Kaur M, Mehta U (1993) Effect of partial and full substitution of bengalgram and blackgram dhals with ricebean on essential amino-acids and mineral contents of vada and pakora. *Journal of Food Science and Technology-Mysore* **30**(6), 454-456.
- Kaur R, Satija CK (1996) Study of incompatibility barriers between interspecific hybrids of *Vigna radiata* X *V. umbellata*. *Crop Improvement* **23**(2), 213-216.
- Kaur R, Satija CK (1998) Cytogenetical and biochemical analysis of interspecific hybrids between *Vigna radiata* and *V. umbellata*. *Indian Journal of Genetics and Plant Breeding* **58**(1), 25-34.
- Kaur S, Singh OS (1988) Response of ricebean to single and combined inoculation with rhizobium and glomus in a p-deficient sterilized soil. *Plant and Soil* **112**(2), 293-295.
- Kaushal RP, Singh BM (1988) Interspecific hybridization between urdbean (*Vigna mungo* (L.) Hepper) and adzukibean (*Vigna angularis* (Willd.) Ohwi and Ohashi). *Euphytica* **39**(1), 53-57.
- Kaushal RP, Singh BM (1989a) Evaluation of blackgram (*Phaseolus mungo*) germplasm for multiple-disease resistance. *Indian Journal of Agricultural Sciences* **59**(11), 726-727.
- Kaushal RP, Singh BM (1989b) Extra-chromosomal inheritance of resistance in *Vigna angularis* to Erysiphe polygoni. *Plant Disease Research* **4**(1), 67-68.
- Kaushal RP, Singh BM (1993) Pathogenic variability in leaf spot and powdery mildew pathogens of legumes. *Indian Phytopathology* **46**(2), 182-184.
- Kaushal RP, Singh BN (1991) Extra-chromosomal inheritance of resistance in adzuki bean (*Vigna angularis* (Willd.) Ohwi & Ohashi) to Erysiphe polygoni D.C. *Legume Research* **14**(2), 99-100.
- Kawai A, Fujita K (2007) Small red bean (azuki) sheds biologically active substances as a prerequisite step for germination, one of which displays the antiviral activity against the rabies virus infectivity and infections in culture. *Microbiology and Immunology* **51**(11), 1071-1079.
- Kawale SB, Kadam SS, Chavan UD, Chavan JK (2005) Effect of processing on insoluble dietary fiber and resistant starch in kidney bean and horse gram. *Journal of Food Science and Technology (Mysore)* **42**(4), 361-362.
- Kawecki TJ (1995) Expression of genetic and environmental variation for life history characters on the usual and novel hosts in *Callosobruchus maculatus* (Coleoptera : Bruchidae). *Heredity* **75**(1), 70-76.
- Keerio HK (1983) Effect of intercropping sorghum with fodder legumes. 645-652.
- Keerio HK, Singh SP (1985) Effect of intercropping on yield of hybrid sorghum under different nitrogen levels. *Pakistan Journal of Agricultural Research* **6**(1), 26-33.
- Kelkar NY, Maheshwari SC, Tyagi AK (1993) Light-dependent accumulation of mRNAs for chloroplast-encoded genes in *Vigna aconitifolia*. *Plant Science (Limerick)* **88**(1), 55-60.
- Kembhavi AA, Buttle DJ, Knight CG, Barrett AJ (1993) The two cysteine endopeptidases of legume seeds: purification and characterization by use of specific fluorometric assays. *Archives of Biochemistry and Biophysics* **303**(2), 208-213.
- Kennedy W, Denny R (1984) Bacterial stem rot, a new disease of adzuki beans in Minnesota. *Plant Disease* **68**(2), 167.
- Kevate BN, Chavan UD, Kadam SS, Chavan JK (2007) Isolation and characterization of starch from Moth bean. *Journal of Arid Legumes* **4**(2), 137-138.
- Khabiruddin M, Gupta SN, Sharma GD (1994b) Grain yield and proximate composition of ricebean (*Vigna umbellata*). *Legume Research* **17**(2), 116-118.
- Khabiruddin M, Gupta SN, Tyagi CS (2002b) Nutritional composition of some improved genotypes of ricebean (*Vigna umbellata*). *Forage Research* **28**(2), 104-105.
- Khair MAM (1989) Studies on irrigated tropical forage legumes in Sudan. 603-604.
- Khalafalla MM, El-Shemy HA, Mizanur RS, Teraishi M, Ishimoto M (2005) Recovery of herbicide-resistant Azuki bean [*Vigna angularis* (Wild.), Ohwi & Ohashi] plants via Agrobacterium- mediated transformation. *African Journal of Biotechnology* **4**(1), 61-67.
- Khalil JK, Sawaya WN, Al-Mohammad HM (1986) Effects of experimental cooking on the yield and proximate composition of three selected legumes. *Journal of Food Science* **51**(1), 233-234, 236.
- Khan MB, Mahboob A, Abdul K (2001a) Effect of planting patterns and different intercropping systems on the productivity of cotton (*Gossypium hirsutum* L.) under irrigated conditions of Faisalabad. *International Journal of Agriculture and Biology* **3**(4), 432-435.
- Khan MB, Mahboob A, Abdul K (2001b) Some competition functions and economics of different cotton-based intercropping systems. *International Journal of Agriculture and Biology* **3**(4), 428-431.

- Khan RA, Nuzhat S, Iqbal A (2007) Analgesic and antitussive activities on methanol extract of *Vigna trilobata* (L) Verdc. *Journal of Basic & Applied Sciences* **3**(2), 81-83.
- Khan SA, Chatterjee BN (1987) Multiple cropping in rain-fed upland entisol under different soil-management conditions in the subhumid tropics. *Soil & Tillage Research* **9**(2), 135-150.
- Khan TN, Erskine W, Kesavan V Improvement of food legumes at the University of Papua New Guinea. 1976, pp. 127-136
- Khanda CM, Mishra PK (1998) Effect of plant density and nitrogen fertilization on growth and yield of rice bean (*Vigna umbellata*). *Indian Journal of Agronomy* **43**(4), 700-703.
- Khanda CM, Mohapatra AK, Misra PK (1999) Effect of nitrogen and phosphorus fertilization on growth, yield and nutrient uptake of winter rice bean (*Vigna umbellata*). *Indian Journal of Agronomy* **44**(4), 791-794.
- Khanda CM, Mohapatra AK, Misra PK (2001) Response of rice bean (*Vigna umbellata*) to row spacing and phosphorus under rainfed condition. *Annals of Agricultural Research* **22**(4), 481-484.
- Khatounian CA (1993) Rice beans. In 'Feijão-arroz. Vol. 16'. pp. 10 pp.)
- Khatri NK, Nanda US, Kakani RK (2003) Extent of yellow mosaic virus disease of moth bean in Bikaner Region. In '' pp. 432-434. (Scientific Publishers (India): Jodhpur)
- Khatri RS (2004) Breeding priorities for genetic improvement in mothbean [*Vigna aconitifolia* (Jacq.) Marechal]. *Annals of Biology* **20**(1), 111-113.
- Khokhar S, Chauhan BM (1986a) Antinutritional factors in moth bean (*Vigna aconitifolia*): varietal differences and effects of methods of domestic processing and cooking. *Journal of Food Science* **51**(3), 591-594.
- Khokhar S, Chauhan BM (1986b) Effect of domestic processing and cooking on in vitro protein digestibility of moth bean. *Journal of Food Science* **51**(4), 1083-1084.
- Kiehn FA, Reimer M (1993) Alternative crops for the prairies. 46 pp.
- Kiessling D, Skiebe K (1987) Studies on the causes of variability in incomplete resistance of dwarf bean, *Phaseolus vulgaris* L., to *Pseudomonas syringae* pv. *phaseolicola* (Burkholder) Young, Dye et Wilkie. *Archiv für Züchtungsforschung* **17**(3), 191-200. [In German]
- Kikuchi K, Koizumi M, Ishida N, Kano H (2006) [Water uptake by dry beans observed by micro-magnetic resonance imaging](#). *Annals of Botany* **98**(3), 545-553.
- Kim JH, Delauney AJ, Verma DPS (1995) [Control of de novo purine biosynthesis genes in ureide-producing legumes: induction of glutamine phosphoribosylpyrophosphate amidotransferase gene and characterization of its cDNA from soybean and \*Vigna\*](#). *Plant Journal* **7**(1), 77-86.
- Kim KC, Choi HS (1987) Effects of temperature on the oviposition, feeding and emergence of the azuki bean weevil (*Callosobruchus chinensis* L.) in the stored beans. *Korean Journal of Plant Protection* **26**(2), 71-81. [In Korean]
- Kim S, Lee Y, Lee S, Moon J, Kim H, Hong E, Kim Y (1998) A new adzuki bean variety "Chilbopat" with large black seed and good seed quality. *RDA Journal of Crop Science (II)* **40**(2), 112-116. [In Korean]
- Kimura S, Laosinchai W, Itoh T, Cui X, Linder CR, Brown RM, Jr. (1999) Immunogold labelling of rosette terminal cellulose-synthesizing complexes in the vascular plant *Vigna angularis*. *Plant Cell* **11**(11), 2075-2085.
- Kimura S, Tokumaru S, Kuge K (2009) *Eremothecium coryli* and *E. ashbyi* cause yeast spot of azuki bean. *Journal of General Plant Pathology* **75**(4), 322-324.
- Kimura SD, Schmidtko K, Tajima R, Yoshida K, Nakashima H, Rauber R (2004) [Seasonal N uptake and N2 fixation by common and adzuki bean at various spacings](#). *Plant and Soil* **258**(1/2), 91-101.
- Kingsolver JM (1999) A new species of *Callosobruchus* (Coleoptera : Bruchidae) from Thailand and China. *Entomological News* **110**(4), 209-213.
- Kishor PBK, Hong Z, Miao G, Hu CAA, Verma DPS (1995a) Overexpression of  $\Delta 1$ -pyrroline-5-carboxylate synthetase increases proline production and confers osmotolerance in transgenic plants. *Plant Physiology* **108**(4), 1387-1394.
- Kishor PBK, Hong ZL, Miao GH, Hu CAA, Verma DPS (1995b) Overexpression of  $\Delta 1$ -pyrroline-5-carboxylate synthetase increases proline production and confers osmotolerance in transgenic plants. *Rice Biotechnology Quarterly* **23**, 21-22.
- Kitazawa K, Suzui T, Yanagita K (1979) Pathogenicity of *Phytophthora vignae* Purss to adzuki-bean and cowpea. *Annals of the Phytopathological Society of Japan* **45**(3), 406-408.
- Kitazawa K, Yanagita K (1984) Adzuki bean wilt caused by *Fusarium oxysporum* Schl.: re-occurrence and confirmation of the causal organism. *Annals of the Phytopathological Society of Japan* **50**(5), 643-645.
- Kitazawa K, Yanagita K (1989) *Fusarium oxysporum* Schl. f.sp. adzukicola n.f.sp., a wilt fungus of *Phaseolus angularis*. *Annals of the Phytopathological Society of Japan* **55**(1), 76-78.

- Kitou M, Yoshida S (1997) Effect of coffee residue on the growth of several crop species. *Journal of Weed Science and Technology* **42**(1), 25-30.
- Klomklao S, Benjakul S, Kishimura H, Osako K, Tanaka M (2010) A heat-stable trypsin inhibitor in adzuki bean (*Vigna angularis*): effect of extraction media, purification and biochemical characteristics. *International Journal of Food Science & Technology* **45**(1), 163-169.
- Kobayashi K (1984) Studies on induced mutants of *Cephalosporium gregatum*, the causal fungus of brown stem rot in adzuki beans. *Annals of the Phytopathological Society of Japan* **50**(4), 522-527.
- Kobayashi K, Hara Y (2004) Effects of the electrolyzed water on the color tone of sekihan. *Nippon Shokuhin Kagaku Kogaku Kaishi = Journal of the Japanese Society for Food Science and Technology* **51**(7), 358-366.
- Kobayashi K, Kondo N, Negishi H, Ui T (1984) Studies on the incidence and distribution of brown stem rot of adzuki beans in Hokkaido. *Memoirs of the Faculty of Agriculture Hokkaido University* **14**(2), 91-97.
- Kobayashi K, Kondo N, Tanaka H, Ui T (1980) Differences in the virulence of type A and type B isolates of *Cephalosporium gregatum*. *Annals of the Phytopathological Society of Japan* **46**(2), 253-255.
- Kobayashi K, Kondo N, Ui T (1981) Appearance of a mutant from *Cephalosporium gregatum* type A isolate in culture. *Annals of the Phytopathological Society of Japan* **47**(4), 577-580.
- Kobayashi K, Kondo N, Ui T, Tachibana H, Aota T (1983) Differences in pathogenicity of *Phialophora gregata* isolates from adzuki bean in Japan and from soybean in the United States. *Plant Disease* **67**(4), 387-388.
- Kobayashi K, Tanaka F, Kondo N, Ui T (1981) A selective medium for isolation of *Cephalosporium gregatum* from soil and populations in Adzuki bean field soils estimated with the medium. *Annals of the Phytopathological Society of Japan* **47**(1), 29-34.
- Kobayashi K, Tanaka F, Ui T, Akai J (1979) A new strain of *Cephalosporium gregatum*, causal fungus of brown stem rot of adzuki beans. *Annals of the Phytopathological Society of Japan* **45**(3), 409-411.
- Kobayashi K, Ui T (1980a) Comparison of *Cephalosporium gregatum* isolates from adzuki bean in Japan and from soybean in USA. *Annals of the Phytopathological Society of Japan* **46**(2), 241-246.
- Kobayashi K, Ui T (1980b) Production of gregatins by an Iowa isolate of *Cephalosporium gregatum*, the cause of brown stem rot of soybeans in U.S.A. *Annals of the Phytopathological Society of Japan* **46**(2), 256-257.
- Kobayashi K, Yamamoto H, Negishi H, Ogoshi A (1991) Formae speciales differentiation of *Phialophora gregata* isolates from adzuki bean and soybean in Japan. *Annals of the Phytopathological Society of Japan* **57**(2), 225-231.
- Kobayashi M, Ohta Y (1983) Induction of stress metabolite formation in suspension cultures of *Vigna angularis*. *Phytochemistry* **22**(5), 1257-1261.
- Kodama A (1989) Karyotype analyses of chromosomes in eighteen species belonging to nine tribes in the Leguminosae. *Bulletin of the Hiroshima Agricultural College* **8**(4), 691-706.
- Koga N (2008) An energy balance under a conventional crop rotation system in northern Japan: perspectives on fuel ethanol production from sugar beet. *Agriculture, Ecosystems & Environment* **125**(1/4), 101-110.
- Koga N, Sawamoto T, Tsuruta H (2006) Life cycle inventory-based analysis of greenhouse gas emissions from arable land farming systems in Hokkaido, northern Japan. *Soil Science and Plant Nutrition* **52**(4), 564-574.
- Koga N, Tsuruta H, Tsuji H, Nakano H (2003) Fuel consumption-derived CO<sub>2</sub> emissions under conventional and reduced tillage cropping systems in northern Japan. *Agriculture, Ecosystems & Environment* **99**(1/3), 213-219.
- Kohli RK, Batish DR (1994b) Exhibition of allelopathy of *Parthenium hysterophorus* L. in agroecosystems. *Tropical Ecology* **35**(2), 295-307.
- Kohli RK, Kanwaljit K, Anita K Inhibition of seed germination of *Vigna umbellata* in response to *Eucalyptus globulus* Labill. allelochemicals. 1990, pp. 1292-1296
- Kohli RK, Rani D (1992b) Identification and bioefficacy of soil chemics of *Parthenium*. 196-198.
- Kojima M, Katagiri Y, Furukawa N, Mouri H, Ohnishi M, Ito S (1991) Sterollipids and sphingolipids in adzuki bean organs. *Research Bulletin of Obihiro University, I* **17**(3), 235-242.
- Kojima M, Kimura H, Ohnishi M, Fujino Y, Ito S (1991) Distribution and characterization of diglycosyldiacylglycerol isomers with different anomeric configuration in higher plants. *Phytochemistry* **30**(4), 1165-1168.
- Kojima M, Nishi S, Saito Y, Hironaka K, Koaze H, Maeda R (2007a) Drink containing adzuki polyphenol inhibits weight gain in female mice on high-fat diets. *Nippon Shokuhin Kagaku Kogaku Kaishi = Journal of the Japanese Society for Food Science and Technology* **54**(5), 229-232.

- Kojima M, Nishi S, Saito Y, Hironaka K, Koaze H, Maeda R (2007b) Effects of short-term and continuous administration of Adzuki polyphenol on blood glucose concentration. *Nippon Shokuhin Kagaku Kogaku Kaishi = Journal of the Japanese Society for Food Science and Technology* **54**(1), 50-53.
- Kojima M, Nishi S, Yamashita S, Saito Y, Maeda R (2006) Smaller increase in serum cholesterol level in rats fed an ethanol extract of Adzuki bean seeds. *Nippon Shokuhin Kagaku Kogaku Kaishi = Journal of the Japanese Society for Food Science and Technology* **53**(7), 380-385.
- Kojima M, Ohnishi M, Ito S, Fujino Y (1989) Characterization of acylmono-, mono-, di, tri-, and tetraglycosylsterol and saponin in adzuki bean (*Vigna angularis*) seeds. *Lipids* **24**(10), 849-853.
- Kojima M, Ohnisi M, Ito S (1991) The fatty acid compositions of seeds of the Leguminosae. *Research Bulletin of Obihiro University, I* **17**(3), 227-233.
- Kojima M, Sasaki S, Ohnishi M, Mano Y, Ito S (1990) Changes in molecular species of glycerolipids from developing *Vigna angularis* seeds. *Phytochemistry* **29**(7), 2091-2096.
- Kojima M, Seki K, Ohnishi M, Ito S (1990) Structure of novel glyceroglycolipids in adzuki bean (*Vigna angularis*) seeds. *Biochemistry and Cell Biology* **68**(1), 59-64.
- Kojima M, Seki T, Ohnishi M, Ito S (1991) [Two digalactosyldiacylglycerols with different anomeric configuration in leguminous seeds](#). *Journal of the Science of Food and Agriculture* **54**(1), 35-41.
- Kojima M, Shiraki H, Ohnishi M, Ito S (1990) Two diglycosyldiacylglycerol isomers in plant leaves, ferns, mosses and seaweeds. *Phytochemistry* **29**(4), 1161-1163.
- Kojima M, Suzuki H, Ohnishi M, Ito S (1990) Chemical characterization of lipids in suspension cell cultures derived from several organs of adzuki beans. *Research Bulletin of Obihiro University, I* **17**(1), 21-29.
- Kojima M, Suzuki H, Ohnishi M, Ito S (1998) Effects of growth temperature on lipids of Adzuki bean cells. *Phytochemistry* **47**(8), 1483-1487.
- Kojima M, Yamashita S, Nishi S, Saito Y, Maeda R (2006) Antioxidative effect and liver protective action of Adzuki polyphenol. *Nippon Shokuhin Kagaku Kogaku Kaishi = Journal of the Japanese Society for Food Science and Technology* **53**(7), 386-392.
- Kokubo A, Yamamoto R, Masuda Y (1993) [Galactose prevents proton excretion but not IAA-induced growth in segments of azuki bean epicotyls](#). *Planta* **190**(2), 284-287.
- Kole C, Panigrahi J (2001b) *Vigna glabrescens* - a natural allopolyploid of *V. radiata* and *V. umbellata*: evidences from seed protein electrophoresis. *Crop Research (Hisar)* **22**(1), 121-128.
- Konarev AV, Tomooka N, Ishimoto M, Vaughan A (1999) Variability of the inhibitors of serine, cysteine proteinases and insect  $\alpha$ -amylases in Vigna and Phaseolus. 173-181.
- Konarev AV, Tomooka N, Vaughan DA (2002) [Proteinase inhibitor polymorphism in the genus Vigna subgenus Ceratotropis and its biosystematic implications](#). *Euphytica* **123**(2), 165-177.
- Kondo N (1995) Studies on adzuki bean wilt caused by *Fusarium oxysporum* f.sp. *adzukicola*. **19**(5), 411-472.
- Kondo N, Fujita S, Murata K, Ogoshi A (1998) Detection of two races of *Phialophora gregata* f.sp. *adzukicola*, the causal agent of adzuki bean brown stem rot. *Plant Disease* **82**(8), 928-930.
- Kondo N, Kobayashi K (1983a) Studies on the ecology of *Cephalosporium gregatum*, the causal fungus of brown stem rot and adzuki beans in Hokkaido. III. The longevity and germination of *Cephalosporium gregatum* conidia in soil. *Memoirs of the Faculty of Agriculture Hokkaido University* **14**(1), 50-55.
- Kondo N, Kobayashi K (1983b) Studies on the ecology of *Cephalosporium gregatum*, the causal fungus of brown stem rot of adzuki beans in Hokkaido. I. Seasonal variations of *Cephalosporium gregatum* populations in soil under host and non-host plant cultivation. *Memoirs of the Faculty of Agriculture Hokkaido University* **14**(1), 39-47.
- Kondo N, Kobayashi K (1983c) Studies on the ecology of *Cephalosporium gregatum*, the causal fungus of brown stem rot of adzuki beans in Hokkaido. II. Effects of infested host residue on the number of *Cephalosporium gregatum* propagules in soil in the field. *Memoirs of the Faculty of Agriculture Hokkaido University* **14**(1), 48-49.
- Kondo N, Kobayashi Y, Sakuma F, Fujita S, Murata K (2002) Regional distribution of two races of *Phialophora gregata* f. sp. *adzukicola*, causal agent of brown stem rot of adzuki bean, and their genetic diversity on Hokkaido, northernmost island of Japan. *Journal of General Plant Pathology* **68**(4), 284-291.
- Kondo N, Kodama F (1989) *Fusarium oxysporum* f.sp. *adzukicola*, causal agent of adzuki bean wilt, and detection of three races of the fungus. *Annals of the Phytopathological Society of Japan* **55**(4), 451-457.
- Kondo N, Kodama F, Ogoshi A (1997a) Survival of *Fusarium oxysporum* f.sp. *adzukicola* in residue of adzuki bean. *Annals of the Phytopathological Society of Japan* **63**(4), 334-336.

- Kondo N, Kodama F, Ogoshi A (1997b) Vegetative compatibility groups of *Fusarium oxysporum* f.sp. *adzukicola* and nonpathogenic *Fusarium oxysporum* on adzuki bean isolated from adzuki bean fields in Hokkaido. *Annals of the Phytopathological Society of Japan* **63**(1), 8-12.
- Kondo N, Nakazawa K, Fujita S, Shimada H, Naito S (2005) New virulent race of *Phialophora gregata* f. sp. *adzukicola* associated with continuous cultivation of adzuki bean cultivar Acc259. *Journal of General Plant Pathology* **71**(5), 360-363.
- Kondo N, Notsu A, Fujita S, Shimada H, Naito S (2005) DNA sequence analysis of ribosomal ITS region 1 of *Phytophthora vignae* f.sp. *adzukicola*, the pathogen that causes stem rot of adzuki bean. *Journal of General Plant Pathology* **71**(6), 414-417.
- Kondo N, Notsu A, Naito S, Fujita S, Shimada H (2004) Distribution of *Phytophthora vignae* f. sp. *adzukicola* races in adzuki bean fields in Hokkaido, Japan. *Plant Disease* **88**(8), 875-877.
- Kondo N, Shimada H, Fujita S (2009) Screening of cultivated and wild adzuki bean for resistance to race 3 of *Cadophora gregata* f. sp. *adzukicola*, cause of brown stem rot. *Journal of General Plant Pathology* **75**(3), 181-187.
- Kondo N, Shimada H, Kodama F (1994) Resistance gene loci to *Fusarium oxysporum* f.sp. *adzukicola* in adzuki bean cultivars Hatsune shozu and Toiku No. 123. *Annals of the Phytopathological Society of Japan* **60**(1), 107-108.
- Konno T (1980) Estimation of factors limiting the growth and yield of beans under different soil and weather conditions. *Research Bulletin of the Hokkaido National Agricultural Experiment Station*(126), 87-103.
- Kono S, Takashima T, Teshigawara S (2002b) A study on host plants and development of Mexican bean beetle *Epilachna varivestis* (Coleoptera: Coccinellidae) in Japan. *Research Bulletin of the Plant Protection Service, Japan* (38), 59-62.
- Kotaru M, Iwami K, Yeh HY, Ibuki F (1992) Occurrence in leguminous seeds, resistance to protease digestion and antigenicity of an  $\alpha$ -amylase inhibitor. *Food Chemistry* **43**(1), 47-50.
- Kothari PR, Kakani RK, Sharma MM (2005) Status of breeder seed production of arid legumes in Rajasthan Agricultural University. **2**(2), 270-273.
- Kothekar VS (1989) Differential radiation sensitivity in moth bean. *Current Science* **58**(13), 758-760.
- Kou KC, Kao CW, Leu LS (1988) The symptoms causal organism and fungicide screening for the control of grape cristulariella leaf spot. *Plant Protection Bulletin (Taichung)* **30**(2), 111-124.
- Krishna G, Mandal AB, Paliwal VK, Yadab KR (1989) Rice bean *Vigna-umbellata* as a feed for adult sheep. *Indian Journal of Animal Nutrition* **6**(4), 365-368.
- Krishna SK, Reddy SVK, Singh KD, Kumar PH, Chandrasekharao C, Krishnamurthy V (2009) Effect of organic and inorganic sources of nitrogen on productivity, quality and economics of FCV tobacco (*Nicotiana tabacum*). *Indian Journal of Agronomy* **54**(3), 336-341.
- Krishnamurthy KV, Godbole DA, Mascarenhas AF (1984) Studies on a drought resistant legume: the moth bean, *Vigna aconitifolia* (Jacq.) Marechal. I. Protoplast culture and organogenesis. *Plant Cell Reports* **3**(1), 30-32.
- Krishnan A, Krishna YSR, Sastri ASRAS (1980) Systems analysis approach for crop planning in Jodhpur district. *Indian Journal of Agricultural Sciences* **50**(5), 412-421.
- Krishnasamy T, Palaniappan J, Nagasamy N (2008) Pollen pistil interaction in inter-specific crosses of *Vigna* sp. *Cytologia* **73**(3), 251-257.
- Krishnaswamy L, Deshpande BD (1975) Acid patterns in germinating seed of *Phaseolus aconitifolius* Jacq. *Taiwania* **20**(2), 234-239.
- Kulkarni AS, Wankhede DB, Syed HM, Ismail S, Kajalwad BM, Sawate AR (2003) Studies on isolation and characterization of starch from two varieties of moth bean. In '' pp. 318-323. (Scientific Publishers (India): Jodhpur)
- Kulkarni H, Karadge BA (1991) Growth and mineral nutrition of moth bean (*Phaseolus aconitifolius* Jacq.) under saline conditions. *Indian Journal of Plant Physiology* **34**(1), 14-24.
- Kulkarni SG, Harode S, Deshpande AD, Borikar PS, Puri SN (1985) Damage and losses caused by *Callosobruchus chinensis* L. to different legumes stored in selected containers. *Agricultural Science Digest, India* **5**(2), 108-110.
- Kulkarni SG, Harode S, Desphande AD, Borkar PS, Puri SN (1988) Effect of containers and treatments on the storage of legumes. *Seed Research* **16**(1), 121-122.
- Kumar AS, Gamborg OL, Nabors MW (1988) Plant regeneration from cell suspension cultures of *Vigna aconitifolia*. *Plant Cell Reports* **7**(2), 138-141.
- Kumar BMD, Shambulingappa KG (1994) Stability analysis for seed yield and its components in rice bean (*Vigna umbellata* (Thunb.) Ohwi and Ohashi). *Legume Research* **17**(1), 47-52.

- Kumar BMD, Shambulingappa KG (1996) Path analysis in rice bean. *Legume Research* **19**(1), 19-22.
- Kumar BMD, Shambulingappa KG, Kulkarni RS (1996) Genetic divergence in rice bean. *Legume Research* **19**(1), 15-18.
- Kumar BMD, Shambulingappa KG, Kulkarni RS, Swamy GSK, Lohithaswa HS (1997) Analysis of genetic variability and character association in rice bean (*Vigna umbellata* Thunb. Ohwi and Ohashi). *Mysore Journal of Agricultural Sciences* **31**(1), 23-28.
- Kumar D (1998) Phenotypic stability for yield and related traits of moth bean in arid situations. *Indian Journal of Pulses Research* **11**(2), 30-34.
- Kumar D (2001) Phenotypic stability of quantitative traits in moth bean. *Indian Journal of Pulses Research* **14**(2), 116-118.
- Kumar D (2002) New plant type with unbranched growth habit in moth bean. *Indian Journal of Pulses Research* **15**(2), 179-180.
- Kumar D (2003) Management of abiotic stresses through resistance breeding. In '!' pp. 337-346. (Scientific Publishers (India): Jodhpur)
- Kumar D (2005) Status and direction of arid legumes research in India. *Indian Journal of Agricultural Sciences* **75**(7), 375-391.
- Kumar D, Rangaswamy NS (1984a) SEM studies on seed surface of wild and cultivated species of *Vigna Savi*. *Proceedings of the Indian Academy of Sciences, Plant Sciences* **93**(1), 35-42.
- Kumar D, Singh NB (2002) 'Moth bean in India.' (Scientific Publishers (India): Jodhpur) xvi + 157 pp.
- Kumar NP, Pandiyan M, Veerabadhiran P (2007a) Interspecific hybridization in *Vigna* species. *Plant Archives* **7**(1), 395-396.
- Kumar NP, Pandiyan M, Veerabadhiran P (2007c) Prefertilization barriers in *Vigna radiata* × *Vigna umbellata*. *Plant Archives* **7**(1), 377-380.
- Kumar S, Saxena SN, Jat RS, Sharma R (2008) Agrobacterium tumefaciens mediated genetic transformation of moth bean [*Vigna aconitifolia* (Jacq.) Marechal]. *Indian Journal of Genetics and Plant Breeding* **68**(3), 327-329.
- Kumar SHA, Uppar DS (2007) Influence of integrated nutrient management on seed yield and quality of mothbean [*Vigna aconitifolia* (Jacq.) Marchel]. *Karnataka Journal of Agricultural Sciences* **20**(2), 394-396.
- Kumar V, Gangwar B (1988) Effect of drainage on performance of grain and forage crops. *Journal of the Andaman Science Association* **4**(1), 74-76.
- Kumawat KC, Kumawat GL (1995) Effect of some insecticides on sucking pests and yellow mosaic incidence of mothbean, *Vigna aconitifolia* (Jacq.) Marechal. *Indian Journal of Virology* **11**(2), 69-72.
- Kumazaki M, Matsuyama S, Suzuki T, Kuwahara Y, Fujii K (2000) Parasitic wasp, *Dinarmus basalis*, utilizes oviposition-marking pheromone of host azuki bean weevils as host-recognizing kairomone. *Journal of Chemical Ecology* **26**(12), 2677-2695.
- Kumhar SR, Ramkrishna K (2006) Comparative symbiotic effectiveness of Bradyrhizobium (*Vigna aconitifolia*) on heterologous hosts. *Legume Research* **29**(1), 57-60.
- Kumhar SR, Ramkrishna K (2007) Symbiotic effectiveness of phage resistant mutants of mothbean *Rhizobium* on different host genetic systems. *Journal of Food Legumes* **20**(1), 83-86.
- Kundu TK, Virupaksha TK, Sudharshana L, Dana S (1993b) Lectins as phylogenetic markers of *Vigna* spp. *Biochemical Systematics and Ecology* **21**(4), 499-504.
- Kunjumon A, Kannan VR, Jasrai YT (1996) Plant regeneration from leaf explants of three grain legumes on same medium. *Journal of Plant Biochemistry and Biotechnology* **5**(1), 27-29.
- Kuno E, Kozai Y, Kubotsu K (1995) Modelling and analyzing density-dependent population processes: comparison between wild and laboratory strains of the bean weevil, *Callosobruchus chinensis* (L.). *Researches on Population Ecology* **37**(2), 165-176.
- Kuo KC, Lee CY, Chiang SS, Zheng YS (1999) Causal agent, pathogenesis and fungicide screening of lima bean anthracnose. *Plant Protection Bulletin (Taipei)* **41**(4), 265-275.
- Kurnar NP, Ano MM, VeRrOpOhiran R (2007) Prefertilization barriers in *Vigna radiata* x *Vigna umbellata*. *Plant Archives* **7**(1), 377-380.
- Kuya N, Kato M, Sato Y, Kaneta T, Sato S (2006) Comparative study of cellular structures implicated in gravisensing in statocytes of primary and lateral roots of *Vigna angularis*. *Protoplasma* **229**(1), 83-91.
- Köhler F, Cardon G, Pöhlman M, Gill R, Schieder O (1989) Enhancement of transformation rates in higher plants by low-dose irradiation: are DNA repair systems involved in the incorporation of exogenous DNA into the plant genome? *Plant Molecular Biology* **12**(2), 189-199.

- Köhler F, Golz C, Eapen S, Kohn H, Schieder O (1987a) Stable transformation of moth bean *Vigna aconitifolia* via direct gene transfer. *Plant Cell Reports* **6**(4), 313-317.
- Köhler F, Golz C, Eapen S, Schieder O (1987b) [Influence of plant cultivar and plasmid-DNA on transformation rates in tobacco and moth bean](#). *Plant Science, Irish Republic* **53**(1), 87-91.
- Lahiri AN (1975) Water use by desertic vegetation and its hydrological implications. *Annals of Arid Zone* **14**(2), 135-148.
- Lahiri AN (1976) Plant responses to soil-water status under arid conditions. 13-24.
- Lahiri AN (1980) Interaction of water stress and mineral nutrition on growth and yield. 341-352.
- Lahuta LB (2006) Biosynthesis of raffinose family oligosaccharides and galactosyl pinitols in developing and maturing seeds of winter vetch (*Vicia villosa* Roth.). *Acta Societatis Botanicorum Poloniae* **75**(3), 219-227.
- Lahuta LB, Gorecki RJ, Horbowicz M (2005) [High concentrations of D-pinitol or D-chiro-inositol inhibit the biosynthesis of raffinose family oligosaccharides in maturing smooth tare \(\*Vicia tetrasperma\* L. Schreb.\) seeds](#). *Acta Physiologiae Plantarum* **27**(4A), 505-513.
- Lahuta LB, Horbowicz M, Gojlo E, Goszcynska J, Gorecki RJ (2005) Exogenously applied D-pinitol and D-chiro-inositol modifies the accumulation of alpha-D-galactosides in developing tiny vetch (*Vicia hirsuta* l. SF Gray) seeds. *Acta Societatis Botanicorum Poloniae* **74**(4), 287-296.
- Lai CH, Yoshida T (1992) Population stability in laboratory ecosystem of stored product insects and parasitoids. *Chinese Journal of Entomology* **12**(3), 183-191.
- Lainson R, Shaw JJ, Naiff RD (1980) Chagas disease in the Amazon Basin Brazil speculations on transmission per os. *Revista do Instituto de Medicina Tropical de São Paulo* **22**(6), 294-297.
- Lairungreang C, Thavarasook C, Tomooka N (1991) Collection of wild Ceratotropis species in northern Thailand. 73-86.
- Lakra RK, Bhagat RK, Choubey S (2000) Effect of crop geometry and nutrient management in maize + ricebean intercropping system. *Journal of Research, Birsa Agricultural University* **12**(2), 157-161.
- Lakshmana D, Reddy BGS, Ramesh S (2007b) [Heterosis studies in rice bean \(\*Vigna umbellata\* \(Thunb.\) Ohwi and Ohashi\)](#). *Legume Research* **30**(3), 209-211.
- Lakshmana D, Reddy BGS, Ramesh S (2007c) [Studies on combining ability in rice bean](#). *Legume Research* **30**(4), 311-312.
- Lakshmi S, Devi LG, Nair MA, Vidya C (2002) Yield and economics of fodder legume hybrid napier intercropping systems. *Forage Research* **28**(1), 13-15.
- Lakshmi YS, Murthy VR (2001) Effect of pre-monsoon and monsoon intercrops on grain equivalents of sunflower in sequence cropping. *Crop Research (Hisar)* **22**(3), 404-407.
- Lakshmi YS, Rao PG, Krishnamurthy VR (2001a) Influence of pre-monsoon and monsoon intercrops with sunflower (*Helianthus annuus* L.) in sequence cropping. *Research on Crops* **2**(2), 148-155.
- Lakshmi YS, Rao PG, Murthy VR (2001b) Effect of pre-monsoon and monsoon intercrops on yield and yield components of sunflower (*Helianthus annuus*) in sequence cropping. *Annals of Agri Bio Research* **6**(2), 141-146.
- Lal A, Warber S, Kirakosyan A, Kaufman PB, Duke JA (2003) Upregulation of isoflavonoids and soluble proteins in edible legumes by light and fungal elicitor treatments. *Journal of Alternative and Complementary Medicine* **9**(3), 371-378.
- Lam-Sanchez A, Durigan JF, Campos SLd, Silvestre SR, Pedroso PAC, Banzatto DA (1990) Effect of planting date on chemical composition and physico-chemical characteristics of grain of cultivars of *Phaseolus vulgaris* L., *Vigna angularis* (Willd) Wright and *Vigna unguiculata* L. Walp. *Alimentos e Nutrição* **2**, 35-44. [In Portuguese]
- Landfeld A, Kyhos K, Strohalm J, Sestak J, Houska M (2006) Heat transfer with heating and cooling granular materials in a scraped-surface homogeniser. *Journal of Food Engineering* **77**(3), 708-712.
- Lang LJ (1991) Genetic resource work on important food legumes in China. 28-35.
- Laurel VB, Ramirez DA, Huelgas VC (1980b) Immunosuppressive chemical treatment in *Vigna radiata* l. Wilczek × *Vigna umbellata* (Thunb.) Ohwi and Ohashi. *Philippine Journal of Crop Science* **5**(4), 128-129.
- Laurena AC, Garcia VV, Mendoza EMT (1984) Localization and determination of condensed tannins in cowpea (*Vigna unguiculata* (L.) Walp.). *Philippine Agriculturist* **67**(3), 329-336.
- Laurena AC, Revilleza MJR, Mendona EMT (1994) [Polyphenols, Phytate, Cyanogenic Glycosides, and Trypsin Inhibitor Activity of Several Philippine Indigenous Food Legumes](#). *Journal of Food Composition and Analysis* **7**(3), 194-202.

- Laurena AC, Rodriguez FM, Sabino NG, Zamora AF, Mendoza EMT (1991) [Amino-acid-composition, relative nutritive-value and \*invitro\* protein digestibility of several philippine indigenous legumes](#). *Plant Foods for Human Nutrition* **41**(1), 59-68.
- Law H, Choi S, Ma C (2008) Study of conformation of vicilin from *Dolichos lablab* and *Phaseolus calcaratus* by Fourier-transform infrared spectroscopy and differential scanning calorimetry. *Food Research International* **41**(7), 720-729.
- Lawn RJ (1979a) [Agronomic studies on \*Vigna\* spp. in south-eastern Queensland. 1. Phenological response of cultivars to sowing date](#). *Australian Journal of Agricultural Research* **30**(5), 855-870.
- Lawn RJ (1979b) [Agronomic studies on \*Vigna\* spp. in south-eastern Queensland. 2. Vegetative and reproductive response of cultivars to sowing date](#). *Australian Journal of Agricultural Research* **30**(5), 871-882.
- Lawn RJ (1983) [Agronomic studies on \*Vigna\* spp. in South-Eastern Queensland. III. Response to sowing arrangement](#). *Australian Journal of Agricultural Research* **34**(5), 505-515.
- Lawn RJ, Bushby HVA (1982a) [Effect of root, shoot and Rhizobium strain on nitrogen fixation in four Asiatic \*Vigna\* species](#). *New Phytologist* **92**(3), 425-434.
- Lawn RJ, Summerfield RJ, Ellis RH, Qi A, Roberts EH, Chay PM, Brouwers JB, Rose JL, Yeates SJ (1995b) [Towards the reliable prediction of time to flowering in six annual crops. VI. Applications in crop improvement](#). *Experimental Agriculture* **31**(1), 89-108.
- Leach GJ, Rees MC, Charles-Edwards DA (1986a) [Relations between summer crops and ground cover legumes in a subtropical environment. 1. Effects of a \*Vigna trilobata\* ground cover on growth and yield of sorghum and sunflower](#). *Field Crops Research* **15**(1), 17-37.
- Leach GJ, Rees MC, Charles-Edwards DA (1986b) [Relations between summer crops and ground cover legumes in a subtropical environment. 2. Effects of sorghum and sunflower on the growth of \*Vigna trilobata\* and \*Medicago scutellata\*](#). *Field Crops Research* **15**(1), 39-55.
- Lee CK, Karunanithy R (1990) [Effects of germination on the chemical composition of \*Glycine\* and \*Phaseolus\* beans](#). *Journal of the Science of Food and Agriculture* **51**(4), 437-445.
- Lee H, Lee E, Rhee Y, Ahn K, Li GX, Jiang C, Lü JX, Kim S (2006) [An oriental herbal cocktail, Ka-mi-kae-kyuk-tang, exerts anti-cancer activities by targeting angiogenesis, apoptosis and metastasis](#). *Carcinogenesis* **27**(12), 2455-2463.
- Lee HB, Lee SY, Hong CK, Kim DY, Huh BL, Kim DR (1992) Studies on the intercropping system on the established sesame field. *Research Reports of the Rural Development Administration, Upland & Industrial Crops* **34**(1), 97-103. [In Korean]
- Lee K (2007) Comparison of weed occurrence and growth of some leguminous plants for green manure cover crop during summer fallow. *Korean Journal of Crop Science* **52**(2), 169-175.
- Lee M, Hwang G, Chen Y, Lin H, Wu C (2006) [Molecular cloning of Indian jujube \(\*Ziziphus mauritiana\*\) allergen Ziz m 1 with sequence similarity to plant class III chitinases](#). *Molecular Immunology* **43**(8), 1144-1151.
- Lee NG, Stein B, Suzuki H, Verma DPS (1993) [Expression of antisense nodulin-35 RNA in \*Vigna aconitifolia\* transgenic root nodules retards peroxisome development and affects nitrogen availability to the plant](#). *Plant Journal* **3**(4), 599-606.
- Lee S, Kim T, Kim B (1995) A simple test for evaluating acid rain tolerance in crops. *Korean Journal of Crop Science* **40**(1), 77-85. [In Korean]
- Lee SC (1989) Survey on the population dynamics and control of *Spodoptera litura* in legumes by using sex pheromone. *Chinese Journal of Entomology* **9**(1), 27-36.
- Lee T, Cheng H (2000) Application of Gray's theory to predict prices of agricultural products - the case of adzuki beans. *Journal of Agriculture and Forestry* **49**(2), 83-92.
- Lee WH (1993) Field studies and cultural practices for adzuki bean in Tainan area. *Taiwan Agriculture* **29**(1), 70-73.
- Lee WT, Ho CC, Lo KC (1990) Mass production of phytoseiids: I. Evaluation on eight host plants for the mass-rearing of *Tetranychus urticae* Koch and *T. kanzawai* Kishida (Acarina: Tetranychidae). *Journal of Agricultural Research of China* **39**(2), 121-132.
- Lee YH, Chin MS, Hong EH (1991) Effects of planting date on the flowering and maturing of adzuki bean (*Vigna angularis* (Willd.) Ohwi & Ohashi) and mungbean (*Vigna radiata* (L.) Wilczek). *Research Reports of the Rural Development Administration, Upland & Industrial Crops* **33**(3), 5-14. [In Korean]
- Lei BJ, Yin GC, Lu CH, Qian H, Zhang KW, Zhou SJ, Wang SL (1991) Study of direct transfer of exogenous DNA into soyabean. *Soybean Science* **10**(1), 58-63.
- Lekhu G, Singhvi PM (2004) Repellency power of some plant extract against *Callosobruchus maculatus* F. infesting moth bean seeds (*Vigna aconitifolia*). *Advances in Plant Sciences* **17**(2), 457-459.

- Lekhu G, Singhvi PM, Menakshi J, Mohita M (2005) Evaluation of certain plant extracts on adult emergence of *Callosobruchus maculatus* infesting, moth bean (*Vigna aconitifolia*). *Journal of Applied Zoological Researches* **16**(2), 203-205.
- Lenzi E, Rauber T, Luchese EB, Galli D, Toledo EA (1990) determination of cadmium in materials consumed by mankind. *Arquivos De Biologia E Tecnologia* **33**(1), 179-194.
- Li B (2005) Application of Qiangjinliniao Decoction in the treatment of acute nephritis in children. *Chinese Journal of Information on Traditional Chinese Medicine* **12**(9), 55-55.
- Li B, Zhao Q, Weber N, Lou Y, Proksch P (2005) The protective effects of total flavonoid extract of Phaseolus angularis Wight on oxidative damages of rat primary hepatocytes in vitro. *Acta Nutrimenta Sinica* **27**(5), 397-400, 405.
- Li Z, Huang C, Zhang X, Cao M, Li Z, Wang G, Wu Z (2005) Studies on transferring P5CS gene into tall fescue (*Festuca arundinacea* Schreb.) via microprojectile bombardment. *Acta Horticulturae Sinica* **32**(4), 653-657.
- Liang HQ, An J, Wu XR (1988) Biochemical analysis of the quality of legume seed proteins. *Acta Agronomica Sinica* **14**(2), 117-123.
- Lin C, Wu S, Wang J, Yang J, Chang C (2001) Evaluation of the antioxidant activity of legumes. *Pharmaceutical Biology* **39**(4), 300-304.
- Lin J, Horng S (2003) Effect of larval density and host on interspecific competition between *Callosobruchus chinensis* and *C. maculatus* (Coleoptera: Bruchidae). *Formosan Entomologist* **23**(1), 13-25.
- Lin P, Lai H (2006) Bioactive compounds in legumes and their germinated products. *Journal of Agricultural and Food Chemistry* **54**(11), 3807-3814.
- Lin ZW, Miao GH, Verma DPS (1995) A cDNA sequence encoding glutamine synthetase is preferentially expressed in nodules of *Vigna aconitifolia*. *Plant Physiology* **107**(1), 279-280.
- Linden JC, Stoner RR (2005) Proprietary elicitor affects seed germination and delays fruit senescence. *Journal of Food, Agriculture & Environment* **3**(3/4), 184-189.
- Liu F, Zhang Y, Jia R, Ren X (2009) Effect of seed soaking with La (NO<sub>3</sub>)<sub>3</sub> on root and seedling growth of adzuki bean under different concentrations NaCl stresses. *Journal of Wuhan Botanical Research* **27**(4), 397-402.
- Liu H (2006) The application of integrated Traditional Chinese Medicine and western medicine in the treatment of Henoch-Schonlein purpura (HSP) in children. *Chinese Journal of Information on Traditional Chinese Medicine* **13**(7), 62-62.
- Liu H, Probst A, Liao B (2005) Metal contamination of soils and crops affected by the Chenzhou lead/zinc mine spill (Hunan, China). *Science of the Total Environment* **339**(1/3), 153-166.
- Liu HQ Present status of the occurrence, injury, and research of soyabean cyst nematodes in Heilongjiang Province. *Soybean Science* **6**(2), 141-150.
- Liu Q, Kuang Y, Zheng Y (1994) Studies on the radiosensitivity of edible vegetable seeds. *Journal of South China Agricultural University* **15**(4), 85-91.
- Liu SD (1991) Biological control of adzuki bean root rot disease caused by Rhizoctonia solani. *Plant Protection Bulletin (Taipei)* **33**(1), 63-71.
- Liu SD (1992) The development and application of an antagonistic fungus as a biocontrol agent against root rot disease of adzuki bean. *Taiwan Agriculture Bimonthly* **28**(3), 87-93.
- Lodha S, Mahander S (1985) Quantitative determination of Macrophomina phaseolina (Tassi) Goid in grass-legume intercropping systems. *Annals of Arid Zone* **23**(3), 259-261.
- Lokesh, Verma PK, Gupta SN, Behl RK (2003b) Path coefficient analysis for various morphological and quality parameters in rice bean [*Vigna umbellata* (Thunb.) Ohwi and Ohashi]. *National Journal of Plant Improvement* **5**(2), 116-119.
- Lokesh, Verma PK, Gupta SN, Deen MK, Behl RK (2003d) Association between morphological and quality parameters in rice bean [*Vigna umbellata* (Thunb.) Ohwi and Ohashi]. *National Journal of Plant Improvement* **5**(1), 54-57.
- Lokesh, Verma PK, Sangwan O, Behl RK (2005a) Association analysis between seed yield and its component characters in rice bean. *Annals of Biology* **21**(2), 165-168.
- Lokesh, Verma PK, Sangwan O, Behl RK (2005b) Genotypic divergence analysis for different characters in rice bean [*Vigna umbellata* (Thunb.) Ohwi and Ohashi]. *Annals of Biology* **21**(2), 159-164.
- Lokesh R, Veeresh LC, Shivashankar G (1990) Rice bean - a potential legume for Karnataka. *Current Research - University of Agricultural Sciences (Bangalore)* **19**(11), 185-187.
- Lokesha R, Veeresh LC (1993) Induced mutagenesis and genetic improvement of rice bean [*Vigna umbellata* (Thunb.) Ohwi and Ohashi]. *Legume Research* **16**(1-2), 37-40.

- Long S, Qian Y (1997) A comparative study on the effects of some black and light-coloured natural foods on scavenging active oxygen species. *Acta Nutrimenta Sinica* **19**(4), 470-473.
- Lubis I, Lubis SHA, Sastrapradja D, Sastrapradja S (1977) Variation in seed protein of *Vigna unguiculata* (L.) Walp). *Annales Bogorienses* **6**(3), 125-132.
- Lumpkin TA, Konovsky JC, Larson KJ, McClary DC (1993) Potential new specialty crops from Asia: azuki bean, edamame soybean, and astragalus. 45-51.
- Lumpkin TA, McClary DC (1994) 'Azuki bean: botany, production and uses.' (CAB International: Wallingford) x + 268 pp.
- Lutova MI, Buravtseva TV, Kamentseva IE, Sharkova VE (1998) Heat tolerance and cell repair capacity in genotypes of *Phaseolus* and *Vigna* of different geographic origin. *Fiziologiya Rastenii (Moscow)* **45**(5), 703-708.
- López A R, Yamada S, Yamanouchi M (1999) Comparison of sodium uptake by and transport in detached plant parts among several crops. *Soil Science and Plant Nutrition* **45**(3), 659-668.
- Mabesa LB, Atutubo EO, Castro-Sandoval E (1982) Protein quality of flours from germinated legumes. *Philippine Agriculturist* **65**(3), 245-252.
- Mabesa LB, Novero FE (1985) Baking qualities of flour from germinated and ungerminated legumes. *Philippine Journal of Crop Science* **10**(2), 63-70.
- Machado M, Tai W, Baker LR (1982a) Cytogenetic analysis of the interspecific hybrid *Vigna radiata* X *V. umbellata*. *Journal of Heredity* **73**(3), 205-208.
- Maciel FL, Gerald LTS, Echeverrigaray S (2001) Random amplified polymorphic DNA (RAPD) markers variability among cultivars and landraces of common beans (*Phaseolus vulgaris* L.) of south-Brazil. *Euphytica* **120**(2), 257-263.
- MacKenzie DR, Ho L, Liu TD, Wu HBF, Oyer EB (1975) Photoperiodism of mung bean and four related species. *HortScience* **10**(5), 486-487.
- Madal N, Dana I (1998) Genetic divergence and nature of heterosis in ricebean. *Annals of Agricultural Research* **19**(1), 18-21.
- Maggio S, Malhotra SP, Dhawan K, Singh R (1999) Purification and characterization of protease inhibitor from rice bean (*Vigna umbellata* T.) seeds. *Journal of Plant Biochemistry and Biotechnology* **8**(1), 61-65.
- Mahabeer S (2007) Biocontrol of root rot disease of mothbean caused by *Macrophomina phaseolina* (Tassi) Goid. *Journal of Food Legumes* **20**(2), 196-197.
- Mahabeer S, Mehta SM (2005) Compatibility of seed protectant chemicals with mothbean *Rhizobium* sp. in vitro. *Plant Disease Research (Ludhiana)* **20**(2), 169-170.
- Mahajan V, Singh J, Sarma BK, Patel CS (1992) Stability for light reception in intergenotypic competition of maize and rice bean. *Indian Journal of Hill Farming* **5**(2), 119-122.
- Mahander S, Singh RP (1986) Effect of fodder and grain legume intercrops on the quality of *Cenchrus ciliaris* Linn. *Indian Journal of Agricultural Sciences* **56**(10), 704-709.
- Mahna SK, Rekha G, Parvateesam M (1993) Genetic effect of sodium azide and hydroxylamine on moth bean (*Vigna aconitifolia*). *Indian Journal of Agricultural Sciences* **63**(9), 551-555.
- Maia FMM, Oliveira JTA, Matos MRT, Moreira RA, Vasconcelos IM (2000) Proximate composition, amino acid content and haemagglutinating and trypsin-inhibiting activities of some Brazilian *Vigna unguiculata* (L.) Walp cultivars. *Journal of the Science of Food and Agriculture* **80**(4), 453-458.
- Maiti S, Mukherjee AK, Nanda MK, Bhattacharya S (2006) Effect of levels of phosphorus and dates of sowing on the productivity of ricebean when grown as a grain legume. *Journal of Crop and Weed* **2**(2), 37-39.
- Major S, Sarma BK (1993) Natural outcrossing in rice bean (*Vigna umbellata*). *Indian Journal of Agricultural Sciences* **63**(8), 520-521.
- Majumdar VL (1997) *Vigna aconitifolia*-a new host for *Curvularia tuberculata*. *Indian Phytopathology* **50**(2), 300.
- Majumdar VL, Jat JR, Gour HN (1996) Effects of biocontrol agents on the growth of *Macrophomina phaseolina*, the incitant of blight of moth bean. *Indian Journal of Mycology and Plant Pathology* **26**(2), 202-203.
- Majumdar VL, Rathore GS (1993) New records of fungi causing leaf spots of moth bean (*Vigna aconitifolia*). *Indian Phytopathology* **46**(2), 195-196.
- Malabadi RB (2002) Plant regeneration from *in vitro* cultured leaf in mothbean. *Journal of Phytological Research* **15**(2), 137-140.
- Malabadi RB, Nataraja K (2002) Large scale production and storability of encapsulated somatic embryos of mothbean (*Vigna aconitifolia* Jacq.). *Journal of Plant Biochemistry and Biotechnology* **11**(1), 61-64.
- Malabadi RB, Nataraja K (2003) Somatic embryogenesis and biochemical analysis of *in vitro*-derived plants of mothbean (*Vigna aconitifolia* Jacq.). *Plant Cell Biotechnology and Molecular Biology* **4**(1/2), 69-74.

- Malabadi RB, Nataraja K (2007) Agrobacterium tumefaciens-mediated genetic transformation of *Vigna aconitifolia* and stable transmission of the genes to somatic seedlings. *International Journal of Agricultural Research* **2**(5), 450-458.
- Malathi VG, Surendranath B, Naghma A, Roy A (2005) Adaptation to new hosts shown by the cloned components of Mungbean yellow mosaic India virus causing cowpea golden mosaic in northern India. *Canadian Journal of Plant Pathology* **27**(3), 439-447.
- Malhotra S, Malik D, Dhindsa KS (1988) Proximate composition and antinutritional factors in rice bean (*Vigna umbellata*). *Plant Foods for Human Nutrition* **38**(1), 75-81.
- Mali PC, Uday B, Satish L (2000) Effect of planting dates and development of yellow mosaic virus on biochemical constituents of moth bean genotypes. *Indian Phytopathology* **53**(4), 379-383.
- Malik G, Dawar S, Sattar A, Dawar A (2005) Efficacy of *Trichoderma harzianum* after multiplication on different substrates in the control of root rot fungi. *International Journal of Biology and Biotechnology* **2**(1), 237-242.
- Malik MA, Saleem MF, Cheema MA (2002) Substitution of nitrogen requirement of wheat (*Triticum aestivum* L.) through green manuring. *International Journal of Agriculture and Biology* **4**(1), 145-147.
- Malik MA, Shah SH, Saleem MF, Jami MS (2002b) Determining suitable stand density and phosphorus level for improving productivity of rice bean (*Vigna umbellata*). *International Journal of Agriculture and Biology* **4**(1), 120-122.
- Malini B, Sarma BK, Annadurai A, Verma DK (2001) Performance of rice bean (*Vigna umbellata*) genotypes in mid-altitude conditions of Meghalaya. *Indian Journal of Hill Farming* **14**(1), 150-152.
- Mandal BK, Dhara MC, Mandal BB, Bhunia SR, Dandapat A (1991) Nodulation in some legumes grown as pure and intercrops. *Indian Agriculturist* **35**(1), 15-19.
- Mandal BK, Dhara MC, Mandal BB, Das SK, Nandy R (1990a) Rice, mungbean, soyabean, peanut, ricebean, and blackgram yields under different intercropping systems. *Agronomy Journal* **82**(6), 1063-1066.
- Mandal BK, Dhara MC, Mandal BB, Patra BC (1989b) Growth of rice, mungbean, soybean, peanut, rice bean and blackgram as sole and intercrops. *Indian Journal of Plant Physiology* **32**(4), 294-298.
- Mandal BK, Rao MV (1983) Effect of cattleshed washings on the yield of some fodder crops. *Indian Journal of Agronomy* **28**(4), 475-476.
- Mandal N, Dana I (1998) Genetic variability and path coefficient analysis in ricebean. *Annals of Agricultural Research* **19**(1), 107-109.
- Mandal S, Saha H, Hajra JN (2003) Preparation of phosphocompost from paddy straw and water hyacinth and its evaluation on paddy and rice bean. *Environment and Ecology* **21**(2), 267-271.
- Mandal SMA (2005) Preliminary field evaluation of ricebean varieties against pod borers. *Insect Environment* **11**(1), 47-48.
- Mandal SR, Mukherjee AK (1999b) Forage yield and qualities of some ricebean (*Vigna umbellata*) germplasms. *Environment and Ecology* **17**(3), 749-751.
- Mandal SR, Mukherjee AK, Pal S, Mandal NN (2002) Comparative response of ricebean and cowpea to S-application in coastal saline zone. *Journal of Interacademia* **6**(1), 21-24.
- Mandal SR, Mukherjee AK, Patra BC (2001) Effect of liming on soil fertility and forage yield and quality of ricebean in acid oxisol. *Environment and Ecology* **19**(4), 771-773.
- Mandeep K, Mehta U (1994) Effect of milling on nutritional quality and chemical characteristics of ricebean. *Indian Journal of Pulses Research* **7**(1), 41-44.
- Mandeep K, Urmil M (1993) Effect of partial and full substitution of Bengalgram and blackgram Dhals with ricebean on essential amino acids and mineral contents of vada and pakora. *Journal of Food Science and Technology (Mysore)* **30**(6), 454-456.
- Mangalekar A, Sontakke S (1989) Partial purification and some properties of nucleolytic enzymes from moth bean (*Phaseolus aconitifolius*). *Indian Journal of Plant Physiology* **32**(4), 336-343.
- Manguiat IJ, Mendoza DM, Tilo SN (1985) Influence of the legume-rhizobia symbiosis on the nitrogen economy of a legume-based cropping system. *Philippine Agriculturist* **68**(1), 44-55.
- Manguiat IJ, Sebiano AG, Guinto DF, Jalalon AT Comparison of *Sesbania rostrata* and *Phaseolus calcaratus* as biofertilizer for corn. 1988, p. S8
- Manguiat IJ, Singleton PW, Rocamora PM, Calo MU, Taleon EE (1997) Effectiveness of *Sesbania rostrata* and *Phaseolus calcaratus* as green manure for upland rice grown in acidic soil. *Plant and Soil* **192**(2), 321-331.
- Manigopa C, Hanif K, Ramkrishna K (2003) Symbiotic properties of hypermotile mutants of *Bradyrhizobium* sp. (*Vigna aconitifolia*). *Indian Journal of Microbiology* **43**(2), 101-105.

- Manigopa C, Ramkrishna K (2006) Field evaluation of cowpea *Rhizobium* mutants for symbiotic effectiveness with mothbean varieties. *Indian Journal of Pulses Research* **19**(2), 210-212.
- Manish P, Tarafdar JC (1999) Critical level of VAM fungal propagules in different arid crops. *Agrochimica* **43**(3/4), 187-192.
- Manju S, Kumar D (2005) Mutants with high productivity induced in mothbean. *Indian Journal of Pulses Research* **18**(2), 244-245.
- Manju S, Seema G, Neetu R, Bhati PS, Kumar D (2003) Induced polygenic variability in moth bean. In '!' pp. 79-83. (Scientific Publishers (India): Jodhpur)
- Manjunath TS, Thimmegowda S, Suryanarayana G, Devakumar N (1996) Response of ricebean to N-nutrition - a modelling approach. *Indian Agriculturist* **40**(4), 247-252.
- Manjunatha M, Bhat NS, Raju GTT, Viraktamath CA (1988) [A new predatory ant \*Aenictus pachycerus\* Smith \(Hymenoptera: Formicidae\) on wireworm \*Agriotes\* sp. \(Coleoptera: Elateridae\)](#). *Current Science* **57**(13), 744.
- Mann HS (1981) Management of arid-land resources for dryland and irrigated crops. 479-493.
- Manning W, Fleming J, Storrie A, Cook T (2009) Weed control in summer crops, 2008-09. iv + 56 pp.
- Mano H, Toquenaga Y, Fujii K (2007) [Scramble-like contest competition in \*Callosobruchus analis\* \(Coleoptera: Bruchidae\)](#). *Journal of Stored Products Research* **43**(3), 211-220.
- Manohar C, Bhattacharya NM (2005a) Association of green forage yield attributes in rice bean *Vigna umbellata* (Thunb.) Ohwi and Ohashi. *Environment and Ecology* **23**(3), 538-540.
- Manohar C, Bhattacharya NM (2005b) Genetic variability in rice bean. *Environment and Ecology* **23**(3), 534-537.
- Manpreet K, Kawatra BL (1999) Molar ratios to predict zinc availability from domestically processed ricebean (*Vigna umbellata*). *Journal of Food Science and Technology (Mysore)* **36**(6), 548-550.
- Manpreet K, Kawatra BL (2002a) [Effect of domestic processing on zinc bioavailability from ricebean \(\*Vigna umbellata\*\) diets](#). *Plant Foods for Human Nutrition* **57**(3/4), 307-318.
- Manpreet K, Kawatra BL (2002b) [Evaluation of iron bioavailability from ricebean \(\*Vigna umbellata\*\) by using anaemic rats](#). *Nutrition Research* **22**(5), 633-640.
- Manyata M, Gajja BL (2004) Analysis of growth of pulses in arid zone of Rajasthan. *Current Agriculture* **28**(1/2), 63-68.
- Mao G (2003) The application of Yiqifuzhengzhilie Decoction in the treatment of chronic prostatitis. *Chinese Journal of Information on Traditional Chinese Medicine* **10**(8), 45-45.
- Mar S, Mori H, Lee J, Fukuda K, Saburi W, Fukuhara A, Okuyama M, Chiba S, Kimura A (2003) Purification, characterization, and sequence analysis of two  $\alpha$ -amylase isoforms from azuki bean, *Vigna angularis*, showing different affinity towards  $\beta$ -cyclodextrin Sepharose. *Bioscience, Biotechnology and Biochemistry* **67**(5), 1080-1093.
- Marappa N (2008b) Screening of mungbean genotypes and its wild relatives for resistant sources to Cercospora leaf spot disease. *Asian Journal of Bio Science* **3**(2), 324-326.
- Marappa N, Savithramma DL, Prabuddha HR, Jayesh KC (2008) Genetic variability study in mungbean and related species for yield and its attributes. *Research on Crops* **9**(2), 364-366.
- Marcelo AV, Cora JE, Fernandes C, Martins MD, Jorge RF (2009) Crop sequences in no-tillage system: effects on soil fertility and soybean, maize and rice yield. *Revista Brasileira De Ciencia Do Solo* **33**(2), 417-428.
- Marium T, Shahnaz D, Mehdi FS, Zaki MJ (2006) Use of Avicennia marina in the control of root infecting fungi on okra and mash bean. *Pakistan Journal of Botany* **38**(3), 811-815.
- Marium T, Shahnaz D, Mehdi FS, Zaki MJ (2007) Use of Avicennia marina (Forsk.) Vierh in the control of root knot nematode Meloidogyne javanica (Treub) Chitwood on okra and mash bean. *Turkish Journal of Biology* **31**(4), 225-230.
- Maruyama C, Araki R, Kawamura M, Kondo N, Kigawa M, Kawai Y, Takanami Y, Miyashita K, Shimomitsu T (2008) Azuki bean juice lowers serum triglyceride concentrations in healthy young women. *Journal of Clinical Biochemistry and Nutrition* **43**(1), 19-25.
- Masuda Y (1980) Auxin-induced changes in noncellulosic polysaccharides of cell walls of monocot coleoptiles and dicot stems. 79-89.
- Matai S, Bagchi DK (1974) Some promising legumes for leaf protein extraction. *Science & Culture* **40**(1), 34-36.
- Mathur AK, Tyagi RNS (1982) Symptomatology of *Colletotrichum truncatum* leaf spot of moth bean. *Indian Phytopathology* **35**(1), 135.
- Mathur AK, Tyagi RNS (1984) Occurrence of Choanephora pod rot on kharif pulses in Rajasthan. *Indian Journal of Mycology and Plant Pathology* **14**(2), 152.

- Mathur AK, Tyagi RNS (1986) Reaction of moth bean cultivars to bacterial blight in Rajasthan. *Indian Journal of Mycology and Plant Pathology* **16**(3), 322.
- Mathur BK, Henry A (2005b) Compound growth rate of area, production and productivity of pulses (moth bean, cowpea, and mung bean) in arid districts of Rajasthan. **2**(1), 50-53. (Scientific Publishers (India): Jodhpur)
- Mathur BK, Henry A (2006) Performance of grain legumes v/s oilseeds and cereals for area, production and productivity in different agro-climatic zones of western Rajasthan. *Journal of Arid Legumes* **3**(1), 83-92.
- Mathur BK, Patel AK, Bohra HC, Mathur AC, Kaushish SK (2005b) Acceptability, palatability and utilization of clusterbean and dewbean fodder v/s buffel grass in sheep. **2**(1), 39-42. (Scientific Publishers (India): Jodhpur)
- Mathur SR, Shukla KB, Sharma MK (2006) Effect of cadmium on seedling growth, lipid peroxidation and photosynthetic pigments of mothbean cultivar. *International Journal of Plant Sciences (Muzaffarnagar)* **1**(2), 200-201.
- Matsuda T, Chonan N (1989) Ultrastructure of seed coats in pulse crops. I. The correlation between cuticular wax and gloss of seed coat. *Scientific Reports of the Faculty of Agriculture, Ibaraki University*(37), 1-9.
- Matsumoto H, Nishi S, Ishizuka K (1991) Action of thiocarbamate herbicide dimepiperate on plant metabolism. *Weed Research (Tokyo)* **36**(3), 257-265.
- Matsumoto H, Tebayashi S, Kuwahara Y, Matsuyama S, Suzuki T, Fujii K (1994) Identification of taxifolin present in the azuki bean as an oviposition stimulant of the azuki bean weevil. *Journal of Pesticide Science* **19**(3), 181-186.
- Matsumoto N, Yoshikawa M, Eda K, Kobayashi A, Yokoshima M, Murakami M, Kanekita H (2008) Simple preprocessing method for multi-determination of 235 pesticide residues in cooked ingredients of foods by GC/MS and LC/MS/MS. *Shokuhin Eiseigaku Zasshi = Journal of the Food Hygienic Society of Japan* **49**(3), 211-222.
- Matsumoto T, Inoue M (1989) On a virus disease of azuki-bean plants occurring on Hachirogata reclaimed land (Ohgata village). *Bulletin of the Akita Prefectural College of Agriculture*(15), 137-144.
- Matsunami T, Kokubun M (2003) Photosynthetic responses of four legume crops to fluctuations of evaporative demand following the rainy season. *Tohoku Journal of Agricultural Research* **54**(3/4), 1-11.
- Matsuzaki M (2009) The effect of soil fumigation and organic matter application on yield of five field crops planted continuously more than ten years. *Japanese Journal of Crop Science* **78**(3), 312-323.
- Matsuzaki M, Hamaguchi H, Shimonasako H (1998) The effect of manure application and soil fumigation on the field crops cultivated continuously. *Research Bulletin of the Hokkaido National Agricultural Experiment Station*(166), 1-65.
- Mayilvaganan M, Singh SP (2003) Functional properties of protein concentrate prepared from Phaseolus aconitifolius (Jacq.). *Biochemical and Cellular Archives* **3**(1/2), 73-80.
- Mayumi K, Shibaoka H (1995) A possible double role for brassinolide in the reorientation of cortical microtubules in the epidermal cells of azuki bean epicotyls. *Plant and Cell Physiology* **36**(1), 173-181.
- Mayumi K, Shibaoka H (1996) The cyclic reorientation of cortical microtubules on walls with a crossed polylamellate structure: effects of plant hormones and an inhibitor of protein kinases on the progression of the cycle. *Protoplasma* **195**(1/4), 112-122.
- Mbagwu FN, Edeoga HO (2006) Observations on the vegetative and floral morphology of some Vigna species (Leguminosae-Papilionoideae). *Pakistan Journal of Biological Sciences* **9**(9), 1754-1758.
- McClary DC, Hang AN, Gilliland GC, Babcock JM, Lumpkin TA, Ogg AG, Tanigoshi LK (1993) Herbicides for azuki [Vigna angularis] production. 590-594.
- McIntosh JL, Ismail IG, Effendi S, Sudjadi M (1982) cropping systems to preserve fertility of red yellow podzolic soils in Indonesia. *Tropical Agriculture Research Series*(15), 297-308.
- Meena RS, Rathore GS, Shekhawat BS, Yadav LD, Agnihotri JP (1984) Efficacy of sowing dates and trap crops in management of yellow mosaic of moth (*Vigna aconitifolia* (Jacq.) Marechal). *Indian Journal of Mycology and Plant Pathology* **14**(3), 304-309.
- Meghwal HP, Veer S (2005) A study of the biology of pulse beetle, Callosobruchus chinensis (Linn.) on moth bean, *Vigna aconitifolia* (Jacq.) Marechal. *Indian Journal of Entomology* **67**(4), 334-335.
- Meghwal HP, Veer S, Singh YP (2007) Relative efficacy of some vegetable oils as grain protectant against Callosobruchus chinensis (Linn.) on moth bean. *Indian Journal of Entomology* **69**(1), 22-25.
- Mehra KL, Yadav MS, Singh UP Screening of germplasm collections of cultivated legumes. 35-36.
- Mehrkhoud F, Fathipour Y, Talebi AA, Kamali K, Naseri B (2008b) Population density and spatial distribution patterns of Tetranychus urticae (Acari, Tetranychidae) and its predator Stethorus gilvifrons (Coleoptera: Coccinellidae) on different agricultural crops. *Journal of the Entomological Research Society* **10**(2), 23-36.

- Mehta U, Verma P, Singh I (1993) Effect of cooking, sprouting and dehulling on the nutritional quality and protein digestibility of rice bean (RBL-1) *Vigna umbellata* in comparison to mung bean (*Phaseolus aureus*). *Journal of Dairying Foods and Home Sciences* **12**(3-4), 165-172.
- Meimandi-Nejad MJ (1977) Crop botany. *FAO Plant Production and Protection Paper*(No.9), 27-61.
- Mekasha C, Dzolkifli O, Yusuf S, Rita M, Noorma O (2006) Effect of heat treatment on developmental stages of *Callosobruchus maculatus* (Coleoptera: Bruchidae) in stored adzuki bean *Vigna angularis*. *International Journal of Tropical Insect Science* **26**(4), 273-279.
- Mendoza EMT, Rodriguez FM, Revilleza MJR (1990) Proximate chemical composition of several Philippine indigenous food legumes. *Philippine Agriculturist* **73**(1), 69-74.
- Meng G, Ma C (2002) [Characterization of globulin from Phaseolus angularis \(red bean\)](#). *International Journal of Food Science & Technology* **37**(6), 687-695.
- Meng G, Ma C, Phillips DL (2003) [Raman spectroscopic study of globulin from Phaseolus angularis \(red bean\)](#). *Food Chemistry* **81**(3), 411-420.
- Meng X, Chen W (2001) Applications of AFLP and ISSR techniques in detecting genetic diversity in the soybean brown stem rot pathogen *Phialophora gregata*. *Mycological Research* **105**(8), 936-940.
- Mengistu A, Tachibana H, Grau CR (1991) Selective medium for isolation and enumeration of *Phialophora gregata* from soybean straw and soil. *Plant Disease* **75**(2), 196-199.
- Menon MV, Nair MA (1993) Performance of some fodder legumes in rice fallows. *Journal of Tropical Agriculture* **31**(2), 246-248.
- Menon MV, Nair MA, Potty NN (1998) Variation in elemental composition and nutritional value of fodder legumes in successive cuttings. *Legume Research* **21**(3/4), 236-238.
- Menon MV, Nair MA, Potty NN (1999a) Effect of preceding fodder crops and fallow on productivity mechanics in first crop season rice. *Journal of Tropical Agriculture* **37**(1/2), 85-86.
- Menon MV, Nair MA, Potty NN (1999b) Effect of summer cropping and fallowing on the productivity of rice lands. *Journal of Tropical Agriculture* **37**(1/2), 83-84.
- Merchant A, Richter A, Popp M, Adams M (2006) Targeted metabolite profiling provides a functional link among eucalypt taxonomy, physiology and evolution. *Phytochemistry* **67**(4), 402-+.
- Mertia RS, Kandpal BK, Kumawat RN (2006) Comparative performance of moth bean (*Vigna aconitifolia*) varieties under rainfed condition. *Journal of Arid Legumes* **3**(1), 15-18.
- Messina FJ (1991) Competitive interactions between larvae from divergent strains of the cowpea weevil (Coleoptera: Bruchidae). *Environmental Entomology* **20**(5), 1438-1443.
- Mhase NL, Mote UN, Shelke SS, Kadam DB (1999) Evaluation of varieties/lines of pulse crops against root-knot nematode, *Meloidogyne incognita* race-2. *Indian Journal of Nematology* **29**(1), 91-92.
- Michanie S, Bryan FL, Alvarez P, Olivo AB (1987) Critical control points for foods prepared in households in which babies had salmonellosis. *International Journal of Food Microbiology* **5**(4), 337-354.
- Micheletto S, Rodriguez-Uribe L, Hernandez R, Richins RD, Curry J, O'Connell MA (2007) [Comparative transcript profiling in roots of Phaseolus acutifolius and P. vulgaris under water deficit stress](#). *Plant Science* **173**(5), 510-520.
- Mikoshiba K (1985) Upland crops in Japan (5). Legumes-soybeans and other legumes. *Farming Japan* **19**(2), 49-57.
- Mimura M, Yasuda K, Yamaguchi H (2001) [RAPD variation in wild, weedy and cultivated azuki beans in Asia](#). *Genetic Resources and Crop Evolution* **47**(6), 603-610.
- Minarsih H, Santoso D, Fitrantri N (2001) Identification of P5CS gene on sugarcane by PCR using heterologous primer. *Menara Perkebunan* **69**(1), 1-9.
- Miranda GV, Santos ICd, Peluzio JM, Santos GRd (1997b) Effect of different plant densities on mung beans (*Vigna radiata* (L.) Wilczeck) and rice beans (*Vigna umbellata* (Thunb.) Ohwi & Ohashi). *Revista Ceres* **44**(252), 241-248. [In Portuguese]
- Misal AM, Patil JV, Chavan UD (2008b) Biochemical basis of resistance to stored grain pest (*Callosobruchus chinensis*) in different *Vigna* species. *Journal of Food Legumes* **21**(4), 262-265.
- Mishra A, Muduli KC, Mishra BK, Patra AK (1995) Inter-relationship between yield and its components in rice bean. *Environment and Ecology* **13**(2), 430-432.
- Mishra D (1987) New host records of fungi from India. *Indian Journal of Mycology and Plant Pathology* **17**(2), 236.
- Mishra D (1995) Phenotypic stability for seed yield in rice bean (*Vigna umbellata*). *Indian Journal of Agricultural Sciences* **65**(12), 904-906.
- Mishra M, Behera DB, Mishra PK (1996b) Response of rice bean *Vigna umbellata* (Thunb.) Ohwi and Ohashi to nitrogen levels and weed management. *Environment and Ecology* **14**(2), 361-363.

- Misra BC, Dey A, Sahu PK (2009a) [Genetic divergence in gamma-rays induced mutants of rice bean \(\*Vigna umbellata\*\)](#). *Legume Research* **32**(1), 25-30.
- Misra RC, Sahu PK, Jali CR, Mishra HP, Misra LD (2008a) [Studies on skewness, kurtosis and transgressive variation in M2 populations of rice bean \(\*Vigna umbellata\*\) varieties](#). *Legume Research* **31**(2), 94-99.
- Mitsuhashi-Kato M, Maeda H, Fujii T (1985) Promotion of rooting in Azukia cuttings by possible glycoproteins extracted from Lentinus edodes culture. *Plant and Cell Physiology* **26**(2), 221-228.
- Mitsuhashi-Kato M, Shibaoka H (1981) Effects of actinomycin D and 2,4-dinitrophenol on the development of root primordia in azuki bean stem cuttings. *Plant and Cell Physiology* **22**(8), 1431-1436.
- Mitsunaga T, Fujii K (1997) The effects of spatial and temporal environmental heterogeneities on persistence in a laboratory experimental community. *Researches on Population Ecology* **39**(2), 249-260.
- Mitsunaga T, Fujii K (1999) An experimental analysis of the relationship between species combination and community persistence. *Researches on Population Ecology* **41**(2), 127-134.
- Miura H (1990) [Varietal differences and inheritance of yield components of adzuki bean cultivated in Hokkaido](#). *Japanese Journal of Crop Science* **59**(4), 696-700.
- Miura H (1991) Genetical control of single-seed size and seed size variation in adzuki bean (*Vigna angularis* (Willd) Ohwi & Ohashi). *SABRAO Journal* **23**(2), 83-90.
- Miyahara T, Akimoto I (1984) List and some characteristics of cereal and pulse varieties preserved in Tohoku National Agricultural Experiment Station. *Miscellaneous Publication, Tohoku National Agricultural Experiment Station*(4), 1-92.
- Miyazawa K (1980) Response of crops to the levels of phosphorus accumulated in the surface layers derived from Andosols in Tokachi District. *Research Bulletin of the Hokkaido National Agricultural Experiment Station*(126), 1-30.
- Mizuno K (1994) Inhibition of gibberellin-induced elongation, reorientation of cortical microtubules and change of isoform of tubulin in epicotyl segments of azuki beans by protein kinase inhibitors. *Plant and Cell Physiology* **35**(8), 1149-1157.
- Mizuno N (1979) Studies on chemical characteristics of serpentine soils and mineral deficiencies and toxicities of crops. (29), 85pp.
- Mochizuki T, Takahashi U, Shimamura S, Fukuyama M (2000) [Secondary aerenchyma formation in hypocotyl in summer leguminous crops](#). *Japanese Journal of Crop Science* **69**(1), 69-73.
- Mohamed SV, Sung J, Jeng T, Wang C (2006) Organogenesis of *Phaseolus angularis* L.: high efficiency of adventitious shoot regeneration from etiolated seedlings in the presence of N6-benzylaminopurine and thidiazuron. *Plant Cell, Tissue and Organ Culture* **86**(2), 187-199.
- Mohan HM, Chittapur BM, Hiremath SM, Chimmad VP (2005) Performance of maize under intercropping with grain legumes. *Karnataka Journal of Agricultural Sciences* **18**(2), 290-293.
- Mohanty SK, Baisakh B (1999) Phenotypic stability for yield and maturity in rice bean *Vigna umbellata* at rainfed condition. *Environment and Ecology* **17**(1), 8-10.
- Mohapatra AK (1998) Effect of time of sowing, spacing and fertility levels on growth and yield of rice bean (*Vigna umbellata*). *Indian Journal of Agronomy* **43**(1), 118-121.
- Mohapatra AK, Halder J (1998) Intercropping of soybean and rice bean with rice and ragi under different planting patterns. *Annals of Agricultural Research* **19**(1), 114-115.
- Mohapatra AK, Paikaray RK, Misra RC, Halder J (1996b) Response of rice bean (*Vigna umbellata*) to nitrogen and phosphorus fertilization in acid red soil. *Annals of Agricultural Research* **17**(4), 445-446.
- Mohapatra BK, Pradhan L (1992) Intercropping fodder legumes with maize in different planting patterns. *Annals of Agricultural Research* **13**(4), 366-371.
- Mohapatra BK, Pradhan L (1993a) Energy relationship in intercropping of maize (*Zea mays*) with cowpea (*Vigna unguiculata*) and rice bean (*Vigna umbellata*). *Indian Journal of Agricultural Sciences* **63**(9), 581-583.
- Mohapatra S, Tripathy SK, Ghosh DC (2005) Effect of sabaigrass (*Eulaliopsis binata*) intercropping with grain legumes on soil fertility, yield and economics. *Indian Journal of Agricultural Sciences* **75**(8), 498-500.
- Mohapatra S, Tripathy SK, Patra AK (2001b) Effect of forage legumes intercropping on growth, yield and economics of sabaigrass (*Eulaliopsis binata*). *Indian Journal of Agricultural Sciences* **71**(1), 14-16.
- Mollah AS, Begum A (2001) A study on transfer factors of Co-60 and Zn-65 from soil to plants in the tropical environment of Bangladesh. *Environmental Monitoring and Assessment* **68**(1), 91-97.
- Mondal SS, Chettri M (1998) Integrated nutrient management for sustaining productivity and fertility under rice (*Oryza sativa*)-based cropping system. *Indian Journal of Agricultural Sciences* **68**(7), 337-340.

- Montenegro B J, Galindo A J (1974) Bean common mosaic virus in Guanajuato state and an evaluation of varietal resistance. *El virus del mosaico comun del frijol en el estado de Guanajuato y evaluacion de la resistencia de variedades.*(No. 18), 89-95.
- Moody K, Robles RP, Floresca ET (1986a) Weed control in legumes in the Philippines. In. ' pp. 212-267)
- Moog FA, Sena EM, Valenzuela FG (1977) Crops and livestock in two Batangas barrios. *Philippine Journal of Animal Industry* **32**(1/4), 11-24.
- Moon J, Lee Y, et al. (2006) A new light green seed coated Azuki bean cultivar, "Yeonkeum" with middle seed size and good seed quality. *Korean Journal of Breeding* **38**(4), 303-304.
- Mora A (1996) Method for adjusting a yield-density model for data derived from the Nelder fan design. *Revista de la Facultad de Agronomía, Universidad Central de Venezuela* **22**(1/2), 31-36. [In Spanish]
- Mora Garcés A (1999) Statistical analysis to data from Nelder Fan Design using an Antedependence model. *Revista de la Facultad de Agronomía, Universidad Central de Venezuela* **25**(1), 75-82. [In Spanish]
- Moriyama M, Oba K (2008) Comparative study on the vitamin C contents of the food legume seeds. *Journal of Nutritional Science and Vitaminology* **54**(1), 1-6.
- Mosha TCE, Bennink MR (2004) [Protein quality of drum-processed cereal-bean-sardine composite supplementary foods for preschool-age children](#). *Journal of the Science of Food and Agriculture* **84**(10), 1111-1118.
- Mosha TCE, Bennink MR (2005) [Protein digestibility-corrected amino acid scores, acceptability and storage stability of ready-to-eat supplementary foods for pre-school age children in Tanzania](#). *Journal of the Science of Food and Agriculture* **85**(9), 1513-1522.
- Mosha TCE, Bennink MR, Ng PKW (2005) Nutritional quality of drum-processed and extruded composite supplementary foods. *Journal of Food Science* **70**(2), C138-C144.
- Moue T (1979) Interspecific hybridization between an A-genome species (Chinese cabbage) and a C-genome species (cabbage) of Brassica. 3. Embryo and ovule culture of interspecific hybrids of Chinese cabbage and cabbage. *Bulletin of the Faculty of Agriculture, Niigata University (Niigata Daigaku Nogakubu Kenkyu Hokoku)*(31), 47-53.
- Muhammad B, Jamali AR, Zahoor A (2006) Genetic resistance in mungbean and mashbean germplasm against mungbean yellow mosaic begomovirus. *Mycopath* **4**(2), 1-4.
- Muhammad Y (2007) Screening of indigenous mothbean lines against yellow mosaic virus. *Sarhad Journal of Agriculture* **23**(2), 445-449.
- Muhammad Y, Najeebulah, Pervez K (2005) Moth bean germplasm evaluation for yield and other important traits. *Indus Journal of Plant Sciences* **4**(2), 241-248.
- Muhammad Y, Najibullah, Pervez K (2007) Genetic divergence in some mothbean [*Vigna aconitifolia* (Jacq) Marechal] genotypes. *Sarhad Journal of Agriculture* **23**(2), 417-421.
- Mukai Y, Sato S (2009) Polyphenol-containing azuki bean (*Vigna angularis*) extract attenuates blood pressure elevation and modulates nitric oxide synthase and caveolin-1 expressions in rats with hypertension. *Nutrition, Metabolism and Cardiovascular Diseases* **19**(7), 491-497.
- Mukerji N, Verma BL (1994) Effect of time lag of first post-sowing irrigation on yield and technological qualities of solid and legume inter-cropped cotton. *Madras Agricultural Journal* **81**(6), 338-340.
- Mukherjee AK (1988b) Role of total available carbohydrate reserve and leaf area content in the stubbles on the regrowth of rice bean *Vigna umbellata*. *Environment and Ecology* **6**(2), 457-459.
- Mukherjee AK, Chatterjee BN (1979a) Herbage growth analysis of rice bean in tropical India. I. Uninterrupted growth condition. *Forage Research* **5**(2), 117-123.
- Mukherjee AK, Chatterjee BN (1979b) Herbage growth analysis of rice bean in tropical India. II. Under interrupted growth condition. *Forage Research* **5**(2), 125-133.
- Mukherjee AK, Maiti S, Nanda MK, Madhumita N (2006) Productivity of rice bean as affected by chemical fertilizer and poultry manure. *Journal of Crop and Weed* **2**(2), 34-36.
- Mukherjee AK, Maiti S, Nanda MK, Simul B, Madhumita N (2005b) Effect of Phosphorus sources on seed production of rice bean (*Vigna umbellata*) cultivars in red lateritic zones of West Bengal. *Journal of Crop and Weed* **1**(1), 5-7.
- Mukherjee AK, Mandal SR, Patra BC, Udita M (2004) Effect of different varieties, dates of sowing and levels of phosphate on seed production of ricebean. *Environment and Ecology* **22**(2), 319-321.
- Mukherjee AK, Mishra GC, Roy G, Nanda MK (1998) Effect of different phosphorus levels on growth, nodulation and dry matter production in ricebean cultivars. *Forage Research* **24**(2), 115-118.
- Mukherjee AK, Roquib MA (1980) Herbage and grain yield potentiality of rice bean (*Vigna umbellata* (Thunb.) Ohwi and Ohashi) -- a review. *Forage Research* **6**(2), 165-170.

- Mukherjee AK, Roy G, Mishra GC, Nanda MK (1998) Effect of cultivars and phosphorus levels on the seed production of ricebean under rainfed ecosystem. *Forage Research* **24**(3), 183-184.
- Mukherjee AK, Sarkar A (1991) Rice bean - a new crop for sustaining pulse production in India. *Environment and Ecology* **9**(1), 207-211.
- Mukhopadhyay B, Roychoudhury N (1987) Biology of green stink bug Nezara viridula.. *Environment and Ecology* **5**(2), 325-327.
- Mulk MM, Jairajpuri MS (1975) Nematodes of leguminous crops in India. III. Three new species of Hoplolaimus Daday, 1905 (Hoplolaimidae). *Indian Journal of Nematology* **5**(1), 1-8.
- Murata K, Narikawa T, Chiba I, Satoh H, Adachi T, Matsukawa I (1985) Breeding the new adzuki bean Erimo. *Bulletin, Hokkaido Prefectural Agricultural Experiment Stations*(53), 103-113.
- Musa MM (1978a) Legume studies. Evaluation of nitrogen fixation in pot experiment. 176-177.
- Musa MM (1978b) Legume studies. Forage crop sequence experiment. 175-176, 177.
- Musa MM, Burhan HO (1974) [The relative performance of forage legumes as rotational crops in the Gezira. Experimental Agriculture](#) **10**(2), 131-140.
- Muszynski S, Hiraiwa S, Tanaka S (1980) Monitoring radiation damage in the seedlings of adzuki bean (*Vigna angularis* (Willd.) Ohwi & Ohashi) after seed irradiation with gamma rays. *Japanese Journal of Breeding* **30**(3), 225-230.
- Muthusamy S, Kanagarajan S, Ponnusamy S (2008) Efficiency of RAPD and ISSR markers system in accessing genetic variation of rice bean (*Vigna umbellata*) landraces. *Electronic Journal of Biotechnology* **11**(3).
- Muzaffar MT, Khan HH (1985) Phytic acid in whole seed and dehulled legumes. *Journal of Agricultural Research, Pakistan* **23**(3), 223-229.
- Mysliwa-Kurdziel B, Prasad MNV, Strzalka K (2004) Photosynthesis in heavy metal stressed plants. *Heavy Metal Stress in Plants: from Biomolecules to Ecosystems, Second Edition*, 146-181.
- Nabhan G (1979) Developing new crops is old idea in Arizona. *Progressive Agriculture in Arizona* **30**(3), 12-15.
- Nagaoka Y, Sawada S (2003) [Representation of seed coat color of azuki bean by using XYZ color system. Japanese Journal of Crop Science](#) **72**(4), 471-474.
- Nagaoka Y, Sawada S, Kato K (2004) [Effects of temperature during ripening period on seed coat color of azuki bean. Japanese Journal of Crop Science](#) **73**(3), 330-335.
- Nagaraj NC, Muniyappa V, Satyan BA (1981) Resistance source for mung bean yellow mosaic virus. 69-72.
- Nagasawa S, Yamada K (1988) An asymptotic regression curve for describing the emergence pattern of the azuki bean weevil, *Callosobruchus chinensis* L. (Coleoptera: Bruchidae). *Applied Entomology and Zoology* **23**(2), 181-185.
- Nagasuna T, Tagawa A, Ogawa Y, Iimoto M (2005) Water absorption rate and volume change during soaking for adzuki beans and soybeans. *Nippon Shokuhin Kagaku Kogaku Kaishi = Journal of the Japanese Society for Food Science and Technology* **52**(12), 566-571.
- Nagpal AK, Gorakh M, Raja P, Kiradoo BD (1998) Utilization of bui (Aerva tomentosa) leaves in the ration of camel. *Indian Journal of Animal Nutrition* **15**(3), 212-215.
- Nagpal AK, Kiradoo BD, Raja P, Gorakh M, Ram K (1998) Comparative studies on stall-feeding and continuous pasture grazing systems on camel production. *Indian Journal of Animal Nutrition* **15**(3), 151-157.
- Nagpal AK, Manju A, Singh GP (2003a) Utilization of mixed and complete rations by lactating camels. *Indian Journal of Animal Nutrition* **20**(2), 178-184.
- Nagpal AK, Manju A, Singh GP (2003b) Utilization of moth chara, wheat straw and bui leaves based complete rations in camel calves. *Indian Journal of Animal Sciences* **73**(11), 1267-1270.
- Nagpal AK, Rai AK, Khanna ND (1993) Nutrient utilization in growing camel calves kept at two watering schedules. *Indian Journal of Animal Sciences* **63**(6), 671-673.
- Nagpal AK, Roy AK, Kiradoo BD, Raja P, Sahani MS (2004) Voluntary feed intake and nutrient utilization of adult female racing camels during exercise and at rest. In '' pp. 795-800. (The Camelid Publishers: Bikaner)
- Nagpal AK, Sahani MS, Roy AK (1998a) Voluntary feed intake, digestibility and utilization of macro-and micro-elements in dry and pregnant camels. *Indian Journal of Animal Nutrition* **15**(3), 158-162.
- Nagpal AK, Sahani MS, Roy AK (2000) Effect of grazing sewan (Lasiurus sindicus) pasture on female camels in arid ecosystem. *Indian Journal of Animal Sciences* **70**(9), 968-971.
- Nagpal AK, Sahani MS, Roy AK, Gorakh M (1998b) Voluntary feed intake and utilization of macro- and micro-nutrients in dry and lactating Bikaneri camels. *International Journal of Animal Sciences* **13**(1), 19-24.
- Nahas E, Machado JO (1982) Studies on the soybean meal grown rhizopus-oligosporus for tempeh fermentation. *Cientifica (Jaboticabal)* **10**(2), 217-224.

- Naidu MR, Bakshi R (1987) A preliminary study on genetic variability for some quantitative traits in mothbean. *Agricultural Science Digest, India* **7**(3), 131-133.
- Naidu MR, Sultan S, Bakshi R (1986) Variability and path-coefficient analysis of yield components in mothbean (*Vigna aconitifolia* Jacq.). *Haryana Agricultural University Journal of Research* **16**(2), 168-171.
- Naik RM, Jadhav SS (2001b) Biochemical basis of resistance to stored grain pests in rice bean (*Vigna umbellata*). *Indian Journal of Agricultural Biochemistry* **14**(1/2), 57-58.
- Naik VN, Pokle DS (1985) Novelties in the flora of Marathwada. *Journal of Economic and Taxonomic Botany* **7**(3), 670-675.
- Naiki T, Ui T (1981) Rhizoctonia solani Kuhn root rot of bean, soybean and adzuki bean seedlings. *Memoirs of the Faculty of Agriculture Hokkaido University* **12**(4), 262-269.
- Nair AK (2000) Effect of farmyard manure and fertilizer levels on yield of maize (*Zea mays*) and succeeding rice bean (*Vigna umbellata*) crop in Sikkim. *Indian Journal of Agricultural Sciences* **70**(4), 239-240.
- Naito S, Kanematsu S (1994) Characterization and pathogenicity of a new anastomosis subgroup AG-2-3 of Rhizoctonia solani Kühn isolated from leaves of soybean. *Annals of the Phytopathological Society of Japan* **60**(6), 681-690.
- Nakajima M, Takita K, Wada H, Miura K, Hasegawa M, Yamaguchi I, Murofushi N (1997) Partial purification and characterization of a gibberellin-binding protein from seedlings of Azukia angularis [*Vigna angularis*]. *Biochemical and Biophysical Research Communications* **241**(3), 782-786.
- Nakamura T, Yokoyama R, Tomita E, Nishitani K (2003) Two azuki bean XTH genes, VaXTH1 and VaXTH2, with similar tissue-specific expression profiles, are differently regulated by auxin. *Plant and Cell Physiology* **44**(1), 16-24.
- Nakamura Y, Shibaoka H (1980) Increase in the gibberellin response by low temperature pretreatment of azuki bean epicotyls. *Botanical Magazine* **93**(1029), 77-87.
- Nakamura Y, Wakabayashi K, Kamisaka S, Hoson T (2002) [Effects of temperature on the cell wall and osmotic properties in dark-grown rice and azuki bean seedlings](#). *Journal of Plant Research* **115**(6), 455-461.
- Nakano S, Soga K, Wakabayashi K, Hoson T (2007) Different cell wall polysaccharides are responsible for gravity resistance in the upper and the basal regions of azuki bean epicotyls. *Biological Sciences in Space* **21**(4), 113-116.
- Nakaseko K (1984) Studies on dry matter production, plant type and productivity in grain legumes. *Memoirs of the Faculty of Agriculture, Hokkaido University* **14**(2), 103-158.
- Nakaseko K, Gotoh K (1981) [Comparative studies of dry matter production, plant type and productivity in soyabean, azuki bean and kidney bean. IV. Dry matter production of azuki bean at various population densities](#). *Japanese Journal of Crop Science* **50**(3), 388-395.
- Nakaseko K, Gotoh K (1982) [Comparative studies on dry matter production, plant type and productivity in soybean, azuki bean and kidney bean. VI. Differences in the early vegetative growth in relation to seed weight and character of cotyledons](#). *Japanese Journal of Crop Science* **51**(1), 110-116.
- Nakaseko K, Gotoh K (1983) [Comparative studies on dry matter production, plant type and productivity in soyabean, azuki bean and kidney bean. VII. An analysis of the productivity among the three crops on the basis of radiation absorption and its efficiency for dry matter accumulation](#). *Japanese Journal of Crop Science* **52**(1), 49-58.
- Nakaseko K, Gotoh K, Asanuma K (1979a) [Comparative studies on dry matter production, plant type and productivity in soyabean, azuki bean and kidney bean. 1. Differences in dry matter accumulation patterns under low population density](#). *Japanese Journal of Crop Science* **48**(1), 82-91.
- Nakaseko K, Gotoh K, Asanuma K (1979b) [Comparative studies on dry matter production, plant type and productivity in soyabean, azuki bean and kidney bean. 2. Relationships between vertical distribution of leaf area and some morphological characteristics](#). *Japanese Journal of Crop Science* **48**(1), 92-98.
- Nakaseko K, Yoshida K (1989) The effect of epi-brassinolide applied at the flowering stage on growth and yield of soyabean and azuki bean. *Memoirs of the Faculty of Agriculture, Hokkaido University* **16**(4), 347-352.
- Nakata M (1991) Participation of acetolactate and acetoin producing system for chlorsulfuron-sensitivity in higher plants. *Weed Research (Tokyo)* **36**(1), 58-67.
- Nanda MK, Mukherjee AK, Tripathi B (1997) Response of fertilizers on different crop sequences. *Journal of Interacademicia* **1**(4), 301-306.
- Nanda MK, Patra B, Mukherjee AK (2000) Yield performance of ricebean as pulse in West Bengal. *Environment and Ecology* **18**(1), 17-19.
- Nanda SS, Alim MA, Dash B, Santra GH (2000) Influence of sowing time and spacing on fodder yield and quality of ricebean (*Vigna umbellata* Thunb.). *Environment and Ecology* **18**(3), 584-586.

- Narain U, Tewari TN, Dwivedi RP (1995) Myrothecium leaf spot of ricebean in India. *Indian Journal of Pulses Research* **8**(1), 106-107.
- Narwal SS, Palaniraj R, Sati SC, Kadian HS, Dahiya DS (2003) Allelopathic plants: 8. Parthenium hysterophorus L. *Allelopathy Journal* **11**(2), 151-170.
- Naseri B, Fathipour Y, Talebi A (2008b) Population fluctuation and thermal requirement for development of Empoasca decipiens (Homoptera, Cicadellidae) on different bean species in natural conditions. *Journal of the Entomological Research Society* **10**(3), 11-23.
- Naseri B, Fathipour Y, Talebi AA (2007b) Comparison of some biological aspects of Empoasca decipiens (Hom.: Cicadellidae) on four bean species. *Journal of Entomological Society of Iran* **27**(1), Pe1-13. [In Persian]
- Naseri B, Fathipour Y, Talebi AA (2007c) Seasonal parasitism of Empoasca decipiens by Anagrus atomus on four bean species in Tehran area. *Applied Entomology and Phytopathology* **75**(1), 4 (En), 1-11 (Pe). [In Persian]
- Nasser LCB, Vieira RF (1997b) Performance of Phaseolus and Vigna species in the cerrado of Planaltina, Distrito Federal. *Cientifica (Jaboticabal)* **25**(1), 223-227. [In Portuguese]
- Nath D, Chowdhury BR, Dasgupta T (2005b) Genetic divergence in some cultivated and wild species of Vigna. *Indian Agriculturist* **49**(1/2), 27-32.
- Navjot S, Satija CK (2003b) Cytomorphological characterization of amphidiploids of *Vigna radiata* × *V. umbellata*. *Crop Improvement* **30**(1), 25-32.
- Nayak SP, Prasad G, Sahoo HK, Mishra SN (1997b) Effects of spacing and varieties on yield and yield attributing characters of ricebean (*Vigna umbellata*). *Environment and Ecology* **15**(3), 730-731.
- Neelam Y, Mahala SC, Yadav VK (2003) Modulation of plant water relations by gibberellic acid in mothbean under salinity. *Indian Journal of Plant Physiology* **8**(3), 297-301.
- Neelam Y, Mahla SC, Yadav VK (2004) Effect of growth regulators on peroxidase and membrane stability under salinity in mothbean seedlings. *Advances in Plant Sciences* **17**(1), 207-211.
- Neelam Y, Mahla SC, Yadav VK (2006) Reversal of salinity stress effects on morpho-physiological parameters of mothbean seedlings by growth regulators. *Indian Journal of Plant Physiology* **11**(3), 310-313.
- Neelam Y, Rathore GS, Majumdar VL, Yadav VK (1994) Stomatal response to Macrophomina phaseolina (Tassi) Goid infection in moth bean (*Vigna aconitifolia*). *Annals of Arid Zone* **33**(1), 91-92.
- Negi A, Boora P (2003) Investigation on effect of domestic processing and cooking methods on the mineral composition of some high yielding varieties of moth bean (*Vigna aconitifolia* Jacq. Marechal). *Forage Research* **29**(2), 97-101.
- Negi A, Boora P, Khetarpaul N (2001) Starch and protein digestibility of newly released moth bean cultivars: effect of soaking, dehulling, germination and pressure cooking. *Nahrung* **45**(4), 251-254.
- Negi KS, Muneem KC (1997) Adzukibean (*Vigna angularis* (Willd.) Ohwi & Ohashi) - a new crop for Uttarakhand Himalaya. *Indian Journal of Forestry* **20**(2), 144-146.
- Negi KS, Pant KC, Muneem KC, Bhag M (1996) Evaluation of rice bean genetic resources. *Indian Journal of Forestry* **19**(2), 156-163.
- Negishi H, Kobayashi K (1984) Effect of soybean cyst nematode on the incidence of brown stem rot (BSR) of adzuki bean. *Annals of the Phytopathological Society of Japan* **50**(4), 500-506.
- Nelson EA, Penner D (2005) Sensitivity of selected crops to isoxaflutole in soil and irrigation water. *Weed Technology* **19**(3), 659-663.
- Neumayr L, Krämer J (1990) Production of enterotoxin A and thermonuclease by *Staphylococcus aureus* in legumes. *International Journal of Food Microbiology* **10**(3-4), 225-234.
- Ng NQ (1992b) Wide crosses of Vigna food legumes. *Biotechnology: Enhancing research on tropical crops in Africa*, 75-80. (Technical Centre for Agricultural and Rural Cooperation: Wageningen)
- Niba LL, Rose N (2003) Effect of soaking solution concentration on resistant starch and oligosaccharide content of adzuki (*V. angularis*), fava (*V. faba*), lima (*P. lunatus*) and mung bean (*V. radiata* L.). *Journal of Food Technology* **1**(1), 4-8.
- Nikitina KV, Budanova VI, Stepanova SI, Yaskina OS (1980) Rapid method of evaluating resistance to bacterial diseases in lupin and Phaseolus. *Selektsiya i Semenovodstvo, USSR*(5), 22-23.
- Nimkar PM, Mandwe DS, Dudhe RM (2005) Physical properties of moth gram. *Biosystems Engineering* **91**(2), 183-189.
- Nishi M, Joginder S, Sachendra B, Avinash B, Anil V (2006) Effect of soil compaction, potassium and cobalt on growth and yield of moth bean. *International Journal of Soil Science* **1**(3), 269-271.
- Nishi M, Joginder S, Sachendra B, Avinash B, Anil V (2007) Effect of phosphorus application and time of harvest on yield and quality of moth bean [*Vigna aconitifolia* (Jacq.) Marechal]. *Journal of Plant Sciences* **2**(2), 247-250.

- Nishi M, Joginder S, Sachendra B, Avinash B, Rahul S, Anil V (2008a) Influence of spacing on seed yield and water relations of moth bean genotypes under arid conditions. *Journal of Food Legumes* **21**(1), 53-55.
- Nishi M, Joginder S, Sachendra B, Avinash B, Rahul S, Anil V (2008b) Response of moth bean genotypes to nutrient management under arid conditions of Indian Thar desert. *Journal of Food Legumes* **21**(1), 69-70.
- Nishiiri K, Matsui S, Watanabe Y (1981) Studies on a crop sequence in Hokkaido with special emphasis on behaviour of agronomic characteristics. I. Growth and yield of soyabean following different crops. *Research Bulletin of the Hokkaido National Agricultural Experiment Station*(130), 113-122.
- Nishimune A (1984) Evaluation of the soil nitrogen supply in upland field crops in Tokachi, Hokkaido. *Research Bulletin of the Hokkaido National Agricultural Experiment Station*(140), 33-91.
- Nishimune A, Konno T, Saito G, Fujita I (1983) The relationship between nitrogen fixation and grain yields of legumes on the main upland soils in Tokachi, Hokkaido. *Research Bulletin of the Hokkaido National Agricultural Experiment Station*(137), 81-106.
- Nishimura K (1993) Oviposition strategy of the parasitic wasp Dinarmus basalis (Hymenoptera, Pteromalidae). *Evolutionary Ecology* **7**(2), 199-206.
- Nishimura K, Jahn GC (1996) Sex allocation of three solitary ectoparasitic wasp species on bean weevil larvae: sex ratio change with host quality and local mate competition. *Journal of Ethology* **14**(1), 27-33.
- Nishitani K (1992) A novel method for detection of endo-xyloglucan transferase. *Plant and Cell Physiology* **33**(8), 1159-1164.
- Nishitani K (1993) Endo-xyloglucan transferase and cell wall extension. *Chemical Regulation of Plants* **28**(2), 135-144.
- Nishitani K, Masuda Y (1980) Modifications of cell wall polysaccharides during auxin-induced growth in azuki bean epicotyl segments. *Plant and Cell Physiology* **21**(1), 169-181.
- Nishitani K, Masuda Y (1981) Auxin-induced changes in the cell wall structure: Changes in the sugar compositions, intrinsic viscosity and molecular weight distributions of matrixpolysaccharides of the epicotyl cell wall of *Vigna angularis*. *Physiologia Plantarum* **52**(4), 482-494.
- Nishitani K, Masuda Y (1982) Roles of auxin and gibberellic acid in growth and maturation of epicotyls of *Vigna angularis*: cell wall changes. *Physiologia Plantarum* **56**(1), 38-45.
- Nishitani K, Masuda Y (1983) Auxin-induced changes in the cell wall xyloglucans: effects of auxin on the two different subfractions of xyloglucans in the epicotyl cell wall of *Vigna angularis*. *Plant and Cell Physiology* **24**(3), 345-355.
- Nishitani K, Nevins DJ (1991) An enzyme probe for the resolution of glucuronoxylan and glucuronoarabinoxylan structures. *Food Hydrocolloids* **5**(1-2), 197-207.
- Nishitani K, Tominaga R (1991) In vitro molecular weight increase in xyloglucans by an apoplastic enzyme preparation from epicotyls of *Vigna angularis*. *Physiologia Plantarum* **82**(4), 490-497.
- Nishiyama Y, Matsuo E, Ishiguro E, Inanaga S, Miyazoto M (1989) Studies on the irradiation effects on some useful plants - sunflower, okra and beans. *Bulletin of the Faculty of Agriculture, Kagoshima University*(39), 233-242.
- Nishizawa K, Teraishi M, Utsumi S, Ishimoto M (2007) [Assessment of the importance of  \$\alpha\$ -amylase inhibitor-2 in bruchid resistance of wild common bean](#). *TAG Theoretical and Applied Genetics* **114**(4), 755-764.
- Nitta T (1990) The stimulation of root activity and a reduction in root diseases of upland crops through improvements in the rhizosphere microflora. *Research Journal of Food and Agriculture* **13**(6), 15-20.
- Nitta T (1991) Diversity of root fungal floras: its implications for soil-borne diseases and crop growth. *JARQ, Japan Agricultural Research Quarterly* **25**(1), 6-11.
- Nozawa H, Yamagata H, Aizono Y, Iwasaki T (1993) Temperature-dependence of the reconstitution of an adzuki bean [*Vigna angularis*] subtilisin inhibitor from its inactive two fragments. *Bioscience, Biotechnology and Biochemistry* **57**(1), 154-155.
- Numata A, Sakuyama K, Kowata H, Kamiyama Y (1991) The relations between sowing date and seed coat colour in azuki bean 'Benidainagon'. *Report of the Tohoku Branch, the Crop Science Society of Japan*(34), 11-12.
- Nupur M, Nirmal M (2003) Random amplified polymorphic DNA (RAPD) assay in mothbean [*Vigna aconitifolia* (Jacq.) Marechal]. *Journal of Interacademicia* **7**(4), 480-483.
- O'Garro LW, Paulraj LP (1997) Onion leaf blight caused by *Xanthomonas campestris*: alternative hosts and resistant onion genotypes. *Plant Disease* **81**(9), 978-982.
- Obokata J, Mikami K, Hayashida N, Nakamura M, Sugiura M (1993) Molecular heterogeneity of photosystem I. psaD, psaE, psaF, psaH, and psaL are all present in isoforms in *Nicotiana* spp. *Plant Physiology* **102**(4), 1259-1267.

- Oda A, Shimizu M, Kuroha T, Satoh S (2005) Induction of xylem sap methylglycine by a drought and rewatering treatment and its inhibitory effects on the growth and development of plant organs. *Physiologia Plantarum* **124**(4), 515-523.
- Ofuya TI, Credland PF (1996) The ability of *Bruchidius atrolineatus* (Pic) (Coleoptera: Bruchidae) to infest and damage seeds of different tropical legumes. *Journal of Stored Products Research* **32**(4), 323-328.
- Ogasawara Y, Hidano Y, Kato Y (2006) Study on carbohydrate composition of apios (*Apis americana* Medikus) flowers and tubers. *Nippon Shokuhin Kagaku Kogaku Kaishi = Journal of the Japanese Society for Food Science and Technology* **53**(2), 130-136.
- Ogawa K, Takeuchi Y, Katayama M (1988) Biomass production and the amounts of absorbed inorganic elements by crops in arable lands in Hokkaido, and its evaluation. *Research Bulletin of the Hokkaido National Agricultural Experiment Station*(149), 57-91.
- Oh C, Kim J, Kim K, Mabry TJ (1995) Rapid gas chromatographic screening of edible seeds, nuts and beans for non-protein and protein amino acids. *Journal of Chromatography* **708**(1), 131-141.
- Oh H, Chun J, Lee M, Min K, Lee S, Cheon C (2000) Role of hsfA gene on host-specificity by *Bradyrhizobium japonicum* in a broad range of tropical legumes. *Canadian Journal of Microbiology* **46**(1), 81-84.
- Ohashi H, Takahashi H (1981) Pollen morphology of *Vigna angularis* (Leguminosae). *Botanical Magazine, Tokyo* **94**(1034), 177-180.
- Ohashi T, Ishimizu T, Akita K, Hase S (2007) In vitro stabilization and minimum active component of polygalacturonic acid synthase involved in pectin biosynthesis. *Bioscience, Biotechnology and Biochemistry* **71**(9), 2291-2299.
- Ohba K, Minoura M, Safarpour MM, Picard GL, Safarpour H (1997) Method for the determination of imazamox and its two hydroxy and glucose conjugate metabolites in adzuki beans by capillary electrophoresis. *Journal of Pesticide Science* **22**(4), 277-281.
- Ohba K, Nirei M, Watanabe S, Han K, Hashimoto N, Shimada K, Sekikawa M, Chiji H, Fukushima M (2005) Effect of an adzuki bean extract on hepatic anti-oxidant enzyme mRNAs in D-galactosamine-treated rats. *Bioscience, Biotechnology and Biochemistry* **69**(10), 1988-1991.
- Ohno T, Miyatake T (2007) Drop or fly? Negative genetic correlation between death-feigning intensity and flying ability as alternative anti-predator strategies. *Proceedings of the Royal Society of London. Series B, Biological Sciences* **274**(1609), 555-560.
- Ohtaki Y, Kiyoohara T, Iwasaki T, Yoshikawa M (1980) Temporary site in the anti-tryptic fragment from adzuki-bean proteinase inhibitor II. *Biochemical and Biophysical Research Communications* **96**(2), 719-724.
- Ohtsuka T, Hikawa M, Saka H (1990) Growth regulating action of isoprothiolane in plants. V. Interactions between RNA, protein synthesis inhibitors and isoprothiolane on adventitious root formation in stem segments of pulse crops. *Japanese Journal of Crop Science* **59**(3), 566-571.
- Oka M, Nagamine T, Egawa Y, Katsuta M, Nakagahra M (1989) Collection of crop genetic resources in the central parts of Japan in 1987. *JICA Ref Series*(2), 119-135.
- Okada T, Kodama F, Fujita S, Kondo N, Naito S (2002) Effect of thiram on seed decay and seedling blight of adzuki bean caused by *Pythium* spp. *53rd Annual Report of the Society of Plant Protection of North Japan*, 99-100.
- Okada T, Kodama F, Fujita S, Murata K, Kondo N, Kobayashi K (2001) Pythium species isolated from ungerminated adzuki bean seeds. Society of Plant Protection of North Japan, Akita.
- Okamura M, Hosoya S (2008) Development of Japanese space foods "sweet red bean paste". *Nippon Shokuhin Kagaku Kogaku Kaishi = Journal of the Japanese Society for Food Science and Technology* **55**(9), 402-404.
- Oli S, Sima B, Sunirmal C (1989b) Cytomorphological studies in some taxa of *Phaseolus* Linn. and *Vigna* Savi. *Cytologia* **54**(1), 97-108.
- Oliveira Gd, Guimarães VM, Lima e Borges EEd, Fialho LdS, Almeida e Oliveira MGd, Rezende STd (2005b) Purification and characterization of  $\alpha$ -galactosidases from *Platymiscium pubescens* Micheli seeds. *Revista Árvore* **29**(4), 535-543. [In Portuguese]
- Oliveira JTA, Pinto VPT, Vasconcelos IM, Fernandes CF, Ramos MV, Ferreira FVA, Rios FJB (2004) In vitro and in vivo digestibility of the albumin and globulin fractions of eight Brazilian cowpea *Vigna unguiculata* (L.) Walp cultivars. *Journal of the Science of Food and Agriculture* **84**(14), 1823-1830.
- Om P, Sharma RS, Singh BP (2002) Effects of phosphorus and row spacing on the yield attributes and yield of moth bean cultivars under rainfed conditions. *Indian Journal of Pulses Research* **15**(2), 142-144.
- Ong HT (1980) Effects of mannitol induced water stress on the ribosomes of intact leaves of azuki bean seedlings. *Biologia Plantarum* **22**(4), 249-254.

- Oogita T, Misumi T, Naitou M, Gotou M, Ogawa N, Tanigawa N (2009) Method of estimating inorganic bromine residue from fumigation with methyl bromide. *Research Bulletin of the Plant Protection Service, Japan*(45), 59-61.
- Ooume K, Inoue Y, Soga K, Wakabayashi K, Fujii S, Yamamoto R, Hoson T (2009) [Cellular basis of growth suppression by submergence in azuki bean epicotyls](#). *Annals of Botany* **103**(2), 325-332.
- Ortiz-R C, Francisco-G E (1987) Stochastic efficiency and the selection of crop varieties. *Ciencia e Investigacion Agraria* **14**(3), 175-188.
- Osaki M (1993) Carbon-nitrogen interaction model in field crop production. 223-226.
- Osaki M (1995) Ontogenetic changes of N, P, and K contents in individual leaves of field crops. *Soil Science and Plant Nutrition* **41**(3), 429-438.
- Osaki M, Shinano T (2001) Plant growth based on interrelation between carbon and nitrogen translocation from leaves. *Photosynthetica* **39**(2), 197-203.
- Osaki M, Shinano T, Tadano T (1992) Carbon-nitrogen interaction in field crop production. *Soil Science and Plant Nutrition* **38**(3), 553-564.
- Osaki M, Yamada S, Tadano T (1995) Effect of sink manipulation on nitrogen accumulation and distribution among organs of Gramineae and Leguminosae. *Soil Science and Plant Nutrition* **41**(1), 33-44.
- Osman AE, Diek AAA (1982) [Effects of defoliation on yield and forage quality of some tropical grasses, legumes and their mixtures](#). *Experimental Agriculture* **18**(2), 157-166.
- Osten VA (1996) Weed management studies in ley legumes for Central Queensland farming systems. In '!' pp. 108-110. (Weed Science Society of Victoria Inc.: Victoria)
- Otanit T, Seike N, Miwa T (2006) Levels of dioxins in rice, wheat, soybean, and adzuki bean cultivated in 1999 to 2002 in Japan and estimation of their intake. *Shokuhin Eiseigaku Zasshi = Journal of the Food Hygienics Society of Japan* **47**(4), 182-188.
- Oven M, Raith K, Neubert RHH, Kutchan TM, Zenk MH (2001) Homo-phytochelatins are synthesized in response to cadmium in azuki beans. *Plant Physiology* **126**(3), 1275-1280.
- Owusu-Domfeh K, Farmer MJ, Christensen DA (1979) The amino acid composition of some Ghanaian legumes. *Ghana Journal of Agricultural Science* **12**, 75-84.
- Ozaki K (1986) Plantlet formation from the calli of primary leaf of azuki bean (*Vigna angularis*). *Japanese Journal of Breeding* **36**(4), 416-419.
- Ozawa Y (1978) On the viability of stored adzuki beans (*Phaseolus angularis*) and their hardness after cooking. *Report of the National Food Research Institute*(33), 37-40.
- Paikaray RK, Misra RC, Sahu PK (1991) Studies on niger based grain legumes intercropping system. *Orissa Journal of Agricultural Research* **4**(1-2), 44-47.
- Pajni HP, Virk N (1985) Comparative nutritional efficiency of different pulses for the development of *Tribolium castaneum* Herbst. (Coleoptera: Tenebrionidae). *Research Bulletin of the Panjab University, Science* **36**(1/2), 7-10.
- Pajni HR, Jit S (1977) Some observations on the oviposition of *Callosobruchus analis* (F.) (Bruchidae: Coleoptera). *Research Bulletin of the Panjab University, Science* **28**(1/2), 93-98.
- Pal SS, Sandhu JS, Inderjit S (2005b) Exploitation of genetic variability in interspecific cross between *Vigna mungo* × *V. umbellata*. *Indian Journal of Pulses Research* **18**(1), 9-11.
- Pan F, Wang P, Hu Z, He S, Feng X (2007) Phylogenetic study of rhizobia strains isolated from rice bean *Vigna umbellata* L.) in Hubei by 16S-23S rDNA RFLP. *Chinese Journal of Applied and Environmental Biology* **13**(1), 78-82.
- Pande HK, Jha KP (1984) Cropping systems for upland areas. 86-97.
- Pandey P, Bansal YK (1995) Culture associated genomic variations in two varieties of rice bean (*Phaseolus calcaratus* Roxy.). *Cell and Chromosome Research* **18**(1), 9-15.
- Pandey SK, Pandey RK (1986) Sterilizing effect of thiourea in relation to density of progeny population in *Callosobruchus chinensis* (L.) (Bruchidae: Coleoptera) through topical application. *Journal of Environmental Biology* **7**(1), 51-59.
- Pandey TD, Nandeha KL, Saxena RR (1998) Crop compatibility and fertility levels in maize forage crops under Bastar agroclimatic conditions. *Forage Research* **24**(1), 57-59.
- Pandiyan M, Ramamoorthi N, Ganesh SK, Jebaraj S, Pagarajan P, Balasubramanian P (2008) Broadening the genetic base and introgression of MYMV resistance and yield improvement through unexplored genes from wild relatives in mungbean. *Plant Mutation Reports* **2**(1), 33-38.

- Pandiyan M, Subbalakshmi B, Muthiah AR, Kumar M (2006a) Cytological analysis *Vigna radiata* x *V. umbellata* L. hybrids. In Plant breeding in post genomics era. Proceedings of Second National Plant Breeding Congress, Coimbatore, India, 1-3 March, 2006. In ' pp. 226) (Indian Society of Plant Breeders: Coimbatore)
- Pandiyan M, Subbalakshmi B, Muthiah AR, Kumar M (2006b) Cytological analysis *Vigna radiata* x *V. umbellata* L. hybrids. In Plant breeding in post genomics era. Proceedings of Second National Plant Breeding Congress, Coimbatore, India, 1-3 March, 2006. ' pp. 226. (Indian Society of Plant Breeders: Coimbatore)
- Panjab S, Joshi NL (1980) Intercropping of pearl millet in arid areas. *Indian Journal of Agricultural Sciences* **50**(4), 338-341.
- Panneerselvam M, Thamizhiniyan P (2008) Effect of AM fungi and dairy effluent on the growth and photosynthetic pigments of *Phaseolus trilobus* Ait. *Plant Archives* **8**(1), 171-173.
- Panneerselvam M, Thamizhiniyan P, Lenin M (2008) Biochemical activity of *Phaseolus trilobus* Ait. in AM fungi and dairy effluent. *Plant Archives* **8**(2), 611-612.
- Pareek BL, Sharma RC, Yadav CPS (1983) Record of insect faunal complex on mothbean, *Vigna aconitifolia* (Jacq.) Marechel in semi arid zone of Rajasthan. *Bulletin of Entomology* **24**(1), 44-45.
- Parihar DR (1979) Outbreak of katra, *Amsacta moorei* pest in the Rajasthan desert. *Annals of Arid Zone* **18**(1/2), 140-141.
- Park KY, Choi BH, Park SU, Kang YK, Moon HG, Lee SH (1987) Effects of continuous cultivation of upland, industrial and forage crops on soil chemical composition and crop yield. *Research Reports of the Rural Development Administration, Crops* **29**(1), 232-239. [In Korean]
- Park S, Nakajima M, Hasegawa M, Yamaguchi I (2005a) Similarities and differences between the characteristics of gibberellin-binding protein and gibberellin 2-oxidases in adzuki bean (*Vigna angularis*) seedlings. *Bioscience, Biotechnology and Biochemistry* **69**(8), 1508-1514.
- Park S, Nakajima M, Sakane M, Xu Z, Tomioka K, Yamaguchi I (2005b) Gibberellin 2-oxidases from seedlings of adzuki bean (*Vigna angularis*) show high gibberellin-binding activity in the presence of 2-oxoglutarate and Co2+. *Bioscience, Biotechnology and Biochemistry* **69**(8), 1498-1507.
- Parr MJ, Tran BMD, Simmonds MSJ, Credland PF (1998) Duration of behaviour patterns during oviposition by the bruchid beetle, *Callosobruchus maculatus*. *Physiological Entomology* **23**(2), 150-157.
- Parvathi S, Kumar VJF (2006) Value added products from rice bean (*Vigna umbellata*). *Journal of Food Science and Technology-Mysore* **43**(2), 190-193.
- Pate JS, Atkins CA, White ST, Rainbird RM, Woo KC (1980) Nitrogen nutrition and xylem transport of nitrogen in ureide-producing grain legumes. *Plant Physiology* **65**(5), 961-965.
- Patel DK, Patel PG, Patel MM (2007) Intercropping studies on rainfed castor under North Gujarat conditions. *Annals of Arid Zone* **46**(1), 99-101.
- Patel DP, Anup D, Rajesh K, Munda GC (2005) Comparative study of photosynthesis and associated parameters in major crops and weed species in mid hills of Meghalaya. *Annals of Plant Physiology* **19**(1), 1-4.
- Patel JD (1976) Comparative seed coat anatomy of some Indian edible pulses. *Phyton, Austria* **17**(3/4), 287-299.
- Patel JD (1977) Comparative anatomy of leaf, node and internode in some pulses. *Flora, German Democratic Republic* **166**(3), 193-201.
- Patel JD, Desai NC, Intwala CG, Kodappully VC (2008) Genetic variability, correlation and path analysis in moth bean. *Journal of Food Legumes* **21**(3), 158-160.
- Patel JD, Rao NV (1975) Shoot apical organization in seedlings of some Indian pulses. *Phyton, Austria* **17**(1/2), 109-119.
- Patel JR (2003) Effect of sowing dates on fodder yield of different crops during winter season. *Journal of Maharashtra Agricultural Universities* **28**(2), 181-182.
- Patel MM, Patel IC, Patel PH, Tikka SBS (2005) Arid legumes based monocropping systems for North Gujarat. **2**(1), 174-176. (Scientific Publishers (India): Jodhpur)
- Patel PN, Jindal JK (1972) Occurrence of bacterial blight disease on moth bean (*Phaseolus aconitifolius*) in India. *Indian Phytopathology* **25**(3), 464-466.
- Patel PN, Jindal JK (1973) Studies on resistance in crops to bacterial diseases in India. VI. Reaction of moth bean (*Phaseolus aconitifolius*) and hyacinth bean (*Dolichos lablab*) lines to bacterial blight and leaf spot diseases incited by *Xanthomonas phaseoli* strains. *Indian Phytopathology* **26**(3), 607-608.
- Pathik SR, Chaudhary HK, Gupta VP (1995) Estimation of additive, dominance and digenic epistatic interaction effects for certain yield characters in adzuki bean. *Indian Journal of Pulses Research* **8**(2), 172-174.
- Pathik SR, Chaudhary HK, Gupta VP (1996) Stability of seed yield and its components in adzuki bean (*Vigna angularis*) in the mid-hills of the north-western Himalayas. *Acta Agronomica Hungarica* **44**(2), 169-176.

- Patidar M, Balak R, Singh MP, Rao AS (2005) Performance of kharif legumes in different soil types of Salodi watershed, Jodhpur district, Rajasthan. **2**(1), 67-70. (Scientific Publishers (India): Jodhpur)
- Patil AS, Jadhav SK (2006a) Cross inoculation studies with rhizobia of rice bean. *Journal of Maharashtra Agricultural Universities* **31**(3), 377.
- Patil AS, Jadhav SK (2006b) Studies on correlation between crystal violet tolerance and nitrogen fixing ability of rhizobia of rice bean. *Journal of Maharashtra Agricultural Universities* **31**(3), 378.
- Patil JD, Pawar RB, Durgude AG, Takate AS, Patil VS (2004) Relative efficacy of different cropping systems and land treatments on runoff, soil erosion and crop yields under dryland conditions. *Journal of Maharashtra Agricultural Universities* **29**(1), 1-3.
- Patil SC, Patil HE, Jambhale VM (2008) Genetic divergence studies in moth bean (*Vigna aconitifolia*). *Journal of Maharashtra Agricultural Universities* **33**(2), 161-164.
- Patil SC, Patil VP, Patil HE, Pawar SV (2007) Correlation and path coefficient studies in moth bean. *International Journal of Plant Sciences (Muzaffarnagar)* **2**(2), 141-144.
- Patnayak BC, Mohan M, Bhatia DR, Hajra A (1979) A note on the nutritional value of cowpea, moth (dewgram) and clusterbean fodders fed as hay to sheep. *Indian Journal of Animal Sciences* **49**(9), 746-748.
- Patra AP (2001) Performance of rainy season legumes as influenced by phosphorus and their relative efficiency on productivity and nitrogen economy of succeeding wheat crop. *Journal of Interacademicia* **5**(4), 458-465.
- Patra AP, Ghosh P, Biswas A, Nayek SS, Ghosh SK, Saha M (2007) Growth and yield of wheat crop as influenced by nitrogen levels, previous rainy season legumes and their recycling in the gangetic alluvial soil of West Bengal. *Journal of Crop and Weed* **3**(2), 15-19.
- Pattnayak PSD, Debnath NC, Basak RK (1990) Effect of partially acidulated Purulia rock phosphate on yield of some legume and non-legume forage crops. *Environment and Ecology* **8**(3), 1008-1011.
- Paulsamy S, Lakshmanachary AS, Manian S (1987) Effect of overgrazing on the phytosociology of a tropical grassland ecosystem. *Indian Journal of Range Management* **8**(2), 103-107.
- Pawar VD, Ingle UM (1987) Production of quick-cooking moth bean (*Phaseolus aconitifolius* Jacq.) 3. Effect of soaking on activity [of] trypsin inhibitor. *Indian Journal of Nutrition and Dietetics* **24**(12), 376-379.
- Pawar VD, Machewad GM, Durge AV, Maitre AS (2009) Processing and characteristics of snacks extruded from rice and corn grits and two malted legumes. *Journal of Food Science and Technology (Mysore)* **46**(5), 494-496.
- Pengelly BC, Blamey FPC, Muchow RC (1999) [Radiation interception and the accumulation of biomass and nitrogen by soybean and three tropical annual forage legumes](#). *Field Crops Research* **63**(2), 99-112.
- Pengelly BC, Muchow RC, Blamey FPC (1999) [Predicting leaf area development in response to temperature in three tropical annual forage legumes](#). *Australian Journal of Agricultural Research* **50**(2), 253-259.
- Peregrine WTH (1983) Chaya (*Cnidoscolus-aconitifolius*) - a potential new vegetable crop for Brunei. *Tropical Pest Management* **29**(1), 39-41.
- Perrino P, Shagarovsky T, Esquivel M, Uranga H, Hammer K (1992) The cultivated races of *Vigna SAVI* in Cuba. *Feddes Repertorium* **103**(7-8), 509-514.
- Perumal S (2006) [The antioxidant activity and free radical-scavenging capacity of phenolics of raw and dry heated moth bean \(\*Vigna aconitifolia\*\) \(Jacq.\) Marechal seed extracts](#). *Food Chemistry* **99**(1), 149-157.
- Pesci P, Beffagna N (1983a) ATPase, ADPase and p-nitrophenylphosphatase activities in microsomal preparations from various plant materials. I. Effects of the presence of magnesium in the homogenization medium. *Atti della Accademia Nazionale dei Lincei, Rendiconti* **74**(1), 34-40.
- Pesci P, Beffagna N (1983b) ATPase, ADPase and p-nitrophenylphosphatase activities in microsomal preparations from various plant materials. II. ATPase, ADPase and acid phosphatase activity and effects of inhibitors and solubilizing treatments. *Atti della Accademia Nazionale dei Lincei, Rendiconti* **74**(2), 99-108.
- Peterbauer T, Brereton I, Richter A (2003) Identification of a digalactosyl ononitol from seeds of adzuki bean (*Vigna angularis*). *Carbohydrate Research* **338**(19), 2017-2019.
- Peterbauer T, Mucha J, Mayer U, Popp M, Glössl J, Richter A (1999) [Stachyose synthesis in seeds of adzuki bean \(\*Vigna angularis\*\): molecular cloning and functional expression of stachyose synthase](#). *Plant Journal* **20**(5), 509-518.
- Peterbauer T, Puschenreiter M, Richter A (1998) Metabolism of galactosylononitol in seeds of *Vigna umbellata*. *Plant and Cell Physiology* **39**(3), 334-341.
- Peterbauer T, Richter A (1998) Galactosylononitol and stachyose synthesis in seeds of adzuki bean. Purification and characterization of stachyose synthase. *Plant Physiology* **117**(1), 165-172.

- Petrova MV (1978a) Node and internode structure in species of Phaseolus and Vigna from the collection of the Vavilov Institute of Plant Industry (VIR). *Byulleten' Vsesoyuznogo Ordena Lenina i Ordena Druzhby Narodov Instituta Rastenievodstva Imeni N. I. Vavilova*(83), 81-90.
- Phansak P, Taylor PWJ, Mongkolporn O (2005) Genetic diversity in yardlong bean (*Vigna unguiculata* ssp *sesquipedalis*) and related *Vigna* species using sequence tagged, microsatellite site analysis. *Scientia Horticulturae* **106**(2), 137-146.
- Phillips N, Imoto S, Ohta Y, Nakanishi K (1992) Partial purification of an endogenous elicitor from suspension-cultured cells of red bean, *Vigna angularis*. *Experientia* **48**(7), 683-687.
- Pietsch SA (1996) Phloem transport: A new 'bidirectional' exudation method. *Journal of Experimental Botany* **47** (Spec. Issue), 1328.
- Pimentel C, Hébert G (1999) Photosynthetic potential and stomatal conductance in bean and cowpea genotypes under water stress. *Revista Brasileira de Fisiologia Vegetal* **11**(1), 7-11. [In Portuguese]
- Pintarak A, Juntorn W (1996a) Improved fallow management. 177-187.
- Pol KM, Awari VR, Ugale SD, Thakur DR (2001b) Periodical correlations of various morpho-physiological and yield contributing characters with grain yield in rice-bean (*Vigna umbellata* (L.)). *Annals of Plant Physiology* **15**(2), 159-162.
- Poonia KL (2004a) Impact of khejri (*Prosopis cineraria*) on soil fertility and its effect on growth and yield attributes of rainfed kharif crops. *Current Agriculture* **28**(1/2), 95-97.
- Poonia KL (2004b) Intercropping of rainfed kharif crops with khejri (*Prosopis cineraria* Linn.). *Current Agriculture* **28**(1/2), 91-93.
- Porpavai S, Palchamy A, Boopathi SNMR, Jayapaul P (2005) [Feasibility studies on intercropping green manures with grain legumes and performance of ratooned green manure in pre-rice season and its effect on rice yield](#). *Legume Research* **28**(4), 303-305.
- Pradhan AC, Samanta G, Agasti MK (2000) Effect of phosphate nutrition and harvesting schedule on the production of forage and mineral matter in rice bean [*Vigna umbellata* (Thunb.) Ohwi & Ohashi] in terai region of West Bengal. *Indian Agriculturist* **44**(1/2), 31-34.
- Prakash BG, Khanure SK (1999b) Isolation of mutants and their frequencies under M2 generation in rice bean (*Vigna umbellata* L. Thunb.). *Madras Agricultural Journal* **86**(10/12), 568-572.
- Prakash BG, Kumar P, Veeresh LC, Shambulingappa KG (1993) Insects and mite pests of rice bean. *Current Research - University of Agricultural Sciences (Bangalore)* **22**(3/5), 57-59.
- Prakash BG, Shambulingappa KG (2000) Effect of gamma rays and EMS on biological end points and estimation of LD<sub>50</sub> value in rice bean. *Karnataka Journal of Agricultural Sciences* **13**(1), 155-157.
- Prakash BG, Shambulingappa KG (1999) Estimation of chlorophyll and viable mutation under M2 generation in rice bean (*Vigna umbellata* L. Thunb.). *Karnataka Journal of Agricultural Sciences* **12**(1/4), 38-42.
- Pramanik A, Somchoudhury AK, Khatua DC (2000) Persistent toxicity of *Bacillus thuringiensis* var. kurstaki to *Spilosoma obliqua* Walker on different host plants under field conditions. *Journal of Entomological Research (New Delhi)* **24**(1), 91-95.
- Pranab R, Nisan B (1987) Reiterated DNA sequences in the genome of mung bean and its cloning. *Plant Physiology & Biochemistry, India* **14**(1), 6-13.
- Prasad G, Bhol BB (1994b) Uptake of nitrogen, phosphorus, potash and calcium by rice bean (*Vigna umbellata*, Thunb, Ohwi and Ohashi) as influenced by row spacing and levels of nitrogen. *Orissa Journal of Agricultural Research* **7**(Supplement), 105-107.
- Prasad G, Bhol BB, Kar BC, Mishra SN (1994a) Response of rice bean (*Vigna umbellata*) to spacing and levels of nitrogen. *Indian Journal of Agronomy* **39**(3), 485-487.
- Prasad KVV, Agrawal SC (1992) Leaf blight of ricebean caused by *Macrophomina phaseolina*. *Indian Journal of Pulses Research* **5**(1), 93.
- Prasad MSL, Prasad MS, Sharma YP, Sangit K (2005) Important diseases of crops in NEH region and their management. In '' Ed. D Prasad) pp. 182-204)
- Prasad MSV, Inamdar JA (1991) Effect of cement kiln dust pollution on growth and yield of *Vigna* spp. *Indian Journal of Ecology* **18**(2), 91-94.
- Prasad MVR, Singh RP (1973) Select your crop for Western Rajasthan. *Indian Farming* **22**(10), 13-15.
- Prati S, Baravelli V, et al. (2007) Composition and content of seed flavonoids in forage and grain legume crops. *Journal of Separation Science* **30**(4), 491-501.
- Praveen K, Kumar D, Tarafdar JC (2000) Reflection of genetic variation in grain yield of moth bean (*Vigna aconitifolia* Jacq. Marechal) on some biochemical properties of soil. *Journal of the Indian Society of Soil Science* **48**(3), 619-621.

- Praveen K, Tarafdar JC (1997) [Rhizobium-induced changes on nitrate reductase activity in rhizosphere and phyllosphere](#). *Current Science* **73**(2), 204-205.
- Praveen K, Tripathi KP, Aggarwal RK (2002) Influence of crops, crop residues and manure on amino acid and amino sugar fractions of organic nitrogen in soil. *Biology and Fertility of Soils* **35**(3), 210-213.
- Praveen S, Manigopa C, Singh DN (2005) Correlation and path analysis for symbiotic improvement in Vigna species. In '' pp. 5-7. (Scientific Publishers (India): Jodhpur)
- Preeti S, Sanjana K, Dhar MK (2006b) [Can phaseolin patterns help resolve the Phaseolus-Vigna complex?](#) *Genetic Resources and Crop Evolution* **53**(8), 1573-1578.
- Prema G, Dhumale DB (1990a) Study of genetic variability in ricebean. *Annals of Plant Physiology* **4**(1), 92-95.
- Prema G, Dhumale DB (1990b) Study of path analysis in ricebean (*Vigna umbellata*). *Annals of Plant Physiology* **4**(2), 211-214.
- Prema G, Dhumale DB, Raut RS (1990) Interrelationship between yield and its components in rice bean (*Vigna umbellata*). *Indian Journal of Agricultural Sciences* **60**(8), 547-549.
- Premchand (1981) Presence of feeding deterrent in velvet bean, *Mucuna cochinchensis* Roxb. *Indian Journal of Entomology* **43**(2), 217-219.
- Price ML, Hagerman AE, Butler LG (1980) [Tannin content of cowpeas, chickpeas, pigeon peas and mung beans](#). *Journal of Agricultural and Food Chemistry* **28**(2), 459-461.
- Purwar JP (2009) Varietal preference of *Spilarctia obliqua* among pulses. *Annals of Plant Protection Sciences* **17**(2), 490-491.
- Purwar JP, Anjuli A, Vishwanath (2007b) Evaluation of host preference of *Spodoptera litura* (Fabricius) among pulses grown in Uttarakhand hills. *Pantnagar Journal of Research* **5**(2), 100-102.
- Pushpa G, Mahishi DM, Satyan BA, Kulkarni RN (1979) Karyomorphology of three species of Phaseolus. *Mysore Journal of Agricultural Sciences* **13**(2), 147-151.
- Puttaswamy, Gowda BLV, Ali TMM (1977) Record of pests infesting moth bean (matki) (*Phaseolus aconitifolius* Jacq.) a potential pulse crop. *Current Research* **6**(4), 69-71.
- Qasim M, Ahmad K, Shah NZ, Razaullah (1994) Studies on the growth and nodulation in some kharif legumes and their effects on soil properties. *Sarhad Journal of Agriculture* **10**(3), 231-236.
- Qi XQ, Chen RY, Wilson KA, Tan-Wilson AL (1994) Characterization of a soybean  $\beta$ -conglycinin-degrading protease cleavage site. *Plant Physiology* **104**(1), 127-133.
- Qi YT, Burkholder WE (1985) Study on sex pheromone biology of the Azuki bean weevil, *Callosobruchus chinensis* (L.). *Contributions from Shanghai Institute of Entomology* **5**, 1-11.
- Qin Y (1999) The grey correlation analysis between contents of some heavy metals in seeds and irrigation water or soil. *Journal of China Agricultural University* **4**(Supplement), 98-102.
- Radhamani J, Vishnol SR, Devi LC, Singh AK (2007) Influence of antioxidants in delaying ageing of arid and non-arid legume seeds. *Journal of Arid Legumes* **4**(2), 77-81.
- Rahman FH, Najar GR (2006) Preparation of phosphocompost from paddy straw-water hyacinth mixture alongwith Mussoorie rock phosphate and its effect on dry matter yield and P uptake by paddy (*Oryza sativa* L.) and rice bean (*Vigna umbellata* Roxb.). *Crop Research (Hisar)* **32**(1), 1-5.
- Rai AK, Khanna ND, Agarwal SP (1991) Effect of feeding *Leucaena leucocephala* with *Phaseolus aconitifolius* on growth and thyroid status of camel calves. 152-156.
- Rai AK, Khanna ND, Agarwal SP (1992) Effect of feeding *Leucaena leucocephala* with *Phaseolus aconitifolius* on growth and thyroid status of camel calves. *Indian Journal of Animal Sciences* **62**(4), 297-301.
- Raiger HL, Dua RP, Phogat BS, Mishra SK (2008b) Assessment of genotype  $\times$  environment interaction in rice bean (*Vigna umbellata*) under Indian plains. *Indian Journal of Agricultural Sciences* **78**(6), 548-550.
- Raj S, Bhagwan S (2005) Response of cultivars and weed management practices on the weeds and yield of moth bean. **2**(1), 28-31. (Scientific Publishers (India): Jodhpur)
- Rajan K, David, Bhanu P, Naveen K, Bhandari JC (2007b) Fodder yield and nutritional attributes of ricebean (*Vigna umbellata*). *Range Management and Agroforestry* **28**(2B), 260-262.
- Rajan S (2009) Genetic differentiation of host limited forms of *Colletotrichum truncatum* from northwestern Himalayas. *Archives of Phytopathology and Plant Protection* **42**(10), 960-966.
- Rajan S, Kaushal RP (1999) Characterization of pathogenic variability in *Colletotrichum truncatum* in Himachal Pradesh. *Indian Phytopathology* **52**(4), 389-392.
- Rajaram N, Janardhanan K (1990b) Chemical analysis and nutritional evaluation of certain under-exploited *Vigna* spp. *Food Sciences and Nutrition* **42F**(4), 213-221.
- Rajen C, Tamang JP (2008b) Microbiological evaluation of maseura, an ethnic fermented legume-based condiment of Sikkim. *Journal of Hill Research* **21**(1), 1-7.

- Rajender S (1994) Pearl millet based intercropping in arid zone of Rajasthan. *Annals of Arid Zone* **33**(2), 155-156.
- Rajender S (1995) Mixed cropping in pearl millet (*Pennisetum glaucum*) in arid zone of Rajasthan. *Indian Journal of Agricultural Sciences* **65**(4), 276-278.
- Rajpurohit TS (1983) Control of root rot of cotton caused by *Rhizoctonia bataticola* (Taub) Butler. *Madras Agricultural Journal* **70**(11), 751-752.
- Rajpurohit TS (2002) Influence of intercropping and mixed cropping with pearl millet, green gram and mothbean on the incidence of stem and root rot (*Macrophomina phaseolina*) of sesame. *Sesame and Safflower Newsletter*(17), 40-41.
- Rajput SS, Jakhar ML (2005) Influence of kanamycin on explant and callus of moth bean. *Indian Journal of Plant Physiology* **10**(2), 115-119.
- Raka S, Aakanksha N (2005) Comparative estimation of nutrients of moth bean using two different methods of cooking. **2**(1), 196-200. (Scientific Publishers (India): Jodhpur)
- Ram S, Dalip K, Suhag KS, Hasija RC (2005) Comparative economics of pulse crops in Rajasthan State. *Environment and Ecology* **23**(Spl-1), 20-24.
- Ramachandra C, Shivaraj B, Gowda A (1993) Studies on the influence of intercrops grown for forage and seed on the seed yield and quality of fodder maize. *Farming Systems* **9**(3-4), 87-92.
- Ramakrishna YS, Singh RP (1978) Influence of climate on crop production in drylands. *Proceedings of the Indian National Science Academy, B* **44**(6), 431-436.
- Ramashrit S, Sarup P (1985a) Effect of insecticides on the eggs of maize stalk borer, *Chilo partellus* (Swinhoe) reared on artificial diet and natural host-plant maize. *Journal of Entomological Research* **9**(1), 1-3.
- Ramashrit S, Sarup P (1985b) Formulation of artificial diets in the laboratory for mass rearing of maize stalk borer, *Chilo partellus* (Swinhoe). *Journal of Entomological Research* **9**(2), 115-121.
- Ramashrit S, Sarup P (1986a) Effect of natural and artificial diets on the susceptibility of moths of the maize stalk borer, *Chilo partellus* (Swinhoe) (Lepidoptera: Crambidae) to different insecticides. *Journal of Entomological Research* **10**(1), 51-56.
- Ramashrit S, Sarup P (1986b) Significance and optimal quantity of ascorbic acid in artifical diets for the mass rearing of maize stalk borer, *Chilo partellus* (Swinhoe) (Lepidoptera: Crambidae). *Journal of Entomological Research* **10**(2), 174-179.
- Ramashrit S, Sarup P (1987) An improved alternative artificial diet for mass rearing of the maize stalk borer, *Chilo partellus* (Swinhoe) (Lepidoptera: Crambidae). *Insect Science and its Application* **8**(2), 181-185.
- Ramzan M, Chahal BS, Judge BK (1990b) Storage losses to some commonly used pulses caused by pulse beetle, *Callosobruchus maculatus* (Fab.). *Journal of Insect Science* **3**(1), 106-108.
- Rana SK, Roquib MA, Mukherjee AK, Chattopadhyay P (1994) Fodder production of Sorghum bicolor on alluvial soil of West Bengal at rainfed conditions as short duration crop. *Environment and Ecology* **12**(1), 14-17.
- Rangasamy SR, Muralidharan V, Rathnasamy R (1988) Breeding behavior of *Vigna radiata*-*V. mungo* hybrid progenies for crop improvement in mungbean. 187-199.
- Ranjit K, Satija CK (1996b) Study of incompatibility barriers between interspecific hybrids of *Vigna radiata* × *V. umbellata*. *Crop Improvement* **23**(2), 213-216.
- Ranjit K, Satija CK (1998b) Cytogenetical and biochemical analysis of interspecific hybrids between *Vigna radiata* and *V. umbellata*. *Indian Journal of Genetics & Plant Breeding* **58**(1), 25-34.
- Rao AS, Singh RS, Joshi NL, Ramakrishna YS (1999) Evapotranspiration requirements of some arid zone crops. 451-456.
- Rao AV (2005a) Strategies for achieving sustainable production of arid legumes. **2**(1), 127-131. (Scientific Publishers (India): Jodhpur)
- Rao AV, Bala K, Tarafdar JC (1990) [Dehydrogenase and phosphatase activities in soil as influenced by the growth of arid-land crops](#). *Journal of Agricultural Science* **115**(2), 221-225.
- Rao AV, Venkateswarlu B (1983) Pattern of nodulation and nitrogen fixation in mothbean. *Indian Journal of Agricultural Sciences* **53**(12), 1035-1038.
- Rao AV, Venkateswarlu B (1987a) Nitrogen fixation as influenced by water stress in selected crop legumes of the Indian arid zone. *Arid Soil Research and Rehabilitation* **1**(2), 89-96.
- Rao AV, Venkateswarlu B (1987b) Use of biofertilisers in the desert ecosystem. *Indian Farming* **37**(1), 21-22.
- Rao AV, Venkateswarlu B, Henry A (1984) Genetic variation in nodulation and nitrogenase activity in guar and moth. *Indian Journal of Genetics and Plant Breeding* **44**(3), 425-428.
- Rao JAVP, Gopalachari NC (1982) Effect of different crop sequences under rainfed conditions with reference to F.C.V. tobacco on the chemical and physical quality characteristics of tobacco leaf. *Tobacco Research* **8**(2), 148-159.

- Rao KJ, Thirumalachar DK (1977) [New record of alternate host plants of groundnut leaf miner \*Stomopteryx subsecivella\* zeller \(syn.: \*S. nerteria\* Meyrick\) \(Lepidoptera: Gelechiidae\)](#). *Current Science* **46**(3), 91-92.
- Rao KN, Rao AS, Reddy R (1985) Effect of different herbicides on control of Cuscuta in legumes. *Pesticides* **19**(6), 61-62.
- Rao KR, Shylesha AN, Thakur NSA (2005) Status and distribution of insect pests on crops and their management by organic methods in North Eastern Hill Region of India. In ' Ed. D Prasad pp. 104-126)
- Rao NSS (1980) Role of bacteria in crop production. *Indian Farming* **30**(7), 71-76.
- Rao SR, Chandel KPS (1991b) [Karyomorphological studies in the cultivated and wild \*Vigna\* species in Indian gene centre](#). *Cytologia* **56**(1), 47-57.
- Rao SR, Raina SN (2005) Cytological evaluation of colchitetrapiosity in moth bean (*Vigna aconitifolia*) and its allied species. *2*(2), 389-396.
- Rao TP, Yano K, Yamauchi A, Tatsumi J, Iijima M (2001) Regulation of phosphorus solubilization by light conditions on shoot in legumes. In 'Developments in Plant and Soil Sciences, Volume 92.' pp. 558-559. (Kluwer Academic Publishers: Dordrecht)
- Rashid KA, Smartt J, Haq N (1988b) Hybridization in the genus *Vigna*. 205-214.
- Rashmi V (2002) Rhizoctonia bataticola is a serious pathogen in the seeds of *Vigna aconitifolia*. *Journal of Phytopathological Research* **15**(1), 81-83.
- Rath BS, Tripathy RK, Nayak SP, Kar BC (1996b) Effect of different row spacings on grain yield of rice bean (*Vigna umbellata* (Thunb) Ohwi and Ohashi) varieties. *Environment and Ecology* **14**(2), 463-465.
- Rath BS, Tripathy RK, Nayak SP, Kar BC (1997) Effect of row spacing on yield and seed quality of rice bean (*Vigna umbellata*) varieties. *Indian Journal of Agronomy* **42**(4), 717-719.
- Rathore BS (1995) Effect of seed dressers on bacterial leaf blight of mothbean (*Vigna aconitifolia*). *Plant Disease Research* **10**(1), 65-68.
- Rathore BS (1997) Relative efficacy of fungicidal seed treatments in the management of root rot and seedling blight of mothbean. *Plant Disease Research* **12**(2), 157-159.
- Rathore BS (2000a) Effect of organic amendments on incidence of seed rot and seedling blight of mothbean. *Plant Disease Research* **15**(2), 217-219.
- Rathore BS (2000b) Evaluation of mothbean entries against seed rot and seedling blight. *Journal of Mycology and Plant Pathology* **30**(1), 126-127.
- Rathore BS (2001) Screening of mothbean genotypes against root rot and seedling blight caused by Macrophomina phaseolina. *Plant Disease Research* **16**(1), 110-112.
- Rathore BS, Chhipa HP (1996) Varietal interaction of mothbean to Rhizobium inoculation. *Indian Journal of Pulses Research* **9**(1), 55-58.
- Rathore BS, Rathore RS (1999) Effect of seed dressers on Macrophomina root rot of mothbean. *Journal of Mycology and Plant Pathology* **29**(3), 389-392.
- Rathore GS, Agnihotri JP (1985) Effect of insecticides on yellow mosaic of moth (*Vigna aconitifolia* (Jacq.) Marechal). *Indian Journal of Virology* **1**(1), 92-94.
- Rathore GS, Sharma RC, Agnihotri JP, Gour HN, Sharma RK (1986) Evaluation of moth bean germplasm for resistance to yellow mosaic. *Indian Journal of Virology* **2**(1), 81-86.
- Rathore MS, Bhati DS, Chandawat MS, Singh I, Chawra RS (2006) Performance of arid legumes and their intercropping system with pearl millet to sustain productivity in arid region of Rajasthan. *Research on Crops* **7**(2), 375-377.
- Rathore MS, Singh I, Chandawat MS (2003) Effect of weed control practices on productivity of dew bean [*Vigna aconitifolia* (Jacq.) Marechal] cultivar RMO 40 under rainfed conditions. *Crop Research (Hisar)* **26**(1), 30-32.
- Rathore SS, Pruthi NK, Naruka IS (2009) Alleviation of sodium chloride stress by growth regulators in seedlings of mothbean [*Vigna aconitifolia* (Jacq.) Marechal]. *Crop Research (Hisar)* **37**(1/3), 40-45.
- Rathore VS, Singh JP, Beniwal RK (2007) Inhibition of germination and seedling growth of crops by aqueous extracts of shrubs. *Annals of Arid Zone* **46**(2), 169-173.
- Rau BHS, Ramana KVR, Singh N (1972) [Studies on nutritive value of leaf proteins and some factors affecting their quality](#). *Journal of the Science of Food and Agriculture* **23**(2), 233-245.
- Raut RS, Dhumale DB, Lokhande VE (1989) Induction of callus and regeneration of plants in rice bean (*Vigna umbellata* Thunb). *Annals of Plant Physiology* **3**(1), 25-28.
- Ravinder G, Eapen S (1986) [Plant regeneration from hypocotyl protoplasts of mothbean \(\*Vigna aconitifolia\*\)](#). *Current Science, India* **55**(2), 100-102.

- Ravindran KC, Indrajith A, Balakrishnan V, Venkatesan K, Kulandaivelu G (2008) Determination of defense mechanism in Phaseolus trilobus Ait. seedlings treated under UV-B radiation. *African Crop Science Journal* **16**(2), 111-118.
- Rawat RR, Sahu HR (1973) A study on the bionomics of Chauliops fallax Scott (Heteroptera: Lygaeidae) at Sehore (Madhya Pradesh). *Journal of the Bombay Natural History Society* **70**(3), 475-479.
- Redden B (1997) New ACIAR project report - adzuki bean: evaluation and utilisation of germplasm in Australia and China. *ACIAR Food Legume Newsletter*(25), 3-5.
- Redden R, Desborough P, Page J, Twyford-Jones P (2000) Adzuki beans: from production to marketing - part 2. *Agricultural Science* **13**(3), 24-27.
- Redden R, Desborough P, Tompkins W, Usher T, Kelly A (2001) [Responses of adzuki bean as affected by row spacing, plant density and variety](#). *Australian Journal of Experimental Agriculture* **41**(2), 235-243.
- Redden RJ, Basford KE, Kroonenberg PM, Islam FMA, Ellis R, Wang S, Cao Y, Zong X, Wang X (2009) Variation in adzuki bean (*Vigna angularis*) germplasm grown in China. *Crop Science* **49**(3), 771-782.
- Reddy BVR, Chakrapani C, Babu CS, Rao KS, Agarwal K, Prasad M (2010) FT-IR, TG, XRD and SEM studies of activated carbons prepared from agricultural waste. *Asian Journal of Chemistry* **22**(3), 1822-1828.
- Reddy MJ, Setty TKP, Jagannatha V, Herle PS (1993) Studies on methods of crop establishment in rice fallows in coastal Karnataka. *Mysore Journal of Agricultural Sciences* **27**(3), 205-208.
- Reddy PVR, Todoriki S, Miyanoshita A, Imamura T, Hayashi T (2006) Effect of soft electron treatment on adzuki bean weevil, *Callosobruchus chinensis* (L.) (Col., Bruchidae). *Journal of Applied Entomology* **130**(6/7), 393-399.
- Reddy VC, Murthy KNK, Shankaranarayana V (2005) Influence of legume intercropping in South African maize under rainfed conditions. *Environment and Ecology* **23**(Spl-1), 195-197.
- Reeder RT, Whitmarsh J, Gray LE, Peterson RE (1986) Inhibition of photosynthetic electron transport by metabolites produced by *Phialophora gregata*. *Physiological and Molecular Plant Pathology* **28**(3), 371-379.
- Rekha D, Dutta SK (1996) Effect of intercropping on infestation of insect pests of green gram. *Journal of the Agricultural Science Society of North East India* **9**(2), 220-223.
- Rele MV, Vartak HG, Jagannathan V (1980) Proteinase inhibitors from *Vigna unguiculata* subsp. *cylindrica*. 1. Occurrence of thiol proteinase inhibitors in plants and purification from *Vigna unguiculata* subsp. *cylindrica*. *Archives of Biochemistry and Biophysics* **204**(1), 117-128.
- Rerkasem B, Rerkasem K, Peoples MB, Herridge DF, Bergersen FJ (1988) [Measurement of N-2 fixation in maize \(Zea-mays L\) ricebean \(\*Vigna-umbellata\* Thunb Ohwi and Ohashi\) intercrops](#). *Plant and Soil* **108**(1), 125-135.
- Rerkasem K, Rerkasem B (1988) [Yields and nitrogen nutrition of intercropped maize and ricebean \(\*Vigna-umbellata\* Thunb Ohwi and Ohashi\)](#). *Plant and Soil* **108**(1), 151-162.
- Rerkasem K, Rerkasem B, Wongwatana S Nitrogen nutrition in legume/non-legume intercrops. 1987, p. 278
- Revilleza MJR, Mendoza EMT, Raymundo LC (1990) [Oligosaccharides in several philippine indigenous food legumes - determination, localization and removal](#). *Plant Foods for Human Nutrition* **40**(1), 83-93.
- Riaz A, Abdul J, Azraf-ul-Haq A, Ehsanullah, Bhatti IH (2007) Evaluation of direct-seeded upland rice-based intercropping systems under strip planting geometry. *Pakistan Journal of Agricultural Sciences* **44**(2), 189-193.
- Richter A, Hoch G, Puschenreiter M, Mayer U, Peterbauer T The role of stachyose synthase in the oligosaccharide metabolism of legume seeds. 2000, Wallingford, pp. 75-84
- Richter A, Peterbauer T, Brereton I (1997) Structure of galactosylononitol. *Journal of Natural Products* **60**(8), 749-754.
- Rifat H, Safdar A (2004) Potential of summer legumes to fix nitrogen and benefit wheat crop under rainfed condition. *Journal of Agronomy* **3**(4), 273-281.
- Ritu M, Satish L (2006) Relative efficacy of on-farm weeds as soil-amendment for managing dry root rot of clusterbean in an arid environment. *Phytopathologia Mediterranea* **45**(3), 215-224.
- Robinson H (1993) A review of the genus Critoniopsis in central and south-America (Vernonieae, Asteraceae). *Proceedings of the Biological Society of Washington* **106**(3), 606-627.
- Robinson RG (1980) Registration of Minoka adzuki bean (Reg. No. 19). *Crop Science* **20**(4), 549.
- Robinson RG (1983) Yield and composition of field bean and adzuki bean in response to irrigation, compost, and nitrogen. *Agronomy Journal* **75**(1), 31-35.
- Rochester IJ, Peoples MB, Constable GA, Gault RR (1998) [Faba beans and other legumes add nitrogen to irrigated cotton cropping systems](#). *Australian Journal of Experimental Agriculture* **38**(3), 253-260.

- Roddick JG, Guan M (1991) Brassinosteroids and root development. 231-245.
- Rodge AB, Ghatge PU, Wankhede DB (2006) Studies on reducing tannin and phytate contents in horsegram (*Dolichos biflorus*) and mothbean (*Vigna aconitifolia*) by soaking and germination. *Journal of Arid Legumes* **3**(2), 28-30.
- Rodge AB, Wankhede DB, Ghatge PU (2005) Preparation, characterization and utilization of protein isolates from mothbean. **2**(2), 348-350.
- Rodriguez FM, Tecson-Mendoza EM (1998) Alkaloids of several Philippine indigenous food legumes: determination and removal. *Philippine Journal of Crop Science* **23**(2), 121-125.
- Rodriguez MS, Mendoza EMT (1991) Nutritional assessment of seed proteins in rice bean *Vigna-umbellata* (thumb) Ohwi and Ohashi. *Plant Foods for Human Nutrition* **41**(1), 1-9.
- Roh S, Tsuchizaki T, Hibi T (1996) Purification, partial characterization and serology of a new strain of azuki bean mosaic potyvirus. *Annals of the Phytopathological Society of Japan* **62**(3), 243-246.
- Roman AD, Herman-y-Lara E, Salgado-Cervantes MA, Garcia-Alvarado MA (2004) Food sorption isotherms prediction using the ross equation. *Drying Technology* **22**(8), 1829-1843.
- Roquib MA, Das M (1995) Genetic divergence in rice bean (*Vigna umbellata*). *Advances in Plant Sciences* **8**(2), 312-318.
- Rossi CE, Ferraz LCCB (2001) Effect of different host plants on the morphometrics of a Brazilian population of *Heterodera glycines*. *Arquivos do Instituto Biológico (São Paulo)* **68**(1), 95-102. [In Portuguese]
- Rout D, Pradhan L, Barik T, Misra SN (1989) Studies on pure stand and cereal-legume association of maize, sorghum, cowpea and ricebean in different proportions. *Orissa Journal of Agricultural Research* **2**(2), 85-90.
- Rout D, Pradhan L, Barik T, Misra SN (1990) Studies on pure stand and cereal-legume association of maize, sorghum, cowpea and rice bean in different proportions. *Indian Agriculturist* **34**(1), 41-46.
- Roy S (1994) Ovipositional preference and development of *Callosobruchus maculatus* Fabricius on seeds of promising dry land legumes. *Agricultural Science Digest* **14**(3-4), 177-180.
- Roychoudhury B, Swagata G, Mandal N, Dasgupta T (2007a) Genetic diversity in some wild and cultivated germplasms in *Vigna* by RAPD marker. Recent trends in horticultural biotechnology, Vol. II. ICAR national symposium on biotechnological interventions for improvement of horticultural crops: issues and strategies. Vellanikkara, Kerala, India, 10-12 January, 2005. In '' (Eds R Keshavachandran, P Nazeem, D Girija, PS John and KV Peter) pp. 515-523) (New India Publishing Agency: New Delhi)
- Rozan P, Kuo Y, Lambein F (2000) Free amino acids present in commercially available seedlings sold for human consumption. A potential hazard for consumers. *Journal of Agricultural and Food Chemistry* **48**(3), 716-723.
- Rudragouda, Angadi SS (2002) Impact of spacings and fertility levels on yield and quality of ricebean (*Vigna umbellata*) genotypes for forage production. *Karnataka Journal of Agricultural Sciences* **15**(4), 685-687.
- Rudragouda, Angadi SS, Hongal MM (2008a) Influence of genotypes, spacings and fertility levels on growth and yield of ricebean [*Vigna umbellata* (Thunb) Ohwi and Ohashi] for fodder production. *Crop Research (Hisar)* **35**(1/2), 153-154.
- Rudragouda, Angadi SS, Rajendra H (2005) Effect of genotypes, spacings and fertility levels on growth and yield of ricebean fodder. *Indian Journal of Pulses Research* **18**(2), 248-249.
- Ruelo J (1980) Countering nematode threat. *Centerpoint*((Fall 1980)), 4.
- Ruma S, Vimla D (2009) Development and nutritional analysis of value added spread instant mix. *Journal of Human Ecology* **28**(3), 187-190.
- Ryoo MI, Chun YS (1993) Oviposition behavior of *Callosobruchus chinensis* (Coleoptera: Bruchidae) and weevil population growth: effects of larval parasitism and competition. *Environmental Entomology* **22**(5), 1009-1015.
- Saab IN, Sachs MM (1995) Complete cDNA and genomic sequence encoding a flooding-responsive gene from maize (*Zea mays* L.) homologous to xyloglucan endotransglycosylase. *Plant Physiology* **108**(1), 439-440.
- Sadana B, Hira CK, Singla N, Grewal H (2006) Nutritional evaluation of rice bean (*Vigna umbellata*) strains. *Journal of Food Science and Technology-Mysore* **43**(5), 516-518.
- Saddler GS (2000) Xanthomonas pisi. [Descriptions of Fungi and Bacteria]. (146), Sheet 1459.
- Saha AK, Setua GC (2006) Effect of integrated nutrient management on quality of mulberry leaves assessed through bioassay. *Uttar Pradesh Journal of Zoology* **26**(3), 287-291.
- Saha DK, Painuli DK, Dhariwal R (2005) Dissemination of technologies and its impact on desert dwellers. **2**(2), 375-380.

- Saharan K, Khetarpaul N, Bishnoi S (2001a) HCl-extractability of minerals from ricebean and fababean: influence of domestic processing methods. *Innovative Food Science & Emerging Technologies* **2**(4), 323-325.
- Saharan K, Khetarpaul N, Bishnoi S (2001b) Processing of newly released ricebean and fababean cultivars: changes in total and available calcium, iron and phosphorus. *International Journal of Food Sciences and Nutrition* **52**(5), 413-418.
- Saharan K, Khetarpaul N, Bishnoi S (2004) Contents and digestibility of carbohydrates of ricebean and fababean as affected by simple inexpensive processing methods. *Nutrition & Food Science* **34**(1), 13-16.
- Sahay G, Sarma BK, Gupta HS, Pathak KA, Prasad MS (1999) Biotic stresses of pulses in North Eastern Hill region of India. *Indian Journal of Hill Farming* **12**(1/2), 8-16.
- Sahu JP, Singh NP, Sharma BB (1995) Performance of ricebean genotypes under varying row spacings. *Indian Journal of Pulses Research* **8**(2), 146-148.
- Sahu PK, Mishra TK, Singh GS, Mishra CHP, Mishra RC (2007) Genetic diversity, character association and path-coefficient study in rice bean. *Environment and Ecology* **25**(Special 3A), 808-812.
- Sahu PK, Misra RC, Jali CR (2008b) Induction of micromutations and their role in affecting character association pattern in rice bean (*Vigna umbellata*). *Environment and Ecology* **26**(1), 178-182.
- Sahu TD, Mandal SR (2004a) Effect of various sources of phosphate on nutrient composition of green manure crops grown during winter season and improvement in soil fertility. *Environment and Ecology* **22**(3), 534-537.
- Sahu TD, Mandal SR (2004b) Influence of different sources of phosphate on micronutrient content of green manure crops and their availability during winter season. *Environment and Ecology* **22**(2), 424-426.
- Saikia P, Sarkar CR, Borua I (1999) Chemical composition, antinutritional factors and effect of cooking on nutritional quality of rice bean *Vigna umbellata* (Thunb; Ohwi and Ohashi). *Food Chemistry* **67**(4), 347-352.
- Saini MK, Keshwa GL (1997) Economics of nitrogen, phosphorus and Mixtalol on rainfed mothbean (*Vigna aconitifolia* (Jacq) Marechal) production. *Agricultural Science Digest (Karnal)* **17**(1), 54-56.
- Saini A, Jawali N (2009) Molecular evolution of 5S rDNA region in *Vigna* subgenus *Ceratotropis* and its phylogenetic implications. *Plant Systematics and Evolution* **280**(3-4), 187-206.
- Saini A, Reddy SK, Jawali N (2008) Intra-individual and intra-species heterogeneity in nuclear rDNA ITS region of *Vigna* species from subgenus *Ceratotropis*. *Genetics Research* **90**(4), 299-316.
- Saini HS (1988) Extractability and evaluation of  $\alpha$ -galactosides of sucrose in leguminous seeds. *Food Chemistry* **28**(2), 149-157.
- Saini ML, Malik JS, Solanki KR, Singh JV (1983) Genotype  $\times$  environmental interaction for green-fodder yield in mothbean. *Indian Journal of Agricultural Sciences* **53**(1), 83-84.
- Saini N, Kiradoo BD, Bhardwaj A, Singh N, Sahani MS (2007) Micronutrient content of certain conventional and unconventional camel feeds of arid western zone of Rajasthan. In '!' pp. 141-143. (College of Veterinary & Animal Science: Rajasthan)
- Saini N, Sharma T, Arya RS (2009) Effect of sulphur supplementation on mineral profile of blood in sheep. *Indian Veterinary Journal* **86**(1), 99-100.
- Saini N, Singh GP (2006) Effect of weaning on growth performance of camel calves. *Indian Journal of Dairy Science* **59**(5), 344-348.
- Sakai A, Honda H, Oshima K, Yamamoto I (1986) Oviposition marking pheromone of two bean weevils, *Callosobruchus chinensis* and *Callosobruchus maculatus*. *Journal of Pesticide Science* **11**(2), 163-168.
- Saker RH, Hoque MI, Islam MM (1992) Studies on the receptive surface of stigma in some *vigna* species. *Bangladesh Journal of Botany* **21**(1), 123-124.
- Sakr MM, Hassan AE, Abderrahman SM (2000) Identification of some *Phaseolus* genotypes by alcohol, glucose-6-phosphate, glutamate and malate dehydrogenase isozymes. *Arab Universities Journal of Agricultural Sciences* **8**(1), 19-30.
- Salgare SA (2007) A fresh light on the embryology of *Phaseolus aconitifolius* Jacquin, obs.: further evidence of a criticism of P. Maheshwari (1950), Souges and Crete's (1952), S.C. Maheshwari (1955), Johri (1963), Rembert (1967a - Ph.d. Thesis, 67b, 69, 71), Bhasin (1971) and Deshpande and Bhasin (1974). *Asian Journal of Bio Science* **2**(1/2), 187-190.
- Salunke BK, Patil KP, Wani MR, Maheshwari VL (2006) Antinutritional constituents of different grain legumes grown in North Maharashtra. *Journal of Food Science and Technology (Mysore)* **43**(5), 519-521.
- Samanta G, Biswas P, Mandal L, Banerjee GC (1984) metabolizable energy value of rice-bean (*Phaseolus-Calcaratus Roxb*) seeds in chicks. *Indian Veterinary Journal* **61**(9), 795-797.

- Sandhu GS, Bassi GS, Sohoo MS (1986b) Evaluation of susceptibility in seeds of ricebean and different genotypes of cowpea to pulse beetle, *Callosobruchus maculatus* (F.). I. Preference for oviposition. *Journal of Research, Punjab Agricultural University* **23**(2), 253-262.
- Sandhu GS, Bassi GS, Sohoo MS (1987b) Evaluation of susceptibility in seeds of ricebean and different genotypes of cowpea to pulse beetle *Callosobruchus maculatus* (F.). II. Growth and development. *Journal of Research, Punjab Agricultural University* **24**(3), 423-432.
- Sangha HS, Mavi GS (1995) Seasonal activity and host-range of blister beetle, *Mylabris phalerata* (Pallas). *Journal of Insect Science* **8**(1), 106-107.
- Sangiri C, Kaga A, Tomooka N, Vaughan D, Srinivas P (2007) [Genetic diversity of the mungbean \(\*Vigna radiata\*, Leguminosae\) gene pool on the basis of microsatellite analysis](#). *Australian Journal of Botany* **55**(8), 837-847.
- Sanjay M, Malhotra SP, Kamal D, Randhir S (1999) Purification and characterization of protease inhibitor from rice bean (*Vigna umbellata* T.) seeds. *Journal of Plant Biochemistry and Biotechnology* **8**(1), 61-64.
- Sankar GRM, Vittal KPR, Chary GR, Ramakrishna YS, Khan IA, Girija A (2006) Statistical assessment of sustainability of genetic resources under different agro-climatic conditions. *Indian Journal of Dryland Agricultural Research and Development* **21**(2), 185-197.
- Sankhla A, Sankhla D, Upadhyaya A, Davies TD, Sankhla N (1990) Influence of lactidichlor-ethyl on growth, chemical composition, and regeneration of moth bean callus cultures. *Comparative Physiology and Ecology* **15**(4), 140-146.
- Sankhla N, Davis TD, Gehlot HS, Upadhyaya A, Sankhla A, Sankhla D (1991) Growth and organogenesis in moth bean callus cultures as influenced by triazole growth regulators and gibberellic acid. *Journal of Plant Growth Regulation* **10**(1), 41-45.
- Sankhla N, Huber W (1975) Effect of salt and abscisic acid on in vivo activity of nitrate reductase in seedlings of *Phaseolus aconitifolius*. *Zeitschrift fur Pflanzenphysiologie* **76**(5), 467-470.
- Sankhla N, Ziegler H, Vyas OP, Stichler W, Trimborn P (1975) Eco-physiological studies on Indian arid zone plants. 5. A screening of some species for the C4-pathway of photosynthetic CO<sub>2</sub>-fixation. *Oecologia* **21**(2), 123-129.
- Sankhla R, Chawan DD (1980) Effect of different seed moisture levels on the germination behaviour of *Phaseolus trilobus* Ait. *Biotogia Plantarum* **22**(5), 388-391.
- Sano-Fujii I (1984) [Effect of bean water content on the production of the active form of \*Callosobruchus maculatus\* \(F.\) \(Coleoptera: Bruchidae\)](#). *Journal of Stored Products Research* **20**(3), 153-161.
- Santosh K, Reager ML, Pareek BL (2006) Yield components of mothbean (*Vigna aconitifolia* (Jacq.) Marechal) as influenced by phosphorus and bio-fertilizers. *Annals of Agricultural Research* **27**(3), 227-229.
- Saono S, Karsono H (1977) Studies on the effect of different rhizobial strains on *Phaseolus calcaratus* Roxb. in sand culture. *Annales Bogorienses* **6**(3), 143-154.
- Saono S, Karsono H (1979) Effect of different rhizobial strains on *Phaseolus calcaratus* Roxb. and *Phaseolus lunatus* grown on various types of soil. *Annales Bogorienses* **7**(1), 21-33.
- Saono S, Karsono H, Abdulkadir S (1975) Notes on the nodulation of some edible legumes of Indonesia. *Annales Bogorienses* **6**(1), 27-41.
- Sapna T, Nag TN (2004) Flavonoids of moth bean cultivars as antimicrobial agent. *Indian Journal of Plant Physiology* **9**(1), 59-61.
- Sapu C, Wungsem R (2009) Study on genetic variability and correlation of morpho-agronomic characters with protein and carbohydrate contents in rice bean. *Agricultural Science Digest* **29**(3), 206-208.
- Saraladevi M, Selvaraju K, Shanmugasundaram P (2008) Efficiency of RAPD and ISSR markers system in assessing genetic variation of rice bean (*Vigna umbellata*) landraces. *EJB, Electronic Journal of Biotechnology* **11**(3), article8.
- Saraswat VN, Rabha MK (1993) Major weeds of rabi and kharif crops and their control. *Indian Farming* **43**(9), 13, 15-23.
- Sarkar A (2006b) Current status and future strategies for improvement of nutritional qualities of rice bean (*Vigna umbellata* (Thunb.) Ohwi and Ohashi). *Journal of Interacademica* **10**(2), 287-292.
- Sarkar A (2006c) Genetic variability and genetic divergence in rice bean: a review. *Environment and Ecology* **24**(1), 76-80.
- Sarkar A, Das B (2007b) Character association and path coefficient analysis in rice bean (*Vigna umbellata* (Thunb.) Ohwi and Ohashi) grown under Terai region of West Bengal. *Journal of Interacademica* **11**(3), 258-264.

- Sarkar A, Mukherjee AK (1991a) Effect of phosphorus on yield and nodulation of greengram (*Phaseolus radiatus*), blackgram (*P. mungo*) and rice bean (*Vigna umbellata*). *Indian Journal of Agricultural Sciences* **61**(5), 328-331.
- Sarkar PK, Debnath NC (1988) Species differences of crops in relation to nutrient composition in alluvial soil. *Environment and Ecology* **6**(2), 348-351.
- Sarkar SK, Ghosh RK, Sounda G, Maitra S, Roy MK, Ghosh KK (1998b) Growth nutrient uptake and water use efficiency of rice bean (*Vigna umbellata*) as affected by different levels of nitrogen, potassium and soil moisture tension. *Journal of Interacademicia* **2**(3), 164-167.
- Sarma BK, Kushwaha RV, Singh M (1991) Screening of rice bean (*Vigna-umbellata*) germplasm tolerant to pod-boring weevil (*Apion-clavipes*). *Indian Journal of Agricultural Sciences* **61**(12), 930-932.
- Sarma BK, Sharma BD, Hazarika GN (1991) Crop genetic resources of North East India: II. Pulses and oilseeds - a review. *Agricultural Reviews* **12**(4), 213-224.
- Sarma BK, Singh M, Pattanayak A (1991) Evaluation of rice bean (*Vigna-umbellata*) germplasm in upland terraces of Meghalaya. *Indian Journal of Agricultural Sciences* **61**(3), 182-184.
- Saroj SD, Sachin H, Shashidhar R, Varsha D, Arun S, Bandekar JR (2007) Radiation processing for elimination of *Salmonella Typhimurium* from inoculated seeds used for sprout making in India and effect of irradiation on germination of seeds. *Journal of Food Protection* **70**(8), 1961-1965.
- Sarjo SD, Shashidhar R, Varsha D, Sachin H, Arun S, Bandekar JR (2006) Microbiological evaluation of sprouts marketed in Mumbai, India, and its suburbs. *Journal of Food Protection* **69**(10), 2515-2518.
- Sarup P, Siddiqui KH, Marhawa KK (1987) Evolution of artificial diets for rearing of the key tissue borers of maize in India. *Journal of Entomological Research* **11**(1), 99-106.
- Sarup P, Siddiqui KH, Marhawa KK (1983) Compounding artificial diets for mass rearing of the maize stalk borer, *Chilo partellus* (Swinhoe). *Journal of Entomological Research* **7**(1), 68-74.
- Sasagawa A, Gomi M, Ohura K, Yamazaki A, Yamada A (2005) Production of Miso based on Koji prepared from mixed different grains using high-pressure treatment. *Nippon Shokuhin Kagaku Kogaku Kaishi = Journal of the Japanese Society for Food Science and Technology* **52**(10), 485-490.
- Sasaki S, Ohnishi M, Mano Y, Kojima M, Ito S, Fujino Y (1989) Characterization of triacylglycerol species in oil-poor beans. *Agricultural and Biological Chemistry* **53**(7), 2021-2023.
- Sastrapradja S, Sutarno H (1977) *Vigna umbellata* (L.) DC. in Indonesia. *Annales Bogorienses* **6**(3), 155-167.
- Satch H (1979) The flowering and ripening of adzuki bean (*Phaseolus angularis* W.F. Wight). *Bulletin of Hokkaido Prefectural Agricultural Experiment Stations*(41), 10-20.
- Sathe SK, Venkatachalam M (2007) Fractionation and biochemical characterization of moth bean (*Vigna aconitifolia* L.) proteins. *LWT - Food Science and Technology* **40**(4), 600-610.
- Satheesha BP, Khalak A, Shivashankar K (1993) Effect of molybdenum seed treatment on yield and economics of rice bean. *Current Research - University of Agricultural Sciences (Bangalore)* **22**(9-10), 125-126.
- Satija CK, Ravi (1996a) Cytomorphological studies in hybrids and amphidiploids of *Vigna radiata* X *V. umbellata*. *Crop Improvement* **23**(1), 19-24.
- Satish L, Gupta GK, Saroj S (1986) Crop disease situation and some new records in Indian arid zone. *Annals of Arid Zone* **25**(4), 311-320.
- Sato H (1979) Introduction of new varieties of summer crops. *Vigna angularis*. *Japanese Journal of Breeding (Ikushugaku Zasshi)* **29**(4), 335-336.
- Sato H (1982) Introduction of new varieties of summer crops. [*Vigna angularis*]. *Japanese Journal of Breeding* **32**(1), 91-92.
- Sato S, Hori Y, Yamate J, Saito T, Kurasaki M, Hatai A (2005) Protective effect of dietary azuki bean (*Vigna angularis*) seed coats against renal interstitial fibrosis of rats induced by cisplatin. *Nutrition* **21**(4), 504-511.
- Sato S, Mukai Y, Yamate J, Kato J, Kurasaki M, Hatai A, Sagai M (2008) Effect of polyphenol-containing azuki bean (*Vigna angularis*) extract on blood pressure elevation and macrophage infiltration in the heart and kidney of spontaneously hypertensive rats. *Clinical and Experimental Pharmacology and Physiology* **35**(1), 43-49.
- Sato S, Yamate J, Hori Y, Hatai A, Nozawa M, Sagai M (2005) Protective effect of polyphenol-containing azuki bean (*Vigna angularis*) seed coats on the renal cortex in streptozotocin-induced diabetic rats. *Journal of Nutritional Biochemistry* **16**(9), 547-553.
- Sato T, Asaka D, Harada T, Matsukawa I (1990) Plant regeneration from adzuki bean epicotyl and callus derived from epicotyl. *Bulletin of Hokkaido Prefectural Agricultural Experiment Stations*(61), 51-60.
- Sato T, Asaka D, Harada T, Matsukawa I (1993) Plant regeneration from protoplasts of adzuki bean (*Vigna angularis* Ohwi & Ohashi). *Japanese Journal of Breeding* **43**(2), 183-191.

- Satoh H, Sawada S, Itoh S (2003) [Rate of acceptance and longevity of pulses and wheat cultivars in Hokkaido](#). *Japanese Journal of Crop Science* **72**(4), 418-423.
- Satwadhar PN, Kadam SS, Salunkhe DK (1981) [Effects of germination and cooking on polyphenols and in vitro protein digestibility of horse gram and moth bean](#). *Qualitas Plantarum -- Plant Foods for Human Nutrition* **31**(1), 71-76.
- Satya V (2004) Screening of select genotypes of mothbean and cowpea for their susceptibility to pulse beetle, *Callosobruchus maculatus* F. (Coleoptera: Bruchidae). *Journal of Arid Legumes* **1**(1), 70-74.
- Satya V (2005) Population built up and quantitative losses due to *Collasobruchus maculatus* F. (Coleoptera: Bruchidae) with special reference to its oviposition behaviour and bio-efficacy of different oils as seed protectants against pulse beetle. **2**(1), 32-36. (Scientific Publishers (India): Jodhpur)
- Satya V (2007) Neem genetic diversity in India and its use as biopesticide and biofertilizer. *Indian Journal of Plant Protection* **35**(2), 185-193.
- Satyan BA, Mahishi DM, Shivashankar G (1982a) Meiosis in the hybrid between greengram and rice bean. *Indian Journal of Genetics and Plant Breeding* **42**(3), 356-359.
- Satyan BA, Reddy PG, Radhamani TR, Prakash KS (1988) Induced genetic variability in amphidiploid generations of mungbean and ricebean crosses. 1988a, . **30**(supplement 1), p. 466
- Satyanarayana J, Singh KM, Singh RN (1995a) Insect pest succession in rice bean. *Bulletin of Entomology (New Delhi)* **36**(1/2), 78-83.
- Satyanarayana J, Singh KM, Singh RN (1995b) Pest complex of rice bean, *Vigna umbellata*. *Bulletin of Entomology (New Delhi)* **36**(1/2), 28-33.
- Satyanarayana J, Singh KM, Singh RN, Subedar S (1995c) Seasonal abundance of rice bean insect pests and natural enemies in relation to abiotic factors. *Bulletin of Entomology (New Delhi)* **36**(1/2), 92-99.
- Satyanarayana J, Singh KM, Singh RN, Subedar S (1997) Influence of intercropping and insecticide. I. Incidence of major pests in rice bean, *Vigna umbellata*. *Indian Journal of Entomology* **59**(3), 316-325.
- Satyanarayana J, Singh KM, Singh RN, Subedar S (1998a) Influence of intercropping and insecticide, II. Natural enemy complex in rice bean, *Vigna umbellata*. *Indian Journal of Entomology* **60**(1), 69-73.
- Satyanarayana J, Singh KM, Singh RN, Subedar S (2001b) Impact of intercropping and insecticide, III yields of rice bean, *Vigna umbellata* (Thunb.) Ohwi and Ohashi. *Indian Journal of Entomology* **63**(2), 114-116.
- Satyavir (1983a) Efficacy of some important insecticides for the control of leafhopper Empoasca kerri Pruthi on mothbean crop. *Indian Journal of Plant Protection* **11**(1/2), 130-133.
- Satyavir (1983b) Efficacy of some important insecticides in the control of Bemisia tabaci (Genn.), a vector of the yellow mosaic disease on mothbean. *Indian Journal of Plant Protection* **11**(1/2), 31-33.
- Savale SR, Sanglikar RV, Sthool VA, Kadam JR (2007) Effect of Leucaena leucocephala barriers and bund on erosion losses and yield of pearl millet+mothbean. *Asian Journal of Soil Science* **2**(2), 142-145.
- Savouré A, Jaoua S, Hua X, Ardiles W, Montagu Mv, Verbruggen N (1995) Isolation, characterization, and chromosomal location of a gene encoding the Δ1-pyrroline-5-carboxylate synthetase in *Arabidopsis thaliana*. *FEBS Letters* **372**(1), 13-19.
- Sawaguchi M (1987) Nutrio-physiological characteristics of Azuki-bean and methods of fertilizer application. *JARQ (Japan Agricultural Research Quarterly)* **21**(2), 77-83.
- Sawaguchi M, Nomura K (1980) Growth and nitrogen uptake by adzuki bean cultivars. *Bulletin of Hokkaido Prefectural Agricultural Experiment Stations*(43), 1-11.
- Sawaguchi M, Nomura K (1981) Changes in carbohydrates of adzuki bean in relation to its growth. *Bulletin of Hokkaido Prefectural Agricultural Experiment Stations*(45), 59-69.
- Sawahel WA, Hassan AH (2002) Generation of transgenic wheat plants producing high levels of the osmoprotectant proline. *Biotechnology Letters* **24**(9), 721-725.
- Sawano M, Shimmen T, Sonobe S (2000) Possible involvement of 65 kDa MAP in elongation growth of azuki bean epicotyls. *Plant and Cell Physiology* **41**(8), 968-976.
- Sawhney RK, Shekhawat NS, Galston AW (1985) Polyamine levels as related to growth, differentiation and senescence in protoplast-derived cultures of *Vigna aconitifolia* and *Avena sativa*. In ' Vol. 3.' pp. 329-337)
- Saxena NB, Mathur SK (1987) Water use of moth bean (*Vigna aconitifolia* L.) in desert. **12**(2), 123-126.
- Saxena NB, Mathur SK (1990) Role of mothbean (*Vigna aconitifolia*) in managing sandy soils. *Transactions of Indian Society of Desert Technology*(15), 49-52.
- Saxena RC (1982) Note on the use of neem kernel for the protection of dew-gram against *Amsacta moorei* Butler. *Indian Journal of Agricultural Sciences* **52**(1), 51-52.
- Saxena RC (1987) Effect of some granular insecticides on germination and growth of the dew-gram *Vigna aconitifolia* (Jacq.) Marechal. *Indian Journal of Entomology* **49**(1), 145-147.

- Saxena SK (1988) 'Lasu' an exclusive top feed for camels during drought. *Indian Farming* **38**(9), 19-20.
- Saxena SN, Kaushik N, Sharma R (2008) Effect of abscisic acid and proline on in vitro flowering in *Vigna aconitifolia*. *Biologia Plantarum* **52**(1), 181-183.
- Saxena SN, Rohit K, Sharma R (2005) Prospects of manipulating in vitro flowering in moth bean for crop improvement. *2*(2), 210-213.
- Saxena SN, Sharma R, Mahla HR (2006) Analysis of induced and naturally available genetic diversity in moth bean (*Vigna aconitifolia* Jacq. Merachal) using RAPD markers. *Journal of Arid Legumes* **3**(1), 60-64.
- Schmidt HE (1992) Bean mosaics. 40-73.
- Schmidt L (1994b) Nutrient content of tropical grain legumes in Madagascar. *Beitrag zum Nährstoffgehalt in Madagaskar angebauter tropischer Körnerleguminosen*. **88**(2), 165-176.
- Seehalak W, Tomooka N, Waranyuwat A, Thipyapong P, Laosuwan P, Kaga A, Vaughan DA (2006) [Genetic diversity of the \*Vigna\* germplasm from Thailand and neighboring regions revealed by AFLP analysis](#). *Genetic Resources and Crop Evolution* **53**(5), 1043-1059.
- Sekar I, Boopathi SNMR, Suresh KK (1998) Fodder production in a hortipastoral system under rainfed condition. *Madras Agricultural Journal* **85**(3/4), 183-185.
- Sekar R, Sulochana CB (1986) [Host range of blackeye cowpea mosaic-virus](#). *Current Science* **55**(23), 1202-1202.
- Sekhar JC, Singh KM, Singh RN, Yeshbir S, Malik KS (1997) Impact of intercropping on the incidence of green jassid, Empoasca kerri on pigeonpea. *Indian Journal of Entomology* **59**(1), 119-123.
- Selvi R, Muthiah AR, Maheswaran M, Shanmugasundaram P (2003) Genetic diversity analysis in the genus *Vigna* based on morphological traits and isozyme markers. *SABRAO Journal of Breeding and Genetics* **35**(2), 103-112.
- Sen R, Sau A (1978) The possible use of bacteriological data in plant systematics. *Science & Culture* **44**(6), 266-268.
- Senecha P, Srivastava HP (1981) Fungi associated with the seeds of moth bean (*Phaseolus aconitifolius* L.). *Geobios, India* **8**(2), 75-76.
- Senecha P, Srivastava HP (1982) Leaf blight of *Vigna aconitifolia* caused by *Macrophomina phaseolina*. *Indian Phytopathology* **35**(1), 160-161.
- Seneviratne GI, Harborne JB (1992b) Constitutive flavonoids and induced isoflavonoids as taxonomic markers in the genus *Vigna*. *Biochemical Systematics and Ecology* **20**(5), 459-467.
- Sequeira RV, Playford CL (2001) [Abundance of \*Helicoverpa\* \(Lepidoptera: Noctuidae\) pupae under cotton and other crops in central Queensland: implications for resistance management](#). *Australian Journal of Entomology* **40**(3), 264-269.
- Seshadri M, Rajekar PK (1981) Comparative study of repetitive DNA in *Phaseolus*. *Indian Journal of Biochemistry & Biophysics* **18**(4), 254-258.
- Setoyama M, Shimada M (2007) Regular oviposition behavior and evolution of decision-making rule in bruchid seed beetle. *Japanese Journal of Ecology* **57**(2), 189-199.
- Setua GC, Das NK, Banerjee ND, Sengupta T, Sudhakar P, Sen S, Saratchandra B (2005) Effect of integrated nutrient management on quality leaf production in mulberry (*Morus alba*) under rainfed, alluvial soil conditions. *Indian Journal of Agricultural Sciences* **75**(8), 474-478.
- Setua GC, Setua M, Banerjee ND, Dutta AK, Debnath S, Ghosh JK, Urs SR (2006b) Integrated weed management for-sustainable leaf production in mulberry (*Morus alba*) under irrigated condition. *Indian Journal of Agricultural Sciences* **76**(4), 258-261.
- Shah RM (1981) Pulses forge ahead in Gujarat. *Pulse Crops Newsletter* **1**(1), 12-13.
- Shah SB (1975) [Chemical examination of the fats from the seeds of \*Phaseolus\* species](#). *Current Science* **44**(1), 11-12.
- Shahnaz D, Sadia A, Marium T, Zaki MJ (2007) Use of sea weed and bacteria in the control of root rot of mash bean and sunflower. *Pakistan Journal of Botany* **39**(4), 1359-1366.
- Shanker RU, Ganeshiah KN (1980a) [Evolutionary significance of pollen to ovule ratio - a study in some pulse crops](#). *Current Science* **49**(6), 244-245.
- Sharma BD, Hore DK (1994) Association of yield attributes in ricebean (*Vigna umbellata*). *Legume Research* **17**(2), 135-137.
- Sharma BD, Hore DK, Dobhal VK (1996) Classificatory analysis in ricebean (*Vigna umbellata*). *Journal of Hill Research* **9**(1), 79-81.
- Sharma BL, Sharma PK, Fageria VD (1993) Suitability of rainy-season crops for late-sown rainfed conditions. *Indian Journal of Agronomy* **38**(1), 22-24.

- Sharma BM (2003) Productivity of grain legumes in agri-silviculture systems under hot arid conditions. In ' ' pp. 279-284. (Scientific Publishers (India): Jodhpur)
- Sharma HC, Franzmann BA (2000) Biology of the legume pod borer, *Maruca vitrata* (Fabricius) and its damage to pigeonpea and Adzuki bean. *Insect Science and its Application* **20**(2), 99-108.
- Sharma JP, Gupta JS (1982) A new leaf spot disease of moth bean. *Indian Phytopathology* **35**(1), 160.
- Sharma JP, Gupta JS (1984) Fungal deterioration of moth bean (*Phaseolus aconitifolius* Jacq.) seeds during storage. *International Journal of Tropical Plant Diseases* **2**(2), 161-167.
- Sharma JP, Gupta JS (1986) Studies on the control of fungal spoilage of moth bean seeds during storage by fungicides. *Journal of the Indian Botanical Society* **65**(4), 456-458.
- Sharma KD, Sunil K, Singh N (2007) Dry matter accumulation and its partitioning in different moth bean [*Vigna aconitifolia* (Jacq.) Marechal] genotypes under moisture stress. *Annals of Biology* **23**(1), 91-95.
- Sharma LA, Singh MRK, Singh NB (1998) Heterosis and combining ability for grain yield and its components in rice bean (*Vigna umbellata* (Thunb) Ohwi and Ohashi). *Indian Journal of Hill Farming* **11**(1/2), 27-33.
- Sharma LC, Reddy BN, Saxena S (1981) Rhizobium inoculants for pulses in Rajasthan. *Pulse Crops Newsletter* **1**(1), 106.
- Sharma PK, Tiwana MS, Gupta BK (1993) Associations of summer legume and non-legume fodders under rainfed conditions. *Agricultural Science Digest (Karnal)* **13**(1), 20-22.
- Sharma R, Mahla HR, Mohapatra T, Bhargava SC, Sharma MM (2003) Isolating plant genomic DNA without liquid nitrogen. *Plant Molecular Biology Reporter* **21**(1), 43-50.
- Sharma RB, Roy AN (1983) The impact of *Citrus sinensis* rind as urban waste components on the field performance of pulses. *Seeds & Farms* **9**(2/3), 23-25.
- Sharma RC (1993) Stability in mothbean for seed yield and associated traits under rainfed conditions. *Indian Journal of Pulses Research* **6**(1), 21-24.
- Sharma RC (1998) Stability in mothbean (*Vigna aconitifolia*). *Indian Journal of Agricultural Sciences* **68**(2), 130-131.
- Sharma RC, Joshi P, Kakani RK (2003) Mutation breeding in moth bean. In ' ' pp. 88-92. (Scientific Publishers (India): Jodhpur)
- Sharma RC, Kakani RK, Solanki SS, Garg DK (2003) Genetic divergence in moth bean. In ' ' pp. 104-106. (Scientific Publishers (India): Jodhpur)
- Sharma RC, Mathur JR (1982) Studies of flower shedding behaviour in mothbean [*Vigna aconitifolia* (Jacq.) Marechal]. *Science and Culture* **48**(2), 64-65.
- Sharma RC, Mathur JR (1983) Jadia - a promising cultivar of mothbean. *Indian Farming* **32**(11), 13-15,26.
- Sharma RC, Mathur JR (1988) Cultivation practices for mothbean in Rajasthan. *Indian Farming* **38**(2), 29-31.
- Sharma RC, Singh RC, Mathur JR (1980) Improvement of moth bean - an important crop of arid and semi-arid zones of Rajasthan and Gujarat. *Annals of Arid Zone* **19**(4), 457-459.
- Sharma RN, Pandey RL, Chitale MW (1998) Genetic variability and character association for yield components in ricebean germplasm. *Indian Journal of Pulses Research* **11**(2), 127-128.
- Sharma RP, Raman KR, Sharma MS, Poddar BK (2008) Effect of cereals and legumes intercropping on production potential, economics and quality of fodder during summer season. *Range Management and Agroforestry* **29**(2), 129-133.
- Sharma RP, Raman KR, Singh AK (2009a) Fodder productivity and economics of pearl millet (*Pennisetum typhoides*) with legumes intercropping under various row proportions. *Indian Journal of Agronomy* **54**(3), 301-305.
- Sharma RP, Raman KR, Singh AK, Poddar BK, Rajesh K (2009b) Production potential and economics of multi-cut forage sorghum (*Sorghum sudanense*) with legumes intercropping under various row proportions. *Range Management and Agroforestry* **30**(1), 67-71.
- Sharma RP, Singh AK, Poddar BK, Raman KR (2008) Forage production potential and economics of maize (*Zea mays*) with legumes intercropping under various row proportions. *Indian Journal of Agronomy* **53**(2), 121-124.
- Sharma VK, Jai D, Gupta VP (1995) Genetic variability for protein and its association with seed yield traits in adzuki bean (*Vigna angularis*) germplasm. *Legume Research* **18**(1), 35-40.
- Sharma VK, Sharma JD, Gupta VP (1995) Genotypic variability and genotype × environment interactions for seed-yield characters in adzuki bean (*Vigna angularis*). *Indian Journal of Agricultural Sciences* **65**(7), 490-493.
- She Y, Rong T, Su S, Liu J, Tian M (2006) Genetic diversity of the cultivated adzuki bean (*Vigna angularis*) germplasm resources analysed with RAMP molecular markers. *Acta Agronomica Sinica* **32**(2), 217-222.

- Shekhawat BS (1992) Effect of sowing dates and row spacings on moth bean varieties. *Indian Journal of Pulses Research* **5**(2), 160-163.
- Shekhawat BS, Shaktawat RPS, Chauhan RPS (2003a) Effect of thiourea application on productivity of mothbean (*Vigna aconitifolia*). *Indian Journal of Agricultural Sciences* **73**(5), 294-295.
- Shekhawat BS, Shaktawat RPS, Shekhawat PS (2003b) Effect of fluchloralin on mothbean [*Vigna aconitifolia* (Jacq) Marechal] and associate weeds. *Indian Journal of Weed Science* **35**(1/2), 141-142.
- Shekhawat BS, Sharma RC, Mathur JR, Agnihotri JP (1978) Evaluation of moth (*Vigna aconitifolius*) germplasm against moth yellow mosaic virus. **8**(1), 50.
- Shekhawat GS, Chundawat GS, Singh S (1972) Response of moth (*Phaseolus aconitifolius*) to nitrogen and phosphorus under rainfed conditions of arid Rajasthan. *Indian Journal of Agronomy* **17**(2), 65-68.
- Shekhawat NS, Galston AW (1983a) Isolation, culture, and regeneration of moth bean *Vigna aconitifolia* leaf protoplasts. *Plant Science Letters* **32**(1/2), 43-51.
- Shekhawat NS, Galston AW (1983b) Regeneration of food legume leaf protoplasts. **72**(1, Supplement), 143.
- Shet MS, Murugiswamy B, Madaiah M (1986) Lipid profile and fatty acid composition of moth bean (*Vigna aconitifolia*) seeds. *Fette Seifen Anstrichmittel* **88**(7), 264-266.
- Shiano T, Osaki M, Komatsu K, Tadano T (1995) Comparison of production efficiency of harvesting organs among field crops. II. CO<sub>2</sub> assimilation and reassimilation in the harvesting organs. *Soil Science and Plant Nutrition* **41**(1), 21-31.
- Shiau Y, Lai L, Horng S (1994) Effects of oviposition behavior on host preference of *Callosobruchus maculatus*. *Chinese Journal of Entomology* **14**(2), 245-253.
- Shibaoka H (1994) Plant hormone-induced changes in the orientation of cortical microtubules: alterations in the cross-linking between microtubules and the plasma membrane. *Annual Review of Plant Physiology and Plant Molecular Biology* **45**, 527-544.
- Shigematsu H, Toko K, Matsuno T, Yamafuji K (1994) Early gravi-electrical responses in bean epicotyls. *Plant Physiology* **105**(3), 875-880.
- Shimada H, Murata K, Fujita S, Chiba I, Hara M, Shirai S, Adachi T (1997) A new adzuki bean variety "Hokuto-dainagon". *Bulletin of Hokkaido Prefectural Agricultural Experiment Stations*(72), 85-95.
- Shimada M (1985) Niche modification and stability of competitive systems. II. Persistence of interspecific competitive systems with parasitoid wasps. *Researches on Population Ecology* **27**(2), 203-216.
- Shimada M (1989) Systems analysis of density-dependent population processes in the azuki bean weevil, *Callosobruchus chinensis*. *Ecological Research* **4**(2), 145-156.
- Shimada M (1990) Preadaptation of a wild bean weevil, *Kytorhinus sharpianus* Bridwell (Coleoptera: Bruchidae), for infesting the storage green gram, *Vigna radiata* Wilczek (Leguminosae). *Applied Entomology and Zoology* **25**(1), 132-134.
- Shimada M (1999) Population fluctuation and persistence of one-host-two-parasitoid systems depending on resource distribution: from parasitizing behavior to population dynamics. *Researches on Population Ecology* **41**(1), 69-79.
- Shimada M, Tuda M (1996) Delayed density dependence and oscillatory population dynamics in overlapping-generation systems of a seed beetle *Callosobruchus chinensis*: matrix population model. *Oecologia* **105**(1), 116-125.
- Shimamura S, Mochizuki T, Inouye J (1997a) [Formation of the lysigenous intercellular space in adventitious roots of leguminous species](#). *Japanese Journal of Crop Science* **66**(2), 208-213.
- Shimizu K, Mitsui Y (1987) Varietal difference of azuki bean and kidney bean in susceptibility to the races of the soybean cyst nematode, *Heterodera glycines* Ichinohe, 1952. *Annual Report of the Society of Plant Protection of North Japan*(38), 109-110.
- Shimizu M, Mitsunaga T (1992) Thiamin and thiamin-binding activity in legume seeds. *Memoirs of the Faculty of Agriculture of Kinki University*(25), 55-59.
- Shinano T, Osaki M, Komatsu K, Tadano T (1993a) Comparison of production efficiency of the harvesting organs among field crops: I. Growth efficiency of the harvesting organs. *Soil Science and Plant Nutrition* **39**(2), 269-280.
- Shinano T, Osaki M, Tadano T (1993b) Comparison of production efficiency among field crops related to nitrogen nutrition and application. 227-230.
- Shinano T, Osaki M, Yamada S, Tadano T (1994) Comparison of root growth and nitrogen absorbing ability between Gramineae and Leguminosae during the vegetative stage. *Soil Science and Plant Nutrition* **40**(3), 485-495.

- Shinde AK, Jamadagni BM, Mhatre VV (2001) Water relations of five grain legumes influenced by potassium and growth regulators under water stress. *Journal of Potassium Research* **17**(1/4), 107-111.
- Shinde VK, Thakare CS (2000) Studies on the host range of Phakopsora pachyrhizi Syd. *Journal of Maharashtra Agricultural Universities* **25**(1), 94-95.
- Shinoda K, Yoshida T (1984) Relationship between adult feeding and emigration from beans of azuki bean weevil, *Callosobruchus chinensis* Linné (Coleoptera: Bruchidae). *Applied Entomology and Zoology* **19**(2), 202-211.
- Shinoda K, Yoshida T (1987) Effect of fungal feeding on longevity and fecundity of the azuki bean weevil, *Callosobruchus chinensis* (L.) (Coleoptera: Bruchidae), in the azuki bean field. *Applied Entomology and Zoology* **22**(4), 465-473.
- Shinoda K, Yoshida T, Igarashi H (1992) Population ecology of the azuki bean beetle, *Callosobruchus chinensis* (L.), (Coleoptera: Bruchidae) on two wild leguminous hosts, *Vigna angularis* var. *nipponensis* and *Dunbaria villosa*. *Applied Entomology and Zoology* **27**(3), 311-318.
- Shinoda K, Yoshida T, Okamoto T (1991) Two wild leguminous host plants of the azuki bean weevil, *Callosobruchus chinensis* (L.) (Coleoptera: Bruchidae). *Applied Entomology and Zoology* **26**(1), 91-98.
- Shinoda WK, Yoshida T (1985) Field biology of the azuki bean weevil, *Callosobruchus chinensis* (L.) (Coleoptera: Bruchidae). I. Seasonal prevalence and assessment of field infestation of Aki-Azuki, an autumn variety of *Phaseolus angularis*. *Japanese Journal of Applied Entomology and Zoology* **29**(1), 14-20.
- Shiozaki H, Tsuchiya K (1985) Differences of response to cool summer damage and wet damage among upland crops. *JARQ (Japan Agricultural Research Quarterly)* **18**(4), 268-274.
- Shiraishi K (2005) Dietary intakes of eighteen elements and K-40 in eighteen food categories by Japanese subjects. *Journal of Radioanalytical and Nuclear Chemistry* **266**(1), 61-69.
- Shirakawa N, Tomioka M, Iwane Y, Takeuchi M, Senda S, Hirota K (1979) Studies on the uracil herbicides 1. Herbicidal properties of 5-bromo-3-butyl-6-methyluracil. *Weed Research, Japan* **24**(3), 199-207.
- Shivadhar S, Singh G, Shyam S (1989) A simple method for measuring leaf area in different crop plants. *Journal of the Andaman Science Association* **5**(2), 146-148.
- Shivani C, Aarti S (2004) Effect of soaking and sprouting on tannin, phytate and in vitro iron in underutilized legumes - horse gram (*Dolichos biflorus*) and moth bean (*Phaseolus aconitifolius*). *Journal of Food Science and Technology (Mysore)* **41**(5), 547-550.
- Shrotriya AK, Verma LS, Ramvir S, Chaudhary DR (1992) Effect of moisture variation on seed physiology. *Indian Journal of Agricultural Sciences* **62**(6), 373-375.
- Shukla GP, Melkania NP, Singh K, Rajpali SK, Arya ON (2003) Genotype x location environment interaction in ricebean [*Vigna umbellata* (Thunb.) Ohwi and Ohashi] with special reference to acidic soil. *Forage Research* **28**(4), 194-200.
- Siddhuraju P, Bravo L, Saura-Calixto F (1998) Chemical composition and carbohydrate digestibility of common and underexploited legumes. 380-381.
- Siddhuraju P, Vijayakumari K, Janardhanan K (1993) Genetic resources of tribal pulses. *Plant Genetic Resources Newsletter*(96), 47-49.
- Siddhuraju P, Vijayakumari K, Janardhanan K (1994) Chemical analysis and nutritional assessment of the less known pulses, *Vigna aconitifolia* (Jacq.) Marechal and *Vigna vexillata* (L.) A. Rich. *Plant Foods for Human Nutrition* **45**(2), 103-111.
- Siddhuraju P, Vijayakumari K, Janardhanan K (1997) Tryptophan contents of total (true) proteins and protein fractions, albumins and globulins of tribal/little known/under-exploited Indian legumes. *Journal of Food Science and Technology (Mysore)* **34**(2), 140-142.
- Siddiqui JI (1981) Record of Apanteles obliquea Wlk. and Meteorus sp. (Hymenoptera: Braconidae) as parasites of the hairy caterpillar *Amsacta moorei* Butl. (Lepidoptera: Arctiidae). *Journal of the Bombay Natural History Society* **78**(1), 181-182.
- Siddiqui KH, Sarup P (1978) Differential response of *Chilo partellus* (Swinhoe) reared on various artificial diets and host plant maize. *Journal of Entomological Research* **2**(2), 123-128.
- Siddiqui KH, Sarup P (1980) Impact of vitamin E on egg production of the maize stalk borer, *Chilo partellus* (Swinhoe) reared in artificial diets. *Journal of Entomological Research* **4**(1), 45-50.
- Siddiqui KH, Sarup P (1981) Significance of ascorbic acid in artificial diets for mass rearing the maize stalk borer, *Chilo partellus* (Swinhoe). *Journal of Entomological Research* **5**(1), 91-93.
- Siddiqui KH, Sarup P (1982) Compounding of artificial diets with formaldehyde affect rearing of the maize stalk borer, *Chilo partellus* (Swinhoe). *Journal of Entomological Research* **6**(2), 197-201.
- Siddiqui KH, Sarup P, Marwaha KK (1985) Effect of streptomycin in artificial diet on mass rearing of the maize stalk borer, *Chilo partellus* (Swinhoe). *Journal of Entomological Research* **9**(1), 60-64.

- Siddiqui KH, Sarup P, Panwar VPS, Marwaha KK (1977) Evolution of base-ingredients to formulate artificial diets for the mass rearing of *Chilo partellus* (Swinhoe). *Journal of Entomological Research* **1**(2), 117-131.
- Siddiqui MA, Ehteshamul-Haque S, Ghaffar A (1999) Use of *Pseudomonas aeruginosa* and fungal antagonists in the control of root knot-root rot disease complex on mungbean and mashbean. *Pakistan Journal of Nematology* **17**(2), 155-167.
- Sidhu MC (2003) Cytogenetic and isozyme studies in interspecific hybrids of *Vigna radiata* and *V. mungo* with *V. trilobata*. *Crop Improvement* **30**(2), 140-145.
- Sidhu N, Satija CK (2003) Cytomorphological characterization of amphidiploids of *Vigna radiata* x *V. umbellata*. *Crop Improvement* **30**(1), 25-32.
- Sihag SK, Khatri RS, Joshi UN (2004) Genetic variability and heritability for grain yield and its components in mothbean [*Vigna aconitifolia* (Jacq.) Marechal]. *Annals of Biology* **20**(2), 219-222.
- Sihag SK, Khatri RS, Singh JV (2005) Correlation and path analysis for grain yield in mothbean [*Vigna aconitifolia* (Jacq.) Marechal]. *2*(2), 306-308.
- Sikkema RH, Soltani N, Shropshire C, Robinson DE (2006) Response of adzuki bean to pre-emergence herbicides. *Canadian Journal of Plant Science* **86**(2), 601-604.
- Sikora RA, Greco N, Silva JFV (2005) Nematode parasites of food legumes. In '' pp. 259-318. (CABI Publishing: Wallingford)
- Simon MV, Benko-Iseppon AM, Resende LV, Winter P, Kahl G (2007) [Genetic diversity and phylogenetic relationships in \*Vigna\* Savi germplasm revealed by DNA amplification fingerprinting](#). *Genome* **50**, 538-547.
- Singh AV, Nath LK (2009a) Evaluation of binder property of Moth bean starch in compressed solid dosage form. *International Journal of PharmTech Research* **1**(2), 365-368.
- Singh AV, Nath LK (2009b) Evaluation of physicochemical character and pasting behaviour of *Phaseolus aconitifolius* Jacq. starch. *Electronic Journal of Environmental, Agricultural and Food Chemistry* **8**(10), 984-990.
- Singh AV, Nath LK, Guha M (2009) Evaluation of physicochemical character and pasting behaviour of *Phaseolus aconitifolius* Jacq. starch. *Asian Journal of Experimental Sciences* **23**(3), 589-595.
- Singh B, Verma MM (1988) RBL-1 a pulse variety of rice bean *Vigna-umbellata* Thunb. Ohwi and Ohashi. *Journal of Research Punjab Agricultural University* **25**(3), 507.
- Singh BM, Hasan U (1980) Effect of temperature and concentrations of 'ABA' on seed germination of *Vigna aconitifolia* (Jacq.) Marechal. *Tropical Grain Legume Bulletin*(17/18), 43-45.
- Singh BM, Shyam KR, Sud VK (1979) Web blight of *Vigna angularis* (Wight) Wilczeck caused by *Rhizoctonia solani* Kuhn. *Tropical Grain Legume Bulletin*(16), 14-15.
- Singh DP (1990a) Distant hybridization in genus *Vigna* - a review. *Indian Journal of Genetics & Plant Breeding* **50**(3), 268-276.
- Singh G, Singh M (1996) Combining ability for yield and its components in rice bean. *Indian Journal of Genetics and Plant Breeding* **56**(4), 520-525.
- Singh G, Singh M, Dhiman KR (1992) Correlation and path analysis in ricebean under mid altitude conditions. *Crop Improvement* **19**(2), 152-154.
- Singh G, Singh SP, Chaudhary BS (1998) Genetic divergence analysis in ricebean. *Indian Journal of Genetics and Plant Breeding* **58**(1), 101-105.
- Singh H, Kushwaha HL, Mishra D (2007) Development of seed drill for sowing on furrow slants to increase the productivity and sustainability of arid crops. *Biosystems Engineering* **98**(2), 176-184.
- Singh H, Mishra D, Nahar NM (2004) Energy use pattern in production agriculture of a typical village in arid zone - Part III. *Energy Conversion and Management* **45**(15/16), 2453-2472.
- Singh HB, Joshi BS, Chandel KPS, Pant KC, Saxena RK (1974) Genetic diversity in some Asiatic *Phaseolus* species and its conservation. *Indian Journal of Genetics and Plant Breeding* **34A**, 52-57.
- Singh KA, Singh LN (1990a) Intercropping of some temperate grasses at mid-hills of Sikkim. *Indian Journal of Agricultural Sciences* **60**(7), 466-470.
- Singh KC, Singh RP (1977) Intercropping of annual grain legumes with sunflower. *Indian Journal of Agricultural Sciences* **47**(11), 563-567.
- Singh KD, Sreenivasulu R (1990) Studies on flue cured tobacco based cropping system in rainfed alfisols of southern Andhra Pradesh. *Annals of Agricultural Research* **11**(1), 80-88.
- Singh KP, Ashwani K, Saharan RP, Ram K (2006) A new boldseeded genotype of mungbean-MRH-5. *National Journal of Plant Improvement* **8**(1), 92-93.

- Singh KP, Monika, Sareen PK, Ashwani K (2003b) Interspecific hybridization studies in *Vigna radiata* (L.) Wilczek and *Vigna umbellata* L. *National Journal of Plant Improvement* **5**(1), 16-18.
- Singh KP, Shrikant (2005) Stability analysis for advance generations of the interspecific hybrid *Vigna radiata* x *Vigna umbellata*. *National Journal of Plant Improvement* **7**(1), 57-58.
- Singh M, Sarma BK (1993) NATURAL OUTCROSSING IN RICE BEAN (VIGNA-UMBELLATA). *Indian Journal of Agricultural Sciences* **63**(8), 520-521.
- Singh MP (1985) Studies on the damage and chemical control of the weevil *Cyrtozemias dispar* Pasc on some legume crops. *Entomon* **10**(4), 261-266.
- Singh MRK, Chakravarti D, Singh NB (1997) Genetic variability, correlation and path analysis in rice bean (*Vigna umbellata* (Thunb) Ohwi and Ohashi) cultivars of Manipur. *Indian Journal of Hill Farming* **10**(1/2), 23-28.
- Singh MRK, Sharma PR, Singh LI (1999) Genetic divergence among rice bean (*Vigna umbellata* (Thunb) Ohwi and Ohashi) cultivars of Manipur hills. *Indian Journal of Genetics and Plant Breeding* **59**(2), 221-225.
- Singh NB, Devi TR (2006a) [Biological effects of gamma rays on M1 generation of ricebean {Vigna umbellata \(Thunb.\) Ohwi and Ohashi}](#). *Legume Research* **29**(4), 266-269.
- Singh PK, Ojah H, Singh JK, Alpana D, Pattanayak A, Verma MR (2006b) Genetic variability and characters association in ricebean (*Vigna umbelata*) genotypes of north east India. *Environment and Ecology* **24S**(Special 3A), 923-926.
- Singh R (1991) Toxicity of endosulfan to *Chilo partellus* (Swinhoe) reared on artificial diets formulated with different quantities of some important dietary ingredients. *Journal of Entomological Research* **15**(3), 169-178.
- Singh R (1992) Differential response of maize stalk borer, *Chilo partellus* (Swinhoe) larvae reared on artificial diets as exemplified by intensity of damage of maize plant. *Journal of Entomological Research* **16**(4), 287-292.
- Singh R, Sarup P (1991) Effect of ascorbic acid in artificial diet (based on green gram + dew gram) on the susceptibility of larvae and adults of *Chilo partellus* (Swinhoe) to endosulfan. *Journal of Entomological Research* **15**(2), 125-134.
- Singh RB, Saha RC, Singh S (2000) Effect of feeding ricebean and sorghum sudan mixed fodder on growth and nutrient utilization in crossbred calves. *Indian Journal of Animal Nutrition* **17**(2), 160-161.
- Singh RC, Faroda AS, Paroda RS, Dahiya DR (1981) Choose your pulse for summer. *Pulse Crops Newsletter* **1**(1), 79-80.
- Singh RJ, Jauhar PP (2005) 'Genetic resources, chromosome engineering, and crop improvement. Vol 1: grain legumes.' (Taylor & Francis: Boca Raton) ix + 376 pp.
- Singh RN (2005) Yield gap in arid land crops of thar desert of Rajasthan. *2*(2), 361-364.
- Singh RP (1980) Cropping systems for drylands of the Indian arid zone. *Annals of Arid Zone* **19**(4), 437-443.
- Singh RP, Singh KC (1980) Relative performance of rainfed moth varieties grown under different fertility levels. *Forage Research* **6**(2), 161-164.
- Singh RP, Singh KC (1981) Relative performance of moth varieties grown during summer season with different fertility levels. *Forage Research* **7**(1), 43-48.
- Singh RV, Arya MPS (1994) Intercropping studies on barnyard millet under rainfed conditions. *Agricultural Science Digest (Karnal)* **14**(3/4), 213-217.
- Singh RV, Arya MPS (1995a) Effect of seed ratio on performance of fingermillet (*Eleusine coracana*)-based mixed-cropping systems. *Indian Journal of Agronomy* **40**(3), 366-370.
- Singh RV, Arya MPS (1995b) Nitrogen requirement of barnyard millet (*Echinochloa frumentacea*) + pulse intercropping systems. *Indian Journal of Agronomy* **40**(2), 172-175.
- Singh RV, Arya MPS (1999a) Effect of seed ratio on barnyard millet (*Echinochloa frumentacea*)-based mixed cropping system. *Indian Journal of Agronomy* **44**(1), 51-55.
- Singh RV, Arya MPS (1999b) Nitrogen requirement of fingermillet (*Eleusine coracana*) plus pulse intercropping system. *Indian Journal of Agronomy* **44**(1), 47-50.
- Singh S, Ram T (1988) Growth and growth efficiency of seedlings in different crops. *Indian Journal of Plant Physiology* **31**(1), 117-119.
- Singh SA (1984) Two fungi attacking pulse crops. *FAO Plant Protection Bulletin* **32**(3), 112.
- Singh SK, Srivastava HP (1984a) A new leaf spot disease of moth bean caused by *Alternaria alternata*. *National Academy of Sciences, India, Science Letters* **7**(8), 243.
- Singh SK, Srivastava HP (1984b) Macrophomina phaseolina diseases of moth bean, *Vigna aconitifolia* (Jacq.) Morechal. *National Academy of Sciences, India, Science Letters* **7**(7), 211-212.
- Singh SK, Srivastava HP (1988) Symptoms of Macrophomina phaseolina infections on moth bean seedlings. *Annals of Arid Zone* **27**(2), 151-152.

- Singh SK, Srivastava HP (1989) Some new fungal diseases of moth bean. *Indian Phytopathology* **42**(1), 164-167.
- Singh SP, Ahuja KN (1990) Intercropping of grain sorghum with fodder legumes under dryland conditions in north-western India. *Indian Journal of Agronomy* **35**(3), 287-296.
- Singh SP, Misra BK, Chandel KPS, Pant KC (1980) Major food constituents of rice-bean (*Vigna-umbellata*). *Journal of Food Science and Technology-Mysore* **17**(5), 238-240.
- Singh SP, Misra BK, Sikka KC, Chandel KPS, Pant KC (1985) Studies on some nutritional aspects of rice bean (*Vigna-umbellata*). *Journal of Food Science and Technology-Mysore* **22**(3), 180-185.
- Singh VP, Tomer YS (1989) A photoinsensitive promising mutant of ricebean (*Vigna umbellata*). *Legume Research* **12**(1), 47-48.
- Sinha R, Kawatra A, Sehgal S (2005) Saponin content and trypsin inhibitor activity of cowpea: Varietal differences and effects of processing and cooking methods. *Journal of Food Science and Technology-Mysore* **42**(2), 182-185.
- Sinha R, Singh KD, Singh AN, Upadhyay PS (1975) Effect of different crop rotations on the yield and quality of chewing tobacco. *Indian Journal of Agricultural Sciences* **45**(3), 128-131.
- Sinsinwar BS (1996) Production potential of forages in rice (*Oryza sativa*)-based cropping system. *Indian Journal of Agronomy* **41**(2), 188-190.
- Siripornadulsil S, Traina S, Verma DPS, Sayre RT (2002) Molecular mechanisms of proline-mediated tolerance to toxic heavy metals in transgenic microalgae. *Plant Cell* **14**(11), 2837-2847.
- Siriwardhane D, Egawa Y, Tomooka N (1991a) Cross-compatibility of cultivated adzuki bean (*Vigna angularis*) and rice bean (*V. umbellata*) with their wild relatives. *Plant Breeding* **107**(4), 320-325.
- Sivalingam PN, Padmalatha KV, Mandal B, Monga D, Ajmera BD, Malathi VG (2007) Detection of begomoviruses by PCR in weeds and crop plants in and around cotton field infected with cotton leaf curl disease. *Indian Phytopathology* **60**(3), 356-361.
- Smartt J (1981) Gene pools in Phaseolus and Vigna cultigenes. *Euphytica* **30**(2), 445-449.
- Smartt J (1985b) Evolution of grain legumes. III. Pulses in the genus Vigna. *Experimental Agriculture* **21**(2), 87-100.
- Smil V (1997) Some unorthodox perspectives on agricultural biodiversity. The case of legume cultivation. *Agriculture Ecosystems & Environment* **62**(2-3), 135-144.
- Soga K, Kotake T, Wakabayashi K, Kamisaka S, Hoson T (2008) Transient increase in the transcript levels of  $\gamma$ -tubulin complex genes during reorientation of cortical microtubules by gravity in azuki bean (*Vigna angularis*) epicotyls. *Journal of Plant Research* **121**(5), 493-498.
- Soga K, Wakabayashi K, Hoson T, Kamisaka S (1999) Hypergravity increases the molecular mass of xyloglucans by decreasing xyloglucan-degrading activity in azuki bean epicotyls. *Plant and Cell Physiology* **40**(6), 581-585.
- Soga K, Wakabayashi K, Hoson T, Kamisaka S (2000) Changes in the apoplastic pH are involved in regulation of xyloglucan breakdown of azuki bean epicotyls under hypergravity conditions. *Plant and Cell Physiology* **41**(4), 509-514.
- Soga K, Wakabayashi K, Kamisaka S, Hoson T (2004) Graviperception in growth inhibition of plant shoots under hypergravity conditions produced by centrifugation is independent of that in gravitropism and may involve mechanoreceptors. *Planta* **218**(6), 1054-1061.
- Soga K, Wakabayashi K, Kamisaka S, Hoson T (2005) Mechanoreceptors rather than sedimentable amyloplasts perceive the gravity signal in hypergravity-induced inhibition of root growth in azuki bean. *Functional Plant Biology* **32**(2), 175-179.
- Soga K, Wakabayashi K, Kamisaka S, Hoson T (2006) Hypergravity induces reorientation of cortical microtubules and modifies growth anisotropy in azuki bean epicotyls. *Planta* **224**(6), 1485-1494.
- Soga K, Wakabayashi K, Kamisaka S, Hoson T (2007) Effects of hypergravity on expression of XTH genes in adzuki bean epicotyls. *Physiologia Plantarum* **131**(2), 332-340.
- Solaiappan U, Krishnan SM, Veerabadran V (1996) Effect of rainfed green-manure crops on succeeding rice (*Oryza sativa*). *Indian Journal of Agronomy* **41**(1), 147-149.
- Solanki SD, Chauhan RM, Tikka SBS (2003) Character associations over environments in moth bean. In '!' pp. 112-114. (Scientific Publishers (India): Jodhpur)
- Solanki SS, Sharma RC, Kakani RK, Garg DK (2003) Estimation of genetic parameters in moth bean. In '!' pp. 84-87. (Scientific Publishers (India): Jodhpur)
- Soltani N, Robinson DE, Shropshire C, Sikkema PH (2006) Adzuki bean (*Vigna angularis*) responses to post-emergence herbicides. *Crop Protection* **25**(6), 613-617.

- Soltani N, Shropshire C, Robinson DE, Sikkema PH (2005) Sensitivity of adzuki bean (*Vigna angularis*) to preplant-incorporated herbicides. *Weed Technology* **19**(4), 897-901.
- Somta C, Somta P, Tomooka N, Ooi PAC, Vaughan DA, Srinives P (2008) Characterization of new sources of mungbean (*Vigna radiata* (L.) Wilczek) resistance to bruchids, *Callosobruchus* spp. (Coleoptera : Bruchidae). *Journal of Stored Products Research* **44**(4), 316-321.
- Somta P, Kaga A, Tomooka N, Isemura T, Vaughan DA, Srinives P (2008) Mapping of quantitative trait loci for a new source of resistance to bruchids in the wild species *Vigna nepalensis* Tateishi & Maxted (*Vigna* subgenus *Ceratotropis*). *TAG Theoretical and Applied Genetics* **117**(4), 621-628.
- Somta P, Kaga A, Tomooka N, Kashiwaba K, Isemura T, Chaitieng B, Srinives P, Vaughan DA (2006) Development of an interspecific *Vigna* linkage map between *Vigna umbellata* (Thunb.) Ohwi and Ohashi and *V. nakashime* (Ohwi) Ohwi & Ohashi and its use in analysis of bruchid resistance and comparative genomics. *Plant Breeding* **125**(1), 77-84.
- Somta P, Talekar NS, Srinives P (2006a) Characterization of *Callosobruchus chinensis* (L.) resistance in *Vigna umbellata* (Thunb.) Ohwi & Ohashi. *Journal of Stored Products Research* **42**(3), 313-327.
- Sonda M, Toko K, Hayashi K, Fujiyoshi T, Ezaki S, Yamafuji K (1990) Relationship between growth and electric oscillations in bean roots. *Plant Physiology* **93**(2), 532-536.
- Soni ML, Beniwal RK, Yadava ND, Talwar HS (2008) Spatial distribution of soil organic carbon under agroforestry and traditional cropping system in Hyper Arid Zone of Rajasthan. *Annals of Arid Zone* **47**(1), 103-106.
- Soni ML, Yadava ND, Beniwal RK, Singh JP, Sunil K (2007) Production potential of arid legumes under grass based stripcropping system in arid rainfed condition of western Rajasthan. *Journal of Arid Legumes* **4**(1), 9-11.
- Sonone AH, Jadhav SJ, Dahat DV, Aher RP (1999) Path analysis in rice bean [*Vigna umbellata* (Thunb) Ohwi and Ohashi]. *Legume Research* **22**(3), 195-197.
- Soufframanien J, Archana J, Gopalakrishna T (2003) Intraspecific variation in the internal transcribed spacer region of rDNA in black gram (*Vigna mungo* (L.) Hepper). *Current Science* **85**(6), 798-802.
- Soufframanien J, Gopalakrishna T (2009) Cross-species amplification of microsatellite loci and diversity analyses in blackgram. *Journal of Food Legumes* **22**(1), 11-17.
- Souma C, Okumura O, Kato J (2007) Effects of hot water extract of adzuki bean on physiological functions of human subjects. *Bulletin of Hokkaido Prefectural Agricultural Experiment Stations*(91), 23-29.
- Souma J, Kobayashi K, Ogoshi A (1991a) Studies on the decrease of the population of Phialophora gregata, the causal fungus of brown stem rot of adzuki beans, by corn cultivation. 1. The populations of Phialophora gregata in the rhizospheres of corn and adzuki bean. *Memoirs of the Faculty of Agriculture, Hokkaido University* **17**(4), 473-477.
- Souma J, Kobayashi K, Ogoshi A (1991b) Studies on the decrease of the population of Phialophora gregata, the causal fungus of brown stem rot of adzuki beans, by corn cultivation. II. Comparisons of the rhizoplane and the rhizosphere microfloras between corn and adzuki bean. *Memoirs of the Faculty of Agriculture, Hokkaido University* **17**(4), 479-488.
- Souma J, Kobayashi K, Ogoshi A (1995) Relationship between root length density and decrease in population of Phialophora gregata f.sp. adzukicola. *Soil Microorganisms*(45), 33-39.
- Sridevi M, Malliah KV (2009) Phosphate solubilization by Rhizobium strains. *Indian Journal of Microbiology* **49**(1), 98-102.
- Srimathi P, Malarkodi K (2002) Influence of seed coat colour on seed quality of rice bean (*Vigna umbellata*). *Agricultural Science Digest* **22**(4), 249-251.
- Srinivasa RN, Khalak A (1993) Nitrogen and seed inoculation with different strains of rhizobia in rice bean. *Current Research - University of Agricultural Sciences (Bangalore)* **22**(2), 30-32.
- Srinivasan T, Durairaj C (2006b) Peroxidase activity as a tool for identifying bruchid [Callosobruchus maculatus (Fabricius)] resistant varieties of rice bean, "*Vigna umbellata* Thunb. (Ohwi & ohashi)". *Plant Protection Bulletin (Faridabad)* **58**(1/2), 14-19.
- Srinivasan T, Durairaj C (2007b) Biochemical basis of resistance in rice bean, *Vigna umbellata* Thunb. (Ohwi and Ohashi), against Callosobruchus maculatus F. *Journal of Entomology* **4**(5), 371-378.
- Srinivasan T, Durairaj C, Senguttuvan K (2008a) A new scoring technique for assessing bruchid resistance in wild legumes. *Journal of Experimental Zoology, India* **11**(2), 321-323.
- Srinivasarao C, Masood A, Ganeshamurthy AN, Singh KK (2003) Potassium requirements of pulse crops. *Better Crops International* **17**(1), 8-11.
- Srinives P, Hual-alai N, Saengchot S, Ngampongsai S (2000a) The use of wild relatives and gamma radiation in mungbean and blackgram breeding. 205-218.

- Srivastava GP, Srivastava VC (2003a) Effect of row spacing on yield of rice bean (*Vigna umbellata*) varieties. *Journal of Research, Birsa Agricultural University* **15**(1), 95-96.
- Srivastava JP, Soni CP (1995) Physiological basis of yield in moth bean (*Vigna aconitifolia* (Jacq.) Marechal) under rainfed conditions. *Journal of Agronomy and Crop Science* **174**(2), 73-77.
- Srivastava LS (1999) Efficacy of fungitoxicants in managing ricebean rust in Sikkim. *Journal of Hill Research* **12**(1), 86-87.
- Srivastava LS (2000) Note: new hosts of *Ampelomyces quisqualis* Ces. from Sikkim. *Journal of Hill Research* **13**(1), 41.
- Srivastava LS, Gupta DK (1990) *Eudarluca caricis* on *Uromyces appendiculatus* - a new host record for India. *Plant Disease Research* **5**(1), 100.
- Srivastava LS, Gupta DK, Verma RN (1992) Some unrecorded ricebean diseases from India. *Plant Disease Research* **7**(1), 72-76.
- Srivastava RP, Srivastava GK (2003b) Nutritional value of pulses. *Indian Journal of Agricultural Biochemistry* **16**(2), 57-65.
- Srivastava RP, Srivastava GK, Gupta RK (2001b) Nutritional quality of rice bean (*Vigna umbellata*). *Indian Journal of Agricultural Biochemistry* **14**(1/2), 55-56.
- Stillwell RC, Wallin WG, Hitchcock LJ, Fox CW (2007) Phenotypic plasticity in a complex world: interactive effects of food and temperature on fitness components of a seed beetle. *Oecologia* **153**(2), 309-321.
- Stokkermans TJW, Orlando R, Koll VSK, Carlson RW, Peters NK (1996) Biological activities and structures of *Bradyrhizobium elkanii* low abundance lipo chitin-oligosaccharides. *Molecular Plant-Microbe Interactions* **9**(4), 298-304.
- Storrie A (1997) Weed control in summer crops 1997-99. A guide to weed control in grain and forage sorghum, maize, sweet corn, soybean, sunflower, adzuki bean, cowpea, lablab, mung bean, pigeon pea, navy bean, peanut and millet. iv + 47 pp.
- Storrie A (1999) Weed control in summer crops 2000-01. iv + 54 pp.
- Storrie A, Edwards J (1998) Herbicide resistance in northern New South Wales. (DPI/211), 4 pp.
- Storrie A, Ferguson N, Cook T (2006) Weed control in summer crops 2005-06. iv + 52 pp.
- Stoyanova M, Ivanov P, Ivanova I, Mitrakov L (1989b) Biochemical characteristics of seed proteins in some species of *Phaseolus*. *Rasteniev'dni Nauki* **26**(6), 14-20. [In Bulgarian]
- Strang J (1980) Voluntary register of grain legume cultivars in Australia. 5. Adzuki bean cv. Bloodwood. *Journal of the Australian Institute of Agricultural Science* **46**(4), 264.
- Su S, Rong T, She Y, Liu J (2005) Identification of genetic diversity of cultivated adzuki bean [*Vigna angularis* var. *angularis*] germplasm resources with AFLP markers. *Journal of Sichuan Agricultural University* **23**(2), 156-162.
- Subedi KD (1998) Relay-planted green manures as a substitute for inorganic fertilizers for rice in the intensive cropping systems in Nepal. *Tropical Agriculture* **75**(4), 422-427.
- Subhash K (2004) Identification of trap crops for reducing of broomrape (*Orobanche aegyptiaca*) infestation in the succeeding mustard (*Brassica juncea*). *Annals of Agricultural Research* **25**(1), 172-173.
- Subhajit D, Sanjeev G (2009) Use of genomic resources in improvement of *Vigna* species. *Journal of Food Legumes* **22**(1), 1-10.
- Subrahmanyam D (1998) Effect of aluminium on growth, lipid peroxidation, superoxide dismutase and peroxidase activities in rice bean and French bean seedlings. *Indian Journal of Plant Physiology* **3**(3), 240-242.
- Subrahmanyam D, Rathore VS (2000) Influence of manganese toxicity on photosynthesis in ricebean (*Vigna umbellata*) seedlings. *Photosynthetica* **38**(3), 449-453.
- Subramanian D (1979a) Morphological studies in *Vigna Savi* and its related genera. *Journal of the Indian Botanical Society* **58**(3), 247-256.
- Subramanian D (1980) Effect of gamma-radiation in *Vigna*. *Indian Journal of Genetics & Plant Breeding* **40**(1), 187-194.
- Subramanian D (1982) Cyto-genetical studies on the effects of colchicine in *Vigna Savi*. and *Macroptilium Urb.* *National Academy of Sciences, Science Letters* **5**(7), 217-221.
- Suchita K, Eapen S (2005) Studies on induction of hairy roots in some leguminous crops with wild strains of *Agrobacterium rhizogenes* and their morphogenic response. *Plant Cell Biotechnology and Molecular Biology* **6**(3/4), 81-88.
- Sudbrink DL, Mack TP, Zehnder GW (1998) Alternate host plants of cowpea curculio, (Coleoptera : Curculionidae) in Alabama. *Florida Entomologist* **81**(3), 373-383.

- Sugawara K, Kobayashi K, Ogoshi A (1997) Influence of the soybean cyst nematode (*Heterodera glycines*) on the incidence of brown stem rot in soybean and adzuki bean. *Soil Biology & Biochemistry* **29**(9/10), 1491-1498.
- Suhita G, Mandal BK, Rana SK (2004) Influence of green manures raised with two levels of phosphate on succeeding crop of rice. *Indian Agriculturist* **48**(3/4), 227-231.
- Sukumaran K, Gnanamanickam SS (1980) Isolation of anti fungal compounds from Indian fodder legume plants. *Indian Journal of Microbiology* **20**(3), 204-207.
- Sultani MI, Gill MA, Anwar MM, Athar M (2007) Evaluation of soil physical properties as influenced by various green manuring legumes and phosphorus fertilization under rain fed conditions. *International Journal of Environmental Science and Technology* **4**(1), 109-118.
- Sultani MI, Gill SM, Mahmood IA, Rafiq MK (2005) Effect of various green manure crops and P levels on the physical properties of soil under rainfed conditions. *Sarhad Journal of Agriculture* **21**(4), 637-646.
- Sultani MI, Mahmood IA, Abdul R, Sartaj (2004) Soil nutrient dynamics as influenced by tropical legumes and P fertilization under rainfed conditions. *Pakistan Journal of Agricultural Sciences* **41**(1/2), 52-57.
- Sultani MI, Shaukat M, Mahmood IA, Joyia MF (2004) Wheat growth and yield response to various green manure legumes and different P levels in Pothowar Region. *Pakistan Journal of Agricultural Sciences* **41**(3/4), 102-108.
- Sumanasinghe VA, Tomooka N, Fukuoka S (1997) Phylogenetic relationships of the subgenus Ceratotropis based on random amplified polymorphic DNA. *Journal of the National Science Council of Sri Lanka* **25**(2), 73-82.
- Sumathi A, Malleshi NG, Rao SV (1995) [Elaboration of amylase activity and changes in paste viscosity of some common Indian legumes during germination.](#) *Plant Foods for Human Nutrition* **47**(4), 341-347.
- Sumathi A, Vishwanatha S, Malleshi NG, Venkat Rao S (1997) [Glycemic response to malted, popped and roller dried wheat-legume based foods in normal subjects.](#) *International Journal of Food Sciences and Nutrition* **48**(2), 103-107.
- Sundaramoorthy S, Kalra N, Sen DN (1992) Allelopathic potentials of *Acacia tortilis* on seed germination and seedling growth of some legumes. 123-124.
- Sundaramoorthy S, Sen DN (1995) Transpiration regulation in some legume crops under different fertility levels in semiarid farming. *Madras Agricultural Journal* **82**(9/10), 539-544.
- Surendra S, Choyal RR, Sushma S, Sudesh (2005) Stimulatory effect of earthworm body fluid (vermiwash) on seed germination and seedling growth of two legumes. *Journal of Phytological Research* **18**(2), 219-222.
- Suwanarat A, Lekhasoonthrakorn N, Rungchuang J, Kritapirom S (1998) Effects of intercropping groundnut and green-manure legumes to corn on the yields of corn and productivity and chemical properties of soil. *Kasetsart Journal, Natural Sciences* **32**(3), 374-384.
- Suzuki H, Fowler TJ, Tierney ML (1993) [Deletion analysis and localization of SbPRP1, a soybean cell wall protein gene, in roots of transgenic tobacco and cowpea.](#) *Plant Molecular Biology* **21**(1), 109-119.
- Suzuki K, Nishimura A, Sugimoto K, Komyoji T (1995) Sensitivity of bean gray mold fungus in Hokkaido to fluazinam. *Annals of the Phytopathological Society of Japan* **61**(4), 399-404.
- Suzuki M (1989) New summer crop cultivars registered by the Ministry of Agriculture, Forestry and Fisheries in 1989: (1) paddy rice, upland rice, sweet potato, adzuki bean, peanut, sugarbeet and buckwheat. *Japanese Journal of Breeding* **39**(4), 499-506.
- Tabiki T, Tsuchiya T (1990) Field evaluation of varietal difference to Phytophthora stem rot of adzuki beans by flooding inoculation method. *Bulletin of Hokkaido Prefectural Agricultural Experiment Stations*(60), 133-142.
- Tabil LG, Jr., Qi H, Chawla KK, Kienholz J, Crossman V, White R (1999) Physical properties of selected special crops grown in Alberta. 16 pp.
- Tabuchi A, Kamisaka S, Hoson T (1997) Purification of xyloglucan hydrolase/endotransferase from cell walls of azuki bean epicotyls. *Plant and Cell Physiology* **38**(6), 653-658.
- Tabuchi A, Mori H, Kamisaka S, Hoson T (2001) A new type of endo-xyloglucan transferase devoted to xyloglucan hydrolysis in the cell wall of azuki bean epicotyls. *Plant and Cell Physiology* **42**(2), 154-161.
- Tadano T, Ozawa K, Sakai H, Osaki M, Matsui H (1993) [Secretion of acid phosphatase by the roots of crop plants under phosphorus-deficient conditions and some properties of the enzyme secreted by lupin roots.](#) *Plant and Soil* **155/156**, 95-98.
- Tadano T, Sakai H (1991) Secretion of acid phosphatase by the roots of several crop species [rice, wheat, soyabeans, *Lupinus albus*, *Vigna angularis*, sugarbeet, tomatoes, cabbages and radishes] under phosphorus-deficient conditions. *Soil Science and Plant Nutrition* **37**(1), 129-140.

- Tagawa A, Kitamura Y, Murata S (1996) Thin layer drying characteristics of adzuki beans. *Transactions of the ASAE* **39**(2), 605-609.
- Tagawa A, Murata S (1995) Thermophysical properties of beans (Part 1). Prediction from results measured by the transient heat flow method using a probe. *Journal of the Japanese Society of Agricultural Machinery* **57**(4), 27-35.
- Takabayashi M (1978) 1977 Evaluation of candidate pesticides. (C-II) Herbicides: upland crops. *Japan Pesticide Information*(35), 47-48.
- Takahama U (1993) Regulation of peroxidase-dependent oxidation of phenolics by ascorbic acid: different effects of ascorbic acid on the oxidation of coniferyl alcohol by the apoplastic soluble and cell wall-bound peroxidases from epicotyls of *Vigna angularis*. *Plant and Cell Physiology* **34**(6), 809-817.
- Takahama U (1994) Changes induced by abscisic acid and light in the redox state of ascorbate in the apoplast of epicotyls of *Vigna angularis*. *Plant and Cell Physiology* **35**(6), 975-978.
- Takahama U (1996) Effects of fusicoccin and indole-3-acetic acid on the levels of ascorbic acid and dehydroascorbic acid in the apoplast during elongation of epicotyl segments of *Vigna angularis*. *Physiologia Plantarum* **98**(4), 731-736.
- Takahama U, Oniki T (1994a) Effects of ascorbate on the oxidation of derivatives of hydroxycinnamic acid and the mechanism of oxidation of sinapic acid by cell wall-bound peroxidases. *Plant and Cell Physiology* **35**(4), 593-600.
- Takahama U, Oniki T (1994b) The association of ascorbate and ascorbate oxidase in the apoplast with IAA-enhanced elongation of epicotyls from *Vigna angularis*. *Plant and Cell Physiology* **35**(2), 257-266.
- Takahama U, Oniki T (1997) Enhancement of peroxidase-dependent oxidation of sinapyl alcohol by an apoplastic component, 4-coumaric acid ester isolated from epicotyls of *Vigna angularis* L. *Plant and Cell Physiology* **38**(4), 456-462.
- Takahama U, Oniki T, Shimokawa H (1996) A possible mechanism for the oxidation of sinapyl alcohol by peroxidase-dependent reactions in the apoplast: enhancement of the oxidation by hydroxycinnamic acids and components of the apoplast. *Plant and Cell Physiology* **37**(4), 499-504.
- Takahashi M, Abe A (1991) Effect of feeding levels of starch on cell wall digestibility of various feeds in the rumen. *Grassland Science* **36**(4), 452-457.
- Takahashi W, Matsushita J, Kobayashi T, Tanaka O, Beppu T (1998) [Plant regeneration from epicotyl segment and callus of \*Vigna angularis\* \(cv. Tanbadainagon\)](#). *Japanese Journal of Crop Science* **67**(4), 561-567.
- Takahashi Y, Kamagata Y, Matsumura T, Uyeda I, Mink GI, Berger PH (1998) Nucleotide sequence of the 3'-terminal region of the genome of the azuki bean mosaic strain of the bean common mosaic virus and its phylogenetic relationship to viruses in the bean common mosaic virus group. *Annals of the Phytopathological Society of Japan* **64**(5), 478-480.
- Takakura S, Yamada E, Ueno K, Kon T (1979) Effects of nematicides on potato cyst nematode, *Globodera rostochiensis* (Woll.) Mulvey & Stone, 1973. *Bulletin of Hokkaido Prefectural Agricultural Experiment Stations*(42), 67-79.
- Takeda K, Shibaoka H (1978) The fine structure of the epidermal cell wall in azuki bean epicotyl. *Botanical Magazine* **91**(1024), 235-245.
- Takeda K, Shibaoka H (1981a) [Changes in microfibril arrangement on the inner surface of the epidermal cell walls in the epicotyl of \*Vigna angularis\* Ohwi et Ohashi during cell growth](#). *Planta* **151**(4), 385-392.
- Takeda K, Shibaoka H (1981b) [Effects of gibberellin and colchicine on microfibril arrangement in epidermal cell walls of \*Vigna angularis\* Ohwi et Ohashi epicotyls](#). *Planta* **151**(4), 393-398.
- Takehara A, Fukuzaki S (2002) Effect of dissolved gas concentration of deaerated water on water absorption and removal of astringency during soaking of adzuki beans. *Nippon Shokuhin Kagaku Kogaku Kaishi = Journal of the Japanese Society for Food Science and Technology* **49**(9), 605-610.
- Takehisa I, Noda C, Mori S, Yamashita M, Nakanishi H, Inoue M, Kamijima O (2001) Genetic variation and geographical distribution of azuki bean (*Vigna angularis*) landraces based on the electrophoregram of seed storage proteins. *Breeding Science* **51**(4), 225-230.
- Takenaga H, Ishibashi S, Aso S, Hyakutake H (1977) Influence of humic acid on oxidative phosphorylation of roots of lowland rice seedlings. *Journal of Agricultural Science, Japan* **22**(1), 108-117.
- Takeshita H, Konno S (1985) Application of Tes-Tape method to examine viability of stored legume seeds. *JARQ (Japan Agricultural Research Quarterly)* **19**(2), 151-152.
- Takesue K, Shibaoka H (1998) [The cyclic reorientation of cortical microtubules in epidermal cells of azuki bean epicotyls: the role of actin filaments in the progression of the cycle](#). *Planta* **205**(4), 539-546.

- Takesue K, Shibaoka H (1999) Auxin-induced longitudinal-to-transverse reorientation of cortical microtubules in nonelongating epidermal cells of azuki bean epicotyls. *Protoplasma* **206**(1/3), 27-30.
- Takeuchi Y, Tsumuraya Y (2001) In vitro biosynthesis of homogalacturonan by a membrane-bound galacturonosyltransferase from epicotyls of azuki bean. *Bioscience, Biotechnology and Biochemistry* **65**(7), 1519-1527.
- Talekar NS, Lee YH (1988) Biology of Ophiomyia centrosematis (Diptera: Agromyzidae), a pest of soybean. *Annals of the Entomological Society of America* **81**(6), 938-942.
- Talekar NS, Ying HL (1988) Biology of Ophiomyia-centrosematis (Diptera, Agromyzidae), a pest of soybean. *Annals of the Entomological Society of America* **81**(6), 938-942.
- Talpada PM, Desai HB, Patel ZN, Shukla PC (1989) Tannin content in some agro-industrial byproducts of Gujarat. *Indian Journal of Animal Nutrition* **6**(2), 181-182.
- Talwar HS, Soni ML, Beniwal RK (2005) Evaluation of crop physiological traits in clusterbean (*Cyamopsis tetragonoloba* L. Taub) and mothbean (*Vigna aconitifolia* Jacq. Marechal). *2*(2), 274-278.
- Talwar HS, Soni ML, Beniwal RK, Yadava ND, Singh JP, Rathore VS, Sunil K (2008) Genotypic variations in stomatal regulation and root traits and their association with yield parameters in moth bean (*Vigna aconitifolia* Jacq.) under rainfed conditions of arid zone. *Annals of Arid Zone* **47**(2), 169-175.
- Tanaka A (1982) Performance of crops on soils low in pH and phosphorus. (15), 157-166.
- Tanaka N, Shimuzu Y, Arima S, Harada J (1994) Fractal analysis of the structures of plant root systems. *Bulletin of the Faculty of Agriculture, Saga University*(76), 81-85.
- Tanaka S, Kondo N, Naito S (2007) Isolation of pathogenicity- and gregatin-deficient mutants of *Phialophora gregata* f.sp. *adzukicola* through *Agrobacterium tumefaciens*-mediated transformation. *Journal of General Plant Pathology* **73**(4), 242-249.
- Tanaka Y (1991) Genetic variation in age-specific fecundity of the adzuki bean weevil *Callosobruchus chinensis* (Coleoptera: Bruchidae) [reared on adzuki beans [*Vigna angularis*]]. *Applied Entomology and Zoology* **26**(2), 263-265.
- Tanaka Y (2000) Realized heritability of behavioural responsiveness to an oviposition deterring pheromone in the azuki bean weevil *Callosobruchus chinensis*. *Entomologia Experimentalis et Applicata* **96**(3), 239-243.
- Tanakamaru S (1990) Agronomic studies on seedling emergence in two-rowed barley. III. Soil resistance and seedling emergence. *Nogaku Kenkyu* **62**(2), 95-105.
- Tanasombat M, Sakurai K, Sumkeaw R, Tungtun T, Vongphrom J (2007) Intercropping of paper mulberry (*Broussonetia papyrifera*) and azuki bean (*Vigna angularis*) on sloping land, Thailand. In '' pp. 519-531. (Kasetsart University: Bangkok)
- Tang C (2008) Thermal denaturation and gelation of vicilin-rich protein isolates from three *Phaseolus* legumes: a comparative study. *LWT - Food Science and Technology* **41**(8), 1380-1388.
- Tang H, Watanabe K, Mitsunaga T (2002) Characterization of storage starches from quinoa, barley and adzuki seeds. *Carbohydrate Polymers* **49**(1), 13-22.
- Tanii A, Baba T (1979) Bacterial stem rot of adzuki bean (*Phaseolus radiatus* var. *aurea* Prain) caused by *Pseudomonas adzukicola* A. Tanii & T. Baba nov. sp. *Bulletin of Hokkaido Prefectural Agricultural Experiment Stations*(42), 29-42.
- Tanimoto E, Watanabe J (1986) Automated recording of lettuce root elongation as affected by auxin and acid pH in a new rhizometer with minimum mechanical contact to roots. *Plant and Cell Physiology* **27**(8), 1475-1487.
- Tanwar GS, Sen DN (1984) Nitrogen content of some Indian desert legumes. *Transactions of Indian Society of Desert Technology and University Centre of Desert Studies* **9**(1), 36-37.
- Tarafdar JC, Kumar D (2003) Moth bean (*Vigna aconitifolia*) cultivars compared for root nutrient uptake efficiency in arid NW India. *Journal of Arid Environments* **55**(2), 215-222.
- Tarafdar JC, Praveen K (1996) The role of vesicular arbuscular mycorrhizal fungi on crop, tree and grasses grown in an arid environment. *Journal of Arid Environments* **34**(2), 197-203.
- Tarafdar JC, Rao AV (1990) Effect of manures and fertilizers on dehydrogenase and phosphatases in the rhizosphere of arid crops. *Polish Journal of Soil Science* **23**(2), 189-193.
- Tarafdar JC, Rao AV (1997) Response of arid legumes to VAM fungal inoculation. *Symbiosis (Rehovot)* **22**(3), 265-274.
- Tarafdar JC, Yadav RS, Sharma BM (2005a) Role of arid legumes in P mobilization from plant unavailable sources. **2**(1), 20-24. (Scientific Publishers (India): Jodhpur)
- Tay DCS, Huang YK (1989) Mungbean genetic resources at the Asian Vegetable Research and Development Center. *Plant Genetic Resources Newsletter*(77), 21-24.

- Taylor JD, Teverson DM, Allen DJ, Pastor Corrales MA (1996) Identification and origin of races of *Pseudomonas syringae* pv. *phaseolicola* from Africa and other bean growing areas. *Plant Pathology* **45**(3), 469-478.
- Tebayashi SI, Matsuyama S, Suzuki T, Kuwahara Y, Nemoto T, Fujii K (1995) Quercimeritin: the third oviposition stimulant of the azuki bean weevil from the host azuki bean. *Journal of Pesticide Science* **20**(3), 299-305.
- Tejinder Pal S, Soni GL, Rattan S (1977) Biochemical evaluation of the common pulses of the Punjab state. *Journal of Research, Punjab Agricultural University* **14**(3), 318-322.
- Terai K (1992) Relation between branchiness and yield components in landraces of white azuki bean [*Vigna angularis*]. *Report of the Tohoku Branch, the Crop Science Society of Japan*(35), 93-94.
- Terai K (1993) [Relationships among branching habit, yield components, and grain yield of native varieties of white azuki bean \(\*Vigna angularis\*\)](#). *Japanese Journal of Crop Science* **62**(4), 491-495.
- Terai K (1996) Varietal differences of fruiting process within individuals among native varieties of white azuki bean [*Vigna angularis*]. *Tohoku Journal of Crop Science*(39), 89-90.
- Terai K, Horie T (1991) [Comparison of yield characters among the native varieties of white adzuki bean \(\*Vigna angularis\*\) in Akita prefecture](#). *Japanese Journal of Crop Science* **60**(1), 8-14.
- Tetarwal JP, Rana KS (2006) Impact of cropping system, fertility level and moisture-conservation practice on productivity, nutrient uptake, water use and profitability of pearl millet (*Pennisetum glaucum*) under rainfed conditions. *Indian Journal of Agronomy* **51**(4), 263-266.
- Thakkar RK, Vyas SR (1972) Absorption spectra and iron contents of leghemoglobins from root nodules of some legumes. *Indian Journal of Agricultural Sciences* **42**(8), 673-675.
- Thakkar RK, Vyas SR (1973) Behaviour of leghemoglobin under different conditions. *Indian Journal of Experimental Biology* **11**(3), 220-221.
- Thang CM, Sanh MV, Wiktorsson H (2008b) Effects of supplementation of mixed cassava (*Manihot esculenta*) and legume (*Phaseolus calcaratus*) fodder on the rumen degradability and performance of growing cattle. *Asian-Australasian Journal of Animal Sciences* **21**(1), 66-74.
- Thangavelu M (1982) Correlation, path and stability analyses in moth (*Vigna aconitifolia* (Jacq.) Marechal). *8*(3), 250.
- Thapa A, Pradhan P, Gurung N, Sharma YR, Jackson GVH, Bridge J (2008b) *Pratylenchus* on ginger in the Himalayas: alternate hosts and survival between crops. *Indian Journal of Nematology* **38**(2), 223-231.
- Thaware BL, Bendale VW, Toro VA (2005) Konkan rice bean-1 (RB-10), a new fodder rice bean variety for Konkan Region of Maharashtra. *Journal of Maharashtra Agricultural Universities* **30**(3), 295-298.
- Thaware BL, Birari SP, Dhoneksh BL, Bendale VW (1997) Genetic divergence in rice bean. *Legume Research* **20**(2), 91-96.
- Thaware BL, Birari SP, Dhoneksh BL, Bendale VW (1999) Path analysis of yield and yield attributes in different environments in rice bean. *Legume Research* **22**(3), 192-194.
- Thaware BL, Birari SP, Dhoneksh BL, Jamadagni BM (1998) Stability analysis for grain yield in rice bean. *Indian Journal of Agricultural Research* **32**(1), 17-20.
- Thaware BL, Birari SP, Dhoneksh BL, Jamadagni BM (2000) Correlation between yield and yield attributes under different environments in rice bean. *Indian Journal of Agricultural Research* **34**(1), 47-50.
- Thaware BL, Birari SP, Majgaonkar SV (1989) Studies on fodder legumes intercrops in kharif ragi. *Annals of Agricultural Research* **10**(4), 426-428.
- Thiyagu K, Jayarnani P, Nadarajan N (2008) [Pollen Pistil Interaction in Inter-Specific Crosses of \*Vigna\* sp.](#) *Cytologia* **73**(3), 251-257.
- Thomas L, Palaniappan SP (1997) Influence of sowing time on biomass production and seed yield of three green manure crops. *International Rice Research Notes* **22**(3), 29-30.
- Thomas L, Palaniappan SP (1998a) Biomass production and nitrogen accumulation of velvet beans, sunnhemp and pillipesara as influenced by plant density and phosphorus application. *Madras Agricultural Journal* **85**(5/6), 268-272.
- Thomas L, Palaniappan SP (1998b) Seed production of velvet beans, sunnhemp and pillipesara as influenced by plant density and phosphorus application. *Madras Agricultural Journal* **85**(1), 35-38.
- Tian J, Fan B (2001) Genetic variability of agronomic traits of adzuki bean in Hebei Province. *Journal of Hebei Agricultural University* **25**(2), 17-20.
- Tiburcio AF, Masdeu MA, Dumortier FM, Galston AW (1986) Polyamine metabolism and osmotic stress. I. Relation to protoplast viability. *Plant Physiology* **82**(2), 369-374.
- Tikka SBS, Asawa BM, Sharma RK, Kumar S (1980) Selection indices in moth bean (*Vigna aconitifolia* Jacq. Marechal). *Tropical Grain Legume Bulletin*(17/18), 46-48.

- Tikka SBS, Assawa BM, Kumar S (1977) Correlated response to selection in moth bean (*Vigna aconitifolia* Jacq. Marechal). *Gujarat agric. Univ. Res. J.* **3**(1), 1-4.
- Tikka SBS, Sudhir K (1976) Association analysis *Vigna aconitifolia* (Jacq.) Marechal. *Science & Culture* **42**(3), 182-183.
- Tikka SBS, Yadavendra JP, Bordia PC, Sudhir K (1976) A correlation and path coefficient analysis of components of grain yield in *Phaseolus aconitifolius* Jacq. *Genetica Agraria* **30**(2), 241-248.
- Tjahjadi C, Breene WM (1984) Isolation and characterization of adzuki bean (*Vigna angularis* cv. Takara) starch. *Journal of Food Science* **49**(2), 558-562.
- Tjahjadi C, Lin S, Breene WM (1988) Isolation and characterization of adzuki bean (*Vigna angularis* cv. Takara) proteins. *Journal of Food Science* **53**(5), 1438-1443.
- Tomar YS, Singh VP, Lal S (1986) Promising new grain legumes for irrigated areas. *Indian Farming* **36**(7), 11-12, 19.
- Tomasbarberan FA, Gil MI, Ferreres F, Rivera D, Obon C, Tomaslorente F (1993) Distribution of 8-hydroxyflavone glycosides and flavonoid aglycones in some Spanish sideritis species. *Biochemical Systematics and Ecology* **21**(4), 487-497.
- Tomer RPS, Singh K (1993) Hard seed studies in rice bean (*Vigna-umbellata*). *Seed Science and Technology* **21**(3), 679-683.
- Tomita KI (1998) Cultivation methods for stable high yields of soyabeans and adzuki beans in Abashiri district. *Bulletin of Hokkaido Prefectural Agricultural Experiment Stations*(75), 65-71.
- Tomokuni M (1981) Heteroptera of the Izu Peninsula central Japan. *Memoirs of the National Science Museum (Tokyo)*(14), 103-116.
- Tomooka N, Egawa Y, Lairungreang C, Thavarasook C (1992) collection of wild *Ceratotropis* species on the Nansei Archipelago, Japan and evaluation of bruchid resistance. *Jarq-Japan Agricultural Research Quarterly* **26**(3), 222-230.
- Tomooka N, Kaga A, Vaughan DA (2006) Diversity and use of the Asian Vigna genetic resources. *Japanese Journal of Tropical Agriculture* **50**(5), 300-303.
- Tomooka N, Kashiwaba K, Vaughan DA, Ishimoto M, Egawa Y (2000b) The effectiveness of evaluating wild species: searching for sources of resistance to bruchid beetles in the genus *Vigna* subgenus *Ceratotropis*. *Euphytica* **115**(1), 27-41.
- Tomooka N, Vaughan D, Kaga A (2002) The *Vigna angularis* complex and related species in East Asia. In '!' pp. 58-76. (International Plant Genetic Resources Institute (IPGRI): Rome)
- Tomooka N, Vaughan D, Xu R, Kashiwaba K, Kaga A (2001) Japanese native *Vigna* genetic resources. *JARQ, Japan Agricultural Research Quarterly* **35**(1), 1-9.
- Tomooka N, Mun Sup Yoon, Koji Doi, Akito Kaga and Duncan Vaughan (2002) AFLP analysis of diploid species in the genus *Vigna* subgenus *Ceratotropis*. *Genetic Resources and Crop Evolution* **49** (5)521-530
- Torikura H, Iwasaki A (1997) Injury of bean seedlings by the chironomid larvae of *Smittia* spp. (Diptera: Chironomidae, Orthocladiinae). *Annual Report of the Society of Plant Protection of North Japan*(48), 171-174.
- Toriyama K (1992) Legumes. *JICA Ref Series*(5), 27-41.
- Tran Dang X, Eiji T, Hiroyuki T, Mitsuhiro M, Tran Dang K, Chung I (2004) Evaluation on phytotoxicity of neem (*Azadirachta indica* A. Juss) to crops and weeds. *Crop Protection* **23**(4), 335-345.
- Tran Duy Q, Tran Thi H, La Tuan N (2005a) Case study project F: In situ conservation of native landraces and their wild relatives in Vietnam (VIE/01/G35) In situ conservation of agricultural biodiversity on-farm: lessons learned and policy implications. Proceedings of Vietnamese National Workshop, Hanoi, Vietnam, 30 March – 1 April 2004. In '!' (Eds NTN Hue, Tv Tuyen, NT Canh, Pv Hien, Pv Chuong, BR Sthapit and D Jarvis) pp. 133) (International Plant Genetic Resources Institute (IPGRI): Rome)
- Tribhuwan S, Dhuria RK (2007) Status of camel nutrition in arid India. In '!' pp. 134-140. (College of Veterinary & Animal Science: Rajasthan)
- Triomphe B (1999) Cropping systems with cover plants in Mexico and Honduras: experiences, lessons and prospects. *Agrobiological Management of Soils and Cropping Systems*, 75-91.
- Tripathi SN, Singh AP, Gill AS (1984) Effect of pure and mixed cropping of *Pennisetum pedicellatum* with forage legumes on forage and crude protein yields. *Indian Journal of Agronomy* **29**(3), 351-355.
- Tripathy MK, Padhi NN (1992b) Pathogenic reactions of rice bean, *Vigna umbellata* (Thunb) Ohwi & Ohashi, to root-knot nematode, *Meloidogyne incognita*. *Orissa Journal of Agricultural Research* **5**(1/2), 17-19.
- Tripathy MK, Padhi NN (1998b) Varietal susceptibility of rice bean, *Vigna umbellata* to root knot nematode, *Meloidogyne incognita*. *Indian Journal of Nematology* **28**(2), 208.

- Trivedi S, Sankhla D, Sankhla N, Upadhyaya A, Weber DJ, Smith BN (1988) Effect of paclobutrazol and salt on growth and lipid constituents of moth bean seedlings. 39-46.
- Trongkongsin K, Norman B, Norman MJT (1976) Forage potential of annual legumes grown in the dry season under irrigation. *Thai Journal of Agricultural Science* **9**(3), 145-163.
- Troszyn'ska A, Lamparski G, Kmita-Głazewska H, Barylko-Pielińska N (2003) Sensory and chemical aspects of astringency of polyphenolic compounds from legume seeds. *Polish Journal of Food and Nutrition Sciences* **12**(Suppl.1), 87-89.
- Trung BC, Yoshida S, Kobayashi Y (1985) Influence of soil moisture stress on the nitrogen nutrition and grain productivity of mungbean. *Japanese Journal of Crop Science* **54**(1), 72-78.
- Tsai BY (2001) Effect of nematode age and plant species on the penetration speed of *Meloidogyne javanica* and *Pratylenchus coffeae*. *Plant Pathology Bulletin* **10**(2), 65-70.
- Tsai S, Tai K, Chu T (1997) Differences in production and dry matter partitioning in adzuki bean (*Phaseolus angularis*) plants in spring and autumn cropping seasons. *Chinese Journal of Agrometeorology* **4**(1), 27-33.
- Tsai SL (1982) Growth analysis of rice stubble azuki bean (*Phaseolus angularis*). *National Science Council Monthly* **10**(12), 967-982.
- Tsai YP, Hwang MT, Chen SP, Liu SS (1986) An ecological study of banana mosaic. *Plant Protection Bulletin (Taichung)* **28**(4), 383-388.
- Tseng TC, Tu JC (1997) Mycoflora and mycotoxins in adzuki and mung beans produced in Ontario, Canada. *Microbios* **90**(363), 87-95.
- Tsubono Y, Kobayashi M, Tsugane S (1997) Food consumption and gastric cancer mortality in five regions of Japan. *Nutrition and Cancer-an International Journal* **27**(1), 60-64.
- Tsuchida H, Mizuno M, Kimura T, Ohara A, Saito F, Ito T (2008) Antibacterial activity of the secondary product 'Shibukiri mizu' produced during azuki paste manufacturing. *Nippon Shokuhin Kagaku Kogaku Kaishi = Journal of the Japanese Society for Food Science and Technology* **55**(12), 606-611.
- Tsuchiya S (1989) Studies on the stem rot disease of adzuki bean caused by *Phytophthora vignae* f.sp. *adzukicola* and its control. (72), 76 pp.
- Tsuji S (1982) Texture profile analysis of processed foods using the tensipresser and the multi-point mensuration method. *Journal of Texture Studies* **13**(2), 135-186.
- Tsuru S (1979) Efficiency of biological nitrogen fixation. *Agriculture and Horticulture* **54**(1), 139-142.
- Tuda M (1993) Density dependence depends on scale; at larval resource patch and at whole population. *Researches on Population Ecology* **35**(2), 261-271.
- Tuda M (1996) Temporal/spatial structure and the dynamical property of laboratory host-parasitoid systems. *Researches on Population Ecology* **38**(2), 133-140.
- Tuda M, Shima K (2002) Relative importance of weather and density dependence on the dispersal and on-plant activity of the predator *Orius minutus*. *Population Ecology* **44**(3), 251-257.
- Tuda M, Shimada M (1993) Population-level analysis on reduction in equilibrium population size of the azuki bean beetle. *Researches on Population Ecology* **35**(2), 231-239.
- Tuda M, Shimada M (1995) Developmental schedules and persistence of experimental host-parasitoid systems at two different temperatures. *Oecologia* **103**(3), 283-291.
- Tun Y, Yamaguchi H (2007a) Phylogenetic relationship of wild and cultivated *Vigna* (subgenus *Ceratotropis*, Fabaceae) from Myanmar based on sequence variations in non-coding regions of trnT-F. *Breeding Science* **57**(4), 271-280.
- Tun Y, Yamaguchi H (2008) Sequence variation of four chloroplast non-coding regions among wild, weedy and cultivated *Vigna angularis* accessions. *Breeding Science* **58**(3), 325-330.
- Tun YT, Yamaguchi H (2007b) Phylogenetic relationship of wild and cultivated *Vigna* (subgenus *Ceratotropis*, fabaceae) from Myanmar based on sequence variations in non-coding regions of trnT-F. *Breeding Science* **57**(4), 271-280.
- Turková V, Klozová E (1985a) Comparison of seed proteins in some representatives of the genus *Vigna*. *Biologia Plantarum* **27**(1), 70-73.
- Tyagi RNS, Mathur AK (1978) Pathological status of kharif pulses in Rajasthan. **8**(1), 20.
- Uday B, Garg BK, Kathju S (2002) Effect of spacing on seed yield and physiological traits in moth bean (*Vigna aconitifolia*). *Indian Journal of Agricultural Sciences* **72**(2), 76-79.
- Ueno T, Kuwahara Y, Fujii K, Taper ML, Toquenaga Y, Suzuki T (1990) D-Catechin: an oviposition stimulant of the azuki bean weevil *Callosobruchus chinensis* in the host azuki bean. *Journal of Pesticide Science* **15**(4), 573-578.

- Ujjinaiah US, Narayanappa C, Prasad SR (1999a) Standardization of the seed testing procedure for rice bean (*Vigna umbellata* (Thunb)). *Seed Research* **27**(1), 14-19.
- Ullah MA, Anwar M, Razzaq A, Zaidi SAR, Bano S, Abbasi KH (2007) Evaluation of Setaria anceps as affected by summer legumes in the Pothowar Plateau of Pakistan. *International Journal of Biology and Biotechnology* **4**(2-3), 197-199.
- Ullah MA, Maqsood A, Razzaq A, Zaidi SAR, Bano S, Abbasi KH (2007) Evaluation of Setaria anceps as affected by summer legumes in the Pothowar Plateau of Pakistan. *International Journal of Biology and Biotechnology* **4**(2/3), 197-199.
- Ullah MA, Tariq AN, Razzaq A (2007b) Effect of rice bean (*Vigna umbellata*) inter-cropping on the yield of perennial grass, *Panicum maximum* cv. Gaton under rain-fed conditions. *Journal of Agriculture and Social Sciences* **3**(2), 70-72.
- Unfried K, Schiebel K, Hemleben V (1991) Subrepeats of rDNA intergenic spacer present as prominent independent satellite DNA in *Vigna radiata* but not in *Vigna angularis*. *Gene* **99**(1), 63-68.
- Upadhyaya A, Davies TD, Sankhla N (1991) Epibrassinolide does not enhance heat shock tolerance and antioxidant activity in moth bean. *Hortscience* **26**(8), 1065-1067.
- Upadhyaya A, Davis TD, Sankhla N (1991) [Heat shock tolerance and antioxidant activity in moth bean seedlings treated with tetcyclacis](#). *Plant Growth Regulation* **10**(3), 215-222.
- Upadhyaya A, Gehlot HS, Davis TD, Sankhla N, Sankhla A, Sankhla D (1989) In vitro growth, metabolism and regeneration of moth bean callus as influenced by flurprimidol. *Biochemie und Physiologie der Pflanzen* **185**(3-4), 245-252.
- Vaishale A, Sadhana J, Seema K, Kishore P, Shashi C (1998) Effect of processing on phytate degradation and mineral solubility in pulses. *Journal of Food Science and Technology (Mysore)* **35**(4), 330-332.
- Valdez-Cepeda RD, Mendoza B, Diaz-Sandoval R, Valdes-Galicia J, Lopez-Martinez JD, De Celis EMR (2003) Power-spectrum behavior of yearly mean grain yields. *Fractals-Complex Geometry Patterns and Scaling in Nature and Society* **11**(3), 295-301.
- Valle LACd, Dias WP, Ferraz S (1996) Reaction of some plant species, mainly legumes, to the soyabean cyst nematode, *Heterodera glycines* Ichinohe. *Nematologia Brasileira* **20**(2), 30-40. [In Portuguese]
- Vamadevan VK, Nayak S, Sreedharan C, Mandal BK Studies of sunlight illumination profile in different crops. In: ' pp. 136)
- Vanaja M, Raju AS (2003) Integrated nutrient management in rice-sunflower cropping system in alfisols. *Journal of Soils and Crops* **13**(2), 211-218.
- Vanaja M, Raju AS (2004) Residual effect of organic, bio and inorganic sources of nutrients on sunflower in rice-sunflower cropping system. *Journal of Soils and Crops* **14**(2), 237-246.
- Vance CP, Boylan KLM, Maxwell CA, Heichel GH, Hardman LL (1985) Transport and partitioning of CO<sub>2</sub> fixed by root nodules of ureide and amide producing legumes. *Plant Physiology* **78**(4), 774-778.
- Varma BK (1995) Some observations on tannin content of rice bean (*Vigna umbellata*). *Journal of Research, Birsa Agricultural University* **7**(1), 89-90.
- Varsha R, Chaudhari US (2002) Psychoactive effect of the leaf extract of *Vigna trilobata* Linn., on a fresh water fish Labeo rohita. *Flora and Fauna (Jhansi)* **8**(2), 101-103.
- Vaughan D, Tomooka N, Kaga A, Doi K, Yoon M (2000a) Research on *Vigna* subgenus *Ceratotropis* genetic resources that provide insights into *in situ* conservation strategies. In: ' pp. 129-150. (National Institute of Agrobiological Resources (NIAR): Tsukuba)
- Vaughan D, Tomooka N, Xu R, Konarev A, Doi K, Kashiwaba K, Kaga A (2000b) The *Vigna angularis* complex in Japan. 159-176.
- Veer S, Yadava CPS (2001) Evaluation of some grain protectants against *Callosobruchus maculatus* on moth bean. *Indian Journal of Entomology* **63**(4), 471-474.
- Vendruscolo ECG, Schuster I, Pileggi M, Scapim CA, Molinari HBC, Marur CJ, Vieira LGE (2007) Stress-induced synthesis of proline confers tolerance to water deficit in transgenic wheat. *Journal of Plant Physiology* **164**(10), 1367-1376.
- Venkateswarlu B, Rao AV (1987) Quantitative effects of field water deficits on N<sub>2</sub> (C<sub>2</sub>H<sub>2</sub>) fixation in selected legumes grown in the Indian desert. *Biology and Fertility of Soils* **5**(1), 18-22.
- Vera-Núñez JA, Infante-Santiago JP, Velasco-Velasco V, Salgado-García S, Palma-López DJ, Grageda-Cabrera OA, Cárdenas R, Peña-Cabriales JJ (2007) Influence of P fertilization on biological nitrogen fixation in herbaceous legumes grown in acid savannah soils from the Tabasco State, Mexico. *Journal of Sustainable Agriculture* **31**(3), 25-42.

- Verdoy D, Coba de la Peña T, Redondo FJ, Lucas MM, Pueyo JJ (2006) [Transgenic \*Medicago truncatula\* plants that accumulate proline display nitrogen-fixing activity with enhanced tolerance to osmotic stress.](#) *Plant, Cell and Environment* **29**(10), 1913-1923.
- Verma DPS, Cheon CI, Hong Z (1994) Small GTP-binding proteins and membrane biogenesis in plants. *Plant Physiology* **106**(1), 1-6.
- Verma DPS, Hu CAA, Delauney AJ (1993) Genetic manipulation for proline overproduction and the control of osmoregulation in plants. 47-58.
- Verma DPS, Kim JH, Wu T (1997) Assimilation of reduced nitrogen in tropical legume nodules: regulation of de novo purine biosynthesis and peroxisome proliferation. 205-209.
- Verma K, Khanna PP (1988) Hard seeds in legumes. *Seed Research* **16**(2), 235-237.
- Verma P, Mehta U (1988) Study of physical characteristics, sensory evaluation and the effect of sprouting, cooking and dehulling on the antinutritional factors of rice bean (*Vigna-umbellata*). *Journal of Food Science and Technology-Mysore* **25**(4), 197-200.
- Verma SK (1983) Host plants of *Amsacta moorei* Butler in the Rajasthan Desert. *Bulletin of Entomology* **24**(1), 49-50.
- Verma SK (2006) Emerging insect pest problems of legumes in arid ecosystem. *Journal of Arid Legumes* **3**(1), 65-69.
- Verma SN, Singh OP (1991) Rice bean *Vigna-umbellata* a new host of soybean girdle beetle *Obereopsis-brevis* S. *Indian Journal of Plant Protection* **19**(1), 104.
- Vibha D, Simlot MM (1997) Effect of trypsin inhibitor on protein quality of black-soybean and mothbean meals. *Journal of Food Science and Technology (Mysore)* **34**(3), 208-211.
- Vieira C (1977) Performance of rice bean (*Vigna umbellata*) varieties in Vicoso, Minas Gerais state Brazil. *Tropical Grain Legume Bulletin*(No.10), 20-21.
- Vieira C, Bevitori R, et al. (1992a) Legumes. In 'Leguminosas. Vol. 16'. pp. 3-75
- Vieira RF (2002) Behavior of azuki bean cultivars under different sowing dates in Ponte Nova and Leopoldina, Minas Gerais, Brazil. *Revista Ceres* **49**(286), 705-712. [In Portuguese]
- Vieira RF, de Faria MN, Ramos JAD, Vieira C, Donzeles SML, de Freitas RTF (1998) Seed germination of six grain legumes during storage at room conditions in Vicoso, Minas Gerais State, Brazil. *Seed Science and Technology* **26**(2), 489-499.
- Vieira RF, Lima RC (2008) Performance of rice bean cultivars in Coimbra and Leopoldina, state of Minas Gerais. *Revista Ceres* **55**(2), 131-134. [In Portuguese]
- Vieira RF, Nasser LCB (1997a) Performance of Phaseolus and Vigna species in the "cerrado" of Goianira, GO, Brazil. *Cientifica (Jaboticabal)* **25**(1), 59-66.
- Vieira RF, Romeiro EMC, Souza LRPd, Donzelli MF, Vieira C (1989) Cooking time for dried beans of the genera Vigna and Phaseolus, food yield and acceptability. *Revista Ceres* **35**(208), 525-533. [In Portuguese]
- Vieira RF, de Santos CM (2000) Behaviour of rice bean cultivars in Ponte Nova, Minas Gerais state. *Comportamento de cultivares de feijão-arroz em Ponte Nova, Minas Gerais.* **47**(273), 573-578.
- Vieira RF, Vieira C (1996c) Performance of beans of Vigna and Phaseolus genera in associated cropping with maize planted simultaneously. *Pesquisa Agropecuaria Brasileira* **31**(11), 781-787.
- Vieira RF, Vieira C (2001) 'Grain legumes.' 206 pp.
- Vieira RF, Vieira C, Deandrade GA (1992) Agronomic comparison of beans Phaseolus and Vigna genera with the common bean (*Phaseolus-vulgaris* L.). *Pesquisa Agropecuaria Brasileira* **27**(6), 841-850.
- Vieira RF, Vieira C, Moura WdM (2000) Behavior of azuki beans at different planting times in Coimbra and Viçosa, state of Minas Gerais. *Revista Ceres* **47**(272), 411-420. [In Portuguese]
- Vietmeyer ND (1986) Lesser-known plants of potential use in agriculture and forestry. *Science, USA* **232**(4756), 1379-1384.
- Vijayakumar C, Khalak A (1993) Effect of rhizobium levels on growth, yield and economics of rice bean. *Current Research - University of Agricultural Sciences (Bangalore)* **22**(9-10), 127-128.
- Vijayakumari K, Siddhuraju P, Janardhanan K (1997) [Chemical composition, amino acid content and protein quality of the little-known legume \*Bauhinia purpurea\* L.](#) *Journal of the Science of Food and Agriculture* **73**(3), 279-286.
- Vijayakumari K, Siddhuraju P, Pugalenthil M, Janardhanan K (1998) [Effect of soaking and heat processing on the levels of antinutrients and digestible proteins in seeds of \*Vigna aconitifolia\* and \*Vigna sinensis\*.](#) *Food Chemistry* **63**(2), 259-264.
- Vikal Y, Satija CK (1992) Cytogenetical characterization of wild and cultivated species of *Vigna*. *Crop Improvement* **19**(2), 104-108.

- Vinh LT, Dworschak E (1985) Phytate content of some foods from plant origin from Vietnam and Hungary. *Nahrung* **29**(2), 161-166.
- Vipin C, Idris M, Tripathi RS (2005) Field evaluation of second generation anticoagulant rodenticides in moth bean crop. **2**(1), 116-119.
- Vipin C, Mohd I, Tripathi RS (2005) Field evaluation of second generation anticoagulant rodenticides in moth bean crop. In '!' pp. 116-119. (Scientific Publishers (India): Jodhpur)
- Vipin C, Tripathi RS (2005) Bio-efficacy of second generation anticoagulant rodenticides in pearl millet-moong-moth bean cropping system in Indian arid region. *Indian Journal of Plant Protection* **33**(2), 167-171.
- Vir S (1982) Varietal preference in moth (*Vigna aconitifolia* Jacq.) for the pulse beetle, *Callosobruchus maculatus* Fabricius. *Bulletin of Grain Technology* **20**(1), 3-7.
- Vir S (1983) Assessment of yield loss in mothbean and cowpea crops due to insect pests, and their control. 1215.
- Vir S (1984a) Assessment of yield loss due to yellow mosaic virus infection in mothbean. *Pesticides, India* **18**(6), 33-34.
- Vir S (1984b) Screening of mothbean *Vigna aconitifolia* Jacq varieties against the pulse beetle *Callosobruchus chinensis* Linn. *Proceedings of the Indian Academy of Sciences, Animal Sciences* **93**(1), 25-28.
- Vir S, Jindal SK, Lodha S (1984) Screening of moth bean cultivars against jassids, whitefly and yellow mosaic virus. *Annals of Arid Zone* **23**(2), 99-103.
- Virendra S, Sharma SK (2005) Response of mothbean (*Vigna aconitifolia*) to zinc and thiourea under dryland condition. *Annals of Agricultural Research* **26**(2), 313-314.
- Vishnu S, Sanjita S, Mathur OP, Purohit GR (2001) Rate and extent of dry matter and nitrogen degradability of some protein sources in goats. *Indian Journal of Animal Nutrition* **18**(1), 90-92.
- Vyas NL, Manjit S, Joshi SP (2003) Evaluation of locally available plant materials for production of oyster mushroom (*Pleurotus florida*) in arid region of India. *Journal of Eco-Physiology* **6**(1/2), 41-43.
- Vyas SP (2005b) Influence of low light on growth, yield and nitrogen metabolism of moth bean (*Vigna aconitifolia* (Jacq.) Marechal.). **2**(1), 37-38. Scientific Publishers (India): Jodhpur
- Vyas SP, Garg BK, Kathju S, Lahiri AN (2003) Relative drought tolerance of moth bean genotypes. In '!' pp. 393-397. (Scientific Publishers (India): Jodhpur)
- Wada K, Oh-Oka H, Matsubara H (1985) Ferredoxin isoproteins and their variation during growth of higher plants. *Physiologie Végétale* **23**(5), 679-686.
- Wael TK (2009) [Pollen grain micro sculpturing and its systematic applications in some taxa of \*Vigna savi\* \(Leguminosae - Papilionoideae\)](#). *Australian Journal of Basic and Applied Sciences* **3**(1), 136-141.
- Wagatsuma T, Ikarashi H, Ishii K (1985) Lead toxicity in crops - comprehensive studies on the environmental improvement for crop productions. *Journal of the Yamagata Agriculture and Forestry Society*(42), 65-74.
- Wagatsuma T, Kaneko M (1987) High toxicity of hydroxy-aluminium polymer ions to plant roots. *Soil Science and Plant Nutrition* **33**(1), 57-67.
- Waghmare SS, Pokharkar RN (1974) Host preference studies of *Aphis craccivora*, Koch. *Research Journal of Mahatma Phule Agricultural University* **5**(1), 28-35.
- Wakabayashi K, Nakano S, Soga K, Hoson T (2009) Cell wall-bound peroxidase activity and lignin formation in azuki bean epicotyls grown under hypergravity conditions. *Journal of Plant Physiology* **166**(9), 947-954.
- Wakelin SA, Warren RA, Kong L, Harvey PR (2008) Management factors affecting size and structure of soil Fusarium communities under irrigated maize in Australia. *Applied Soil Ecology* **39**(2), 201-209.
- Waliyar F, Sharma SB, Traore A (1998) Host preferences of plant parasitic nematodes associated with growth variability problem of groundnut in Niger. *International Journal of Nematology* **8**(1), 1-4.
- Wanek W, Richter A (1995) [Purification and characterization of myoinositol 6-O-methyltransferase from \*Vigna umbellata\* Ohwi et Ohashi](#). *Planta* **197**(3), 427-434.
- Wanek W, Richter A (1997) Biosynthesis and accumulation of D-ononitol in *Vigna umbellata* in response to drought stress. *Physiologia Plantarum* **101**(2), 416-424.
- Wang C, Lee P, Wu Y Field augmentation of *Orius strigicollis* (Heteroptera: Anthocoridae) for the control of thrips in Taiwan. 2001, Taipei, pp. 1-9
- Wang C, Tschen JSM (1997) The development of adzuki bean mycorrhiza associated with *Glomus mosseae*. *Bulletin of National Pingtung Polytechnic Institute* **6**(3), 163-174.
- Wang FJ, Wang TY, Wang FN (1991) Germplasm resources of grain legumes in Hainan Island. *Crop Genetic Resources*(No. 1), 7-10.
- Wang L, Cheng X, Wang S, Liang H, Zhao D, Xu N (2009a) Genetic diversity of adzuki bean germplasm resources revealed by SSR markers. *Acta Agronomica Sinica* **35**(10), 1858-1865.

- Wang L, Cheng X, Wang S, Liu C, Liang H (2009b) Transferability of SSR from adzuki bean to mungbean. *Acta Agronomica Sinica* **35**(5), 816-820.
- Wang L, Cheng X, Wang S, Xu N, Liang H, Zhao D (2009c) Genetic diversity among adzuki bean germplasm revealed by SSR markers. *Scientia Agricultura Sinica* **42**(8), 2661-2666.
- Wang ML, Barkley NA, Gillaspie GA, Pederson GA (2008) [Phylogenetic relationships and genetic diversity of the USDA Vigna germplasm collection revealed by gene-derived markers and sequencing](#). *Genetics Research* **90**(6), 467-480.
- Wang P, Pan F, Hu Z, He S, Feng X (2004) Biodiversity and phylogeny of rhizobia strains isolated from rice bean (*Vigna umbellata* L.). *Abstracts of the General Meeting of the American Society for Microbiology* **104**, 399.
- Wang QY, Jin WL (1995) Research on flower bud differentiation in adzuki bean. *Journal of Nanjing Agricultural University* **18**(1), 15-20.
- Wang SM, Redden RJ, Hu Jiapeng JP, Desborough PJ, Lawrence PL, Usher T (2001) [Chinese adzuki bean germplasm: 1. Evaluation of agronomic traits](#). *Australian Journal of Agricultural Research* **52**(6), 671-681.
- Wang XC, Tian J, Li H (1992) Identification of salt tolerance in adzuki bean germplasm resources. *Crop Genetic Resources*(3), 25-26.
- Wang XW, Kaga A, Tomooka N, Vaughan DA (2004) [The development of SSR markers by a new method in plants and their application to gene flow studies in azuki bean \[Vigna angularis \(Willd.\) Ohwi & Ohashi\]](#). *Theoretical and Applied Genetics* **109**(2), 352-360.
- Wang Z, Dong M, Yu Z (1995) Studies on growth characters and productivities of different adzuki bean ecotypes in two sowing dates. *Journal of Zhejiang Agricultural University* **21**(6), 637-641.
- Wani SH, Singh NB, Singh NG, Singh KN, Laishram JM (2007b) Stability of seed yield and its component characters in rice bean (*Vigna umbellata*) Thunb. Ohwi and Ohashi. *Environment and Ecology* **25**S(Special 4), 1094-1098.
- Wankhede DB (2003) The industrial potential of arid legume processing industry in India in the post GATT scenario - problems and opportunities. In ' pp. 287-297. (Scientific Publishers (India): Jodhpur)
- Wankhede DB, Syed HM, Syed KA (2003) Studies on effect of processing on the functional properties of moth bean, horse gram and cowpea. In ' pp. 301-306. (Scientific Publishers (India): Jodhpur)
- Washitani I, Masuda M (1990) A comparative study of the germination characteristics of seeds from a moist tall grassland community. *Functional Ecology* **4**(4), 543-557.
- Watanabe K, Taniguchi E (1986) [Syntheses and plant growth-regulating activity of 2-substituted phenylacetic acid derivatives](#). *Agricultural and Biological Chemistry* **50**(8), 2047-2052.
- Watanabe N (1984) Comparative studies on ecology of the adzuki bean weevil, *Callosobruchus chinensis* (L.) and the cowpea weevil, *C. maculatus* (F.) (Coleoptera: Bruchidae). I. Effect of seed size on creeping of females into the pile. *Japanese Journal of Applied Entomology and Zoology* **28**(4), 223-228.
- Watanabe N (1988) A new proposal for measurement of the adjusted mean crowding through consideration of size variability in habitat units. *Researches on Population Ecology* **30**(2), 215-225.
- Watanabe S, Sato M, Shiina T, Usui N, Kato K (1986) Long-term storage and distribution of crop seeds for genetic resources at the Germplasm Seed Storage Center. Yatabe.
- Watanabe T (1981) Detection of *Pythium deliense* in the Ryukyu Islands and its ecological implication. *Annals of the Phytopathological Society of Japan* **47**(4), 562-565.
- Wati RK, Theppakorn T, Benjakul S, Rawdkuen S (2009) Three-phase partitioning of trypsin inhibitor from legume seeds. *Process Biochemistry* **44**(12), 1307-1314.
- Weder JKP (2002a) Identification of food and feed legumes by RAPD-PCR. *Lebensmittel-Wissenschaft und -Technologie* **35**(6), 504-511.
- Weder JKP (2002b) [Identification of plant food raw material by RAPD-PCR: legumes](#). *Journal of Agricultural and Food Chemistry* **50**(16), 4456-4463.
- Wei S (1998) Identification of *Vigna angularis* germplasm resistance to *Cercospora canescens*. *Crop Genetic Resources*(3), 40.
- Weiss A, Hertel C, Grothe S, Ha D, Hammes WP (2007) Characterization of the cultivable microbiota of sprouts and their potential for application as protective cultures. *Systematic and Applied Microbiology* **30**(6), 483-493.
- Whitbread AM, Clem RL (2006) [Graze to grain - measuring and modelling the effects of grazed pasture leys on soil nitrogen and sorghum yield on a Vertisol soil in the Australian subtropics](#). *Australian Journal of Agricultural Research* **57**(5), 489-500.

- Wijeratne PM (1998) Variation in egg and adult production of *Callosobruchus maculatus* (F.) and *Callosobruchus chinensis* (L.) and the effect of egg density and oviposition site limitation. *Tropical Agricultural Research and Extension* **1**(2), 136-142.
- Wijeratne PM, Smith RH (1994) Frequency related oviposition behaviour and host seed preference of Brazil and Yemen strains of cowpea weevil (*Callosobruchus maculatus* (F.)). *Tropical Agriculturist* **150**, 49-64.
- Wilson KA (1980) The release of proteinase inhibitors from legume seeds during germination. *Phytochemistry* **19**(12), 2517-2519.
- Wong KH, Cheung PCK (1998) [Nutritional assessment of three Chinese indigenous legumes in growing rats. Nutrition Research](#) **18**(9), 1573-1580.
- Wongvithoonyaporn P, Bucke C, Svasti J (1998) Separation, characterization, and specificity of alpha-mannosidases from *Vigna umbellata*. *Bioscience Biotechnology and Biochemistry* **62**(4), 613-621.
- Wongwattana C, Ishizuka K (1988a) Herbicidal activity, absorption and translocation of clomeprop in plant seedlings. *Weed Research, Japan* **33**(3), 191-199.
- Wongwattana C, Ishizuka K (1988b) Metabolism of clomeprop in plant seedlings. *Weed Research, Japan* **33**(3), 200-208.
- Woo KC, James SH (1981) A pathway of ureide synthesis in N<sub>2</sub>-fixing nodules of tropical legumes. 390.
- Wood IM, Russell JS (1979) Grain legumes. 232-262.
- Wu H, Yuan F, Chen L, Duan Y (2001) Advances in soyabean cyst nematode and mechanism of soyabean resistance to *Heterodera glycines*. *Soybean Science* **20**(4), 285-289.
- Wu M, Chou L, Wang L (2002) Antioxidant activities of azuki (*Vigna angularis*) beans and pods. *Food Science and Agricultural Chemistry* **4**(2), 74-80.
- Wu SJ, Wang JS, Lin CC, Chang CH (2001) Evaluation of hepatoprotective activity of legumes. *Phytomedicine* **8**(3), 213-219.
- Wu YL, Hsu CC, Chen KF (1985) The new azuki bean variety Kaohsiung 3. *Taiwan Agriculture Bimonthly* **21**(1), 26-28.
- Wungsem R (2008) Germplasm evaluation of ricebean of Nagaland. *Journal of Food Legumes* **21**(2), 143-144.
- Xie Y, Jin W, Wang Q (2009) Morphological and anatomical studies on vegetative organs of cultivated, semi-wild and wild Adzuki beans. *Jiangsu Journal of Agricultural Sciences* **25**(1), 54-58.
- Xie Y, Xu J, Lu W, Lin G (2009) Adzuki bean: a new resource of biocatalyst for asymmetric reduction of aromatic ketones with high stereoselectivity and substrate tolerance. *Bioresource Technology* **100**(9), 2463-2468.
- Xin C, Volkaert H, Chatwachirawong P, Srinives P (2008) Utilization of information from international observation trials for the introduction of new crops: an introduction of azuki bean varieties from China to Thailand. *Journal of Crop Science and Biotechnology* **11**(1), 51-56.
- Xu D, Jia Z, Wang H, Zhu D, Pan X, Chen D, Zhou Y (2001) A new middle cowpea variety - 'Yangjiang No.40'. *Acta Horticulturae Sinica* **28**(1), 90.
- Xu HX, Jing T, Tomooka N, Kaga A, Isemura T, Vaughan DA (2008) [Genetic diversity of the azuki bean \(\*Vigna angularis\* \(Willd.\) Ohwi & Ohashi\) gene pool as assessed by SSR markers](#). *Genome* **51**(9), 728-738.
- Xu N, Cheng X, Wang L, Wang S, Liu C, Sun L, Mei L (2009) Screening and application of SSR molecular markers for genetic diversity analysis of Chinese adzuki bean germplasm resources. *Acta Agronomica Sinica* **35**(2), 219-227.
- Xu N, Cheng X, Wang S, Wang L, Zhao D (2008) Establishment of an adzuki bean (*Vigna angularis*) core collection based on geographical distribution and phenotypic data in China. *Acta Agronomica Sinica* **34**(8), 1366-1373.
- Xu R, Tomooka N, Vaughan DA (2000a) AFLP markers for characterizing the adzuki bean complex. *Crop Science* **40**(3), 808-815.
- Xu R, Tomooka N, Vaughan DA, Doi K (2000b) [The \*Vigna angularis\* complex: genetic variation and relationships revealed by RAPD analysis, and their implications for in situ conservation and domestication](#). *Genetic Resources and Crop Evolution* **47**(2), 123-134.
- Xu W (1999) Experiments on epidemiology and life habit of the cowpea weevil (*Callosobruchus maculatus* F.). *Zhejiang Nongye Kexue*(5), 222-224.
- Xu Z, Nakajima M, Suzuki Y, Yamaguchi I (2002) Cloning and characterization of the abscisic acid-specific glucosyltransferase gene from adzuki bean seedlings. *Plant Physiology* **129**(3), 1285-1295.
- Xu ZW, Huang WN, Zhang FM (1992) The accumulation of ureide and effect of some nitrogenous and carbonic compounds in germinating seedlings of adzuki bean (*Vigna angularis*). *Journal of Fujian Agricultural College* **21**(2), 186-189.

- Yadav GL (1993) Effect of spacing and phosphorus levels on yield of moth bean (*Vigna aconitifolia*). *Indian Journal of Agronomy* **38**(3), 513-514.
- Yadav GL (1998) Response of mothbean varieties to row spacings and phosphorus levels. *Indian Journal of Pulses Research* **11**(1), 38-42.
- Yadav GL, Jat BL (2005) Intercropping of mothbean varieties with pearl millet for sustainable crop production in arid eco-system. *Indian Journal of Pulses Research* **18**(2), 252-253.
- Yadav MK, Vyas SR (1971) Response of root-nodule rhizobia to saline, alkaline and acid conditions. *Indian Journal of Agricultural Sciences* **41**(10), 875-881.
- Yadav MS, Bhagat PD, Tomar AS, Prasad RD (2007) Impact of moth bean var. RMO-40 at farmers field in arid zone. *Journal of Arid Legumes* **4**(2), 139-141.
- Yadav OP, Henry A, Yadav BS (2003) Collection and evaluation of germplasm of three arid legumes from Rajasthan. In ' pp. 126-130. (Scientific Publishers (India): Jodhpur)
- Yadav RS, Kumar D, Sharma BM, Tarafdar JC (2003) Genetic variation in phytase and acid phosphatase content of legume seeds. In ' pp. 131-135. (Scientific Publishers (India): Jodhpur)
- Yadav RS, Sharma SK, Poonia BL, Dahama AK (2004) Selectivity and phytotoxicity of oxadiargyl on cumin and weeds and its residual effect on succeeding mothbean and pearl millet. *Indian Journal of Weed Science* **36**(1/2), 83-85.
- Yadav RS, Tarafdar JC (2001) Influence of organic and inorganic phosphorus supply on the maximum secretion of acid phosphatase by plants. *Biology and Fertility of Soils* **34**(3), 140-143.
- Yadav SR, Bhargava MC (2005a) Developmental response of Callosobruchus maculatus (Fab.) to different varieties/genotypes of mothbean [*Vigna aconitifolia* (Jacq.) Marechal]. *Journal of Plant Protection and Environment* **2**(1), 68-74.
- Yadav SR, Bhargava MC (2005b) Evaluation of seed protectants against the pulse beetle, Callosobruchus maculatus (Fab.) infesting stored mothbean [*Vigna aconitifolia* (Jacq.) Marechal]. *Journal of Plant Protection and Environment* **2**(2), 12-16.
- Yadav US, Walia RK (1988) On the biology of pigeonpea cyst nematode, *Heterodera cajani* Koshy, 1967. *Indian Journal of Nematology* **18**(1), 35-39.
- Yadava ND (1992) Studies on strip cropping of pearl millet + arid legumes in rainfed arid western Rajasthan. *Current Agriculture* **16**(1-2), 89-92.
- Yadava ND, Beniwal RK (2003) Competitive relationship of arid legume and pearl millet inter cropping system under rainfed condition of Western Rajasthan. In ' pp. 272-274. (Scientific Publishers (India): Jodhpur)
- Yadava ND, Beniwal RK (2006a) Performance of mothbean varieties to spacing and fertilizer application in arid western Rajasthan under rainfed condition. *Journal of Arid Legumes* **3**(2), 64-65.
- Yadava ND, Beniwal RK (2006b) Production potential of arid legume based intercropping system in arid western Rajasthan. *Current Agriculture* **30**(1/2), 49-52.
- Yadava ND, Gupta IC (1988) Strip cropping with sesame pays in arid zone. *Indian Farming* **38**(3), 6-7.
- Yadava ND, Gupta IC (1989) Effect of planting systems on different kharif pulses under rainfed condition during subnormal rainfall year and drought year in western Rajasthan. *Current Agriculture* **13**(1-2), 89-93.
- Yadava ND, Gupta IC (1992) Studies on drought resistance on kharif crops of western Rajasthan. *Annals of Agricultural Research* **13**(2), 202-204.
- Yadava ND, Rathore VS, Beniwal RK (2005) Production potential of legume based intercropping system under hyper arid condition of Rajasthan. **2**(2), 230-232.
- Yadava ND, Soni ML, Beniwal RK (2005) Productivity of arid legumes on deposited and eroded soils under arid rainfed situation of western Rajasthan. **2**(2), 237-240.
- Yamada E, Sakuma F, Hashizume K, Takahashi M, Fukuhara Y, Kobayashi K, Kondo N (2005a) Effect of *Pratylenchus penetrans* on the infection of brown stem rot in azuki bean. *Japanese Journal of Nematology* **35**(2), 71-77.
- Yamada E, Sakuma F, Hashizume K, Takahashi M, Fukuhara Y, Kobayashi K, Kondo N (2005b) Seasonal occurrence of *Pratylenchus penetrans*, *Heterodera glycines* and *Phialophora gregata* f.sp. *adzukicola* as affected by the cultivation of green manure crops. *Japanese Journal of Nematology* **35**(2), 79-86.
- Yamada S, Takeoka A, Yamanouchi M (2002) 22Na<sup>+</sup> and 36Cl<sup>-</sup> mobility in salinized excised leaves of several crop plants. *Soil Science and Plant Nutrition* **48**(1), 23-29.
- Yamada T, Moriyama R, Hattori K, Ishimoto M (2005) Isolation of two  $\alpha$ -amylase inhibitor genes of tepary bean (*Phaseolus acutifolius* A. Gray) and their functional characterization in genetically engineered azuki bean. *Plant Science* **169**(3), 502-511.

- Yamada T, Teraishi M, Hattori K, Ishimoto M (2001) Transformation of adzuki bean by *Agrobacterium tumefaciens*. *Plant Cell, Tissue and Organ Culture* **64**(1), 47-54.
- Yamagata H (1981) Fruits and perspectives of mutation breeding. *Gamma Field Symposia*(20), 11-40.
- Yamagata M, Matsumoto S, Koga N, Ae N (2001) Prominent nitrogen uptake response to cattle feces by specific plant species. In 'Developments in Plant and Soil Sciences, Volume 92.' pp. 856-857. (Kluwer Academic Publishers: Dordrecht)
- Yamagi N, Yoshida T, Inaba K, Mitsunaga T (1996) Changes in trypsin inhibitors of leguminous seeds during germination. *Memoirs of the Faculty of Agriculture of Kinki University*(29), 69-75.
- Yamaguchi H (1992) Wild and weed azuki beans in Japan. *Economic Botany* **46**(4), 384-394.
- Yamaguchi H, Nikuma Y (1996) Biometric analysis on classification of weed, wild and cultivated azuki beans. *Weed Research, Japan* **41**(1), 55-62.
- Yamaguchi T, Wakizuka T (1987) Stimulation of growth in aged crops' seeds by treatment with plant growth regulators. *International Botanical Congress Abstracts* **17**, 139.
- Yamamoto H (1994) Studies on taxonomy of the causal fungus of brown stem rot of adzuki bean and soyabean, Phialophora gregata (Allington et Chamberlain) Gams. *Memoirs of the Faculty of Agriculture, Hokkaido University* **19**(1), 57-98.
- Yamamoto H, Kobayashi K, Ogoshi A (1990) Isozyme polymorphism in Phialophora gregata isolates from adzuki bean and soybean in Japan. *Annals of the Phytopathological Society of Japan* **56**(5), 584-590.
- Yamamoto H, Kobayashi K, Ogoshi A (1991) Comparison of culture characteristics of Phialophora gregata f.sp. adzukicola and sojae in Japan. *Transactions of the Mycological Society of Japan* **32**(4), 529-533.
- Yamamoto H, Kobayashi K, Ogoshi A (1993) Restriction fragment length polymorphisms in mitochondrial DNA of Phialophora gregata in Japan. *Transactions of the Mycological Society of Japan* **34**(4), 465-471.
- Yamamoto R, Inouhe M, Masuda Y (1988) Galactose inhibition of auxin-induced growth of mono- and dicotyledonous plants. *Plant Physiology* **86**(4), 1223-1227.
- Yamamoto Y, Sano CM, Tatsumi Y, Sano H (2006) [Field analyses of horizontal gene flow among \*Vigna angularis\* complex plants](#). *Plant Breeding* **125**(2), 156-160.
- Yamanouchi M (1979) Boron nutrition in crop plants. *Bulletin of the Faculty of Agriculture, Tottori University* **31**, 37-91.
- Yamanouchi M, Akitomo M, Fujimoto T, Nagai T (1994) Effects of different kinds of sodium salts on the absorption of K<sup>+</sup>, Ca<sup>2+</sup> and Mg<sup>2+</sup> in 12 crop varieties. *Japanese Journal of Soil Science and Plant Nutrition* **65**(2), 157-164.
- Yamashita K (1991) Occurrence of azuki bean mosaic disease in Nanbu region of Aomori Prefecture. *Annual Report of the Society of Plant Protection of North Japan*(42), 40-41.
- Yamazaki M, Lopez PL, Kaku K (1988) The bioavailability of nutrients in some philippine feedstuffs to poultry. *Jarp* **22**(3), 229-234.
- Yamazaki T, Takaoka M, Katoh E, Hanada K, Sakita M, Sakata K, Nishiuchi Y, Hirano H (2003) A possible physiological function and the tertiary structure of a 4-kDa peptide in legumes. *European Journal of Biochemistry* **270**(6), 1269-1276.
- Yamini KN, Gomathinayagam P, Devasena N, Mohanbabu R (2001a) Isozyme analysis of interspecific hybrids of *Vigna* spp. *Journal of Soils and Crops* **11**(1), 36-39.
- Yamini KN, Gomathinayagam P, Devasena N, Punitha D (2001b) Studies on crossability of green gram with V. sublobata, V. aconitifolia and V. trilobata. *Journal of Soils and Crops* **11**(1), 17-23.
- Yang JL, You JF, Li YY, Wu P, Zheng SJ (2007) Magnesium enhances aluminum-induced citrate secretion in rice bean roots (*Vigna umbellata*) by restoring plasma membrane H<sup>+</sup>-ATPase activity. *Plant and Cell Physiology* **48**(1), 66-73.
- Yang JL, Zhang L, Li YY, You JF, Wu P, Zheng SJ (2006) [Citrate transporters play a critical role in aluminium-stimulated citrate efflux in rice bean \(\*Vigna umbellata\*\) roots](#). *Annals of Botany* **97**(4), 579-584.
- Yang R, Hsieh FS, Horng S (2006) Effects of search experience in a resource-heterogeneous environment on the oviposition decisions of the seed beetle, *Callosobruchus maculatus* (F.). *Ecological Entomology* **31**(4), 285-293.
- Yang RJ, Han YG (1994) Geographical distribution of *Vigna angularis* var. nipponensis in the Liaoning province of China and artificial crossing between it and V. angularis var. angularis. *Acta Agronomica Sinica* **20**(5), 607-613.
- Yano A, Yasuda K, Yamaguchi H (2004) A test for molecular identification of Japanese archaeological beans and phylogenetic relationship of wild and cultivated species of subgenus Ceratotropis (Genus *Vigna*,

- Papilionaceae) using sequence variation in two non-coding regions of the trnL and trnF genes. *Economic Botany* **58**, S135-S146.
- Yao A, Feng B, Chai Y, Gao J, Liu J, Xue S, Zhu R (2008) Effects of different tillage techniques on yield and water use efficiency in minor grain crops. *Agricultural Research in the Arid Areas* **26**(1), 97-101.
- Yao BY (1986) The development of systems to give three rice crops per year. *Jiangsu Agricultural Science (Jiangsu Nongye Kexue)*(9), 7-8.
- Yao X, Zhang J (2008) Research on preparation of bean mash with chestnut kernel. *Modern Food Science and Technology* **24**(9), 927-929.
- Yasuda K, Yamaguchi H (1996a) Phylogenetic analysis of the subgenus Ceratotropis (genus Vigna) and an assumption of the progenitor of azuki bean using isozyme variation. *Breeding Science* **46**(4), 337-342.
- Yasuda K, Yamaguchi H (1998) Life history of wild and weed azuki beans under different weeding conditions. *Journal of Weed Science and Technology* **43**(2), 114-121.
- Yasuda K, Yamaguchi H (2006) Growth and seed production in wild and weed azuki beans under shading conditions. *Journal of Weed Science and Technology* **51**(2), 61-68.
- Yasui T, Hase S (1978) Quantitative determination of oligosaccharides in beans and their products by gas-liquid chromatography. *Journal of the Japanese Society of Starch Science* **25**(4), 234-239.
- Yasui T, Tateishi Y, Ohashi H (1985a) Distribution of low molecular weight carbohydrates in the subgenus Ceratotropis of the genus Vigna (Leguminosae). *Botanical Magazine* **98**(1049), 75-87.
- Ye JS, Shan FH (1994) A study on the high-yield cultivation technology for adzuki bean. *Beijing Agricultural Sciences* **12**(2), 16-19.
- Ye XY, Ng TB (2002a) A new antifungal peptide from rice beans. *Journal of Peptide Research* **60**(2), 81-87.
- Ye XY, Ng TB (2002b) Purification of angularin, a novel antifungal peptide from adzuki beans. *Journal of Peptide Science* **8**(3), 101-106.
- Yee E, Kidwell KK, Sills GR, Lumpkin TA (1999) Diversity among selected *Vigna angularis* (azuki) accessions on the basis of RAPD and AFLP markers. *Crop Science* **39**(1), 268-275.
- Yeh MS (1986) seed coat variation in Phaseolus and Vigna species (Leguminosae) by SEM. *Journal of the Agricultural Association of China*(136), 46-54.
- Yin S, Zhang Y, Tao P, Deng X, Yin B (2008) Effects of day-length at seedling stage on growth, development and yield of adzuki bean. *Scientia Agricultura Sinica* **41**(8), 2286-2293.
- Yogesh V, Satija CK (1992b) Cytogenetical characterization of wild and cultivated species of Vigna. *Crop Improvement* **19**(2), 104-108.
- Yokoi Y, Kimura M, Hogetsu K (1978) Quantitative relationships between growth and respiration. 1. Components of respiratory loss and growth efficiencies of etiolated red bean seedlings. *Botanical Magazine, Tokyo* **91**(1021), 31-41.
- Yokoyama T, Tomooka N, Okabayashi M, Kaga A, Boonkerd N, Vaughan DA (2006b) Variation in the nod gene RFLPs, nucleotide sequences of 16S rRNA genes, Nod factors, and nodulation abilities of Bradyrhizobium strains isolated from Thai Vigna plants. *Canadian Journal of Microbiology* **52**(1), 31-46.
- Yoneyama T, Kondo M (1990) Sesbania spp., Aeschynomene indica and Crotalaria spp. are amide-exporters. *Soil Science and Plant Nutrition* **36**(4), 689-693.
- Yoneyama T, Ladha JK, Watanabe I (1987) Nodule bacteroids and Anabaena: natural <sup>15</sup>N enrichment in the legume-Rhizobium and Azolla-Anabaena symbiotic systems. *Journal of Plant Physiology* **127**(3/4), 251-259.
- Yoneyama T, Ohtani T (1983) Variations of natural <sup>13</sup>C abundances in leguminous plants. *Plant and Cell Physiology* **24**(6), 971-977.
- Yoneyama T, Yamada N, Kojima H, Yazaki J (1984) Variations of natural <sup>15</sup>N abundances in leguminous plants and nodule fractions. *Plant and Cell Physiology* **25**(8), 1561-1565.
- Yoon M, Doi K, Kaga A, Tomooka N, Vaughan DA (2000a) [Analysis of the genetic diversity in the \*Vigna minima\* complex and related species in East Asia](#). *Journal of Plant Research* **113**(1112), 375-386.
- Yoon MS, Lee J, Kim CY, Baek HJ (2007a) [Genetic relationships among cultivated and wild \*Vigna angularis\* \(Willd.\) Ohwi et Ohashi and relatives from Korea based on AFLP markers](#). *Genetic Resources and Crop Evolution* **54**(4), 875-883.
- Yopp JH, Mandava NB, Sasse JM (1981) Brassinolide, a growth-promoting steroidal lactone. I. Activity in selected auxin bioassays. *Physiologia Plantarum* **53**(4), 445-452.
- Yopp JH, Mandava NB, Thompson MJ, Sasse JM (1981) Activity of brassinosteroid in selected bioassays in combination with chemicals known to synergize or retard responses to auxin and gibberellin. *138-145*.

- Yoshiba Y, Kiyosue T, Katagiri T, Ueda H, Mizoguchi T, Yamaguchi-Shinozaki K, Wada K, Harada Y, Shinozaki K (1995) [Correlation between the induction of a gene for Δ1-pyrroline-5-carboxylate synthetase and the accumulation of proline in \*Arabidopsis thaliana\* under osmotic stress.](#) *Plant Journal* **7**(5), 751-760.
- Yoshida H, Saiki M, Tomiyama Y, Mizushina Y (2008) Positional distribution of fatty acids in triacylglycerols and phospholipids from adzuki beans (*Vigna angularis*). *European Journal of Lipid Science and Technology* **110**(2), 158-163.
- Yoshida H, Tomiyama Y, Yoshida N, Mizushina Y (2009a) [Characteristics of lipid components, fatty acid distributions and triacylglycerol molecular species of adzuki beans \(\*Vigna angularis\*\)](#). *Food Chemistry* **115**(4), 1424-1429.
- Yoshida H, Tomiyama Y, Yoshida N, Mizushina Y (2009b) Profiles of lipid components, fatty acid compositions and triacylglycerol molecular species of adzuki beans (*Vigna angularis*). *Journal of the American Oil Chemists' Society* **86**(6), 545-552.
- Yoshida H, Tomiyama Y, Yoshida N, Saiki M, Mizushina Y (2008) [Lipid classes, fatty acid compositions and triacylglycerol molecular species from adzuki beans \(\*Vigna angularis\*\)](#). *Journal of Food Lipids* **15**(3), 343-355.
- Yoshida H, Yoshida N, Tomiyama Y, Saiki M, Mizushina Y (2008) [Distribution profiles of tocopherols and fatty acids of phospholipids in adzuki beans \(\*Vigna angularis\*\)](#). *Journal of Food Lipids* **15**(2), 209-221.
- Yoshida K (1987) Studies on seed size variation in grain legumes. *Memoirs of the Faculty of Agriculture, Hokkaido University* **15**(4), 385-434.
- Yoshida K, Bando A, Yanagisawa K, Gotoh K (1986) [Effects of TIBA \(2,3,5-triiodobenzoic acid\) on growth and yield of azuki bean, \*Vigna angularis\*.](#) *Japanese Journal of Crop Science* **55**(3), 345-351.
- Yoshida K, Sato H (1990) [The extent and its source of variation for characteristics related to seed quality of adzuki beans. I. Seed size variation among growers' lots in Hokkaido area.](#) *Japanese Journal of Crop Science* **59**(3), 450-454.
- Yoshida K, Sato H, Sato M (1995) [The extent of variation and its source for characteristics related to seed quality of adzuki beans. III. The water uptake of seeds and hardseededness.](#) *Japanese Journal of Crop Science* **64**(1), 7-13.
- Yoshida K, Sato H, Ueshima H, Ishii N, Sato M (1991) [The extent and source of variation in seed quality characteristics in adzuki bean. 2. Variation in seed coat colour among growers batches in Hokkaido.](#) *Japanese Journal of Crop Science* **60**(2), 234-240.
- Yoshida K, Sato Y, Okuno R, Kameda K, Isobe M, Kondo T (1996) Structural analysis and measurement of anthocyanins from colored seed coats of *Vigna*, *Phaseolus*, and *Glycine* legumes. *Bioscience, Biotechnology and Biochemistry* **60**(4), 589-593.
- Yoshida T, Gichuki EM (1983) Use of solar heat to control stored-product insect pests: bean depth. *Scientific Reports of the Faculty of Agriculture Okayama University*(61), 5-8.
- Yoshida T, Hayashi K, Toko K, Yamafuji K (1988) [Effect of anoxia on the spatial pattern of electric potential formed along the root.](#) *Annals of Botany* **62**(5), 497-507.
- Yoshida T, Igarashi H, Shinoda K (1987) Life history of *Callosobruchus chinensis* (L.) (Coleoptera, Bruchidae). 471-477.
- Yoshikawa M, Ogura S, Tatsumi M (1977) [Some properties of proteinase inhibitors from adzuki beans \(\*Phaseolus angularis\*\)](#). *Agricultural and Biological Chemistry* **41**(11), 2235-2239.
- Yoshioka SI (1981) Studies on applying waste water discharged from potato starch factories to arable land. Part 3. Effects of waste water application on the yield of upland crops and soil mineral nutrients. *Research Bulletin of the Hokkaido National Agricultural Experiment Station*(130), 99-112.
- Yoshiyasu Y, Nakajima T (2004) Host suitability of two leguminous plants for *Matsumuraes falcana* (Lepidoptera: Tortricidae). *Esakia*(44), 211-215.
- Yost R, Evans D (1988) Green manure and legume covers in the tropics. (055), 44□pp.
- Yousif AM, Batey IL, Larroque OR, Curtin B, Bekes F, Deeth HC (2003) Effect of storage of adzuki bean (*Vigna angularis*) on starch and protein properties. *Lebensmittel-Wissenschaft und -Technologie* **36**(6), 601-607.
- Yousif AM, Deeth HC (2003) [Effect of storage time and conditions on the cotyledon cell wall of the adzuki bean \(\*Vigna angularis\*\)](#). *Food Chemistry* **81**(2), 169-174.
- Yousif AM, Deeth HC, Caffin NA, Lisle AT (2002) Effect of storage time and conditions on the hardness and cooking quality of adzuki (*Vigna angularis*). *Lebensmittel-Wissenschaft und -Technologie* **35**(4), 338-343.
- Yousif AM, Kato J, Deeth HC (2003) Effect of storage time and conditions on the seed coat colour of Australian adzuki beans. *Food Australia* **55**(10), 479-484.

- Yousif AM, Kato J, Deeth HC (2007) Effect of storage on the biochemical structure and processing quality of adzuki bean (*Vigna angularis*). *Food Reviews International* **23**(1), 1-33.
- Yu S, Li R, Zhang R, Song Y, Zhang Z, Zhang D (2002) Fluorescence in situ hybridization of  $\Delta'$ -pyrroline-5-carboxylate synthetase (P5CS) gene on rice chromosome. *Journal of Huazhong Agricultural University* **21**(1), 1-4.
- Yunes ANA, Andrade MTd, Sales MP, Morais RA, Fernandes KVS, Gomes VM, Xavier-Filho J (1998) [Legume seed vicilins \(7S storage proteins\) interfere with the development of the cowpea weevil \(\*Callosobruchus maculatus\* \(F\)\)](#). *Journal of the Science of Food and Agriculture* **76**(1), 111-116.
- Zade VR, Dighe RS, Zode NG, Kausal RT (1992) Standardization of seed testing procedure for rice bean. *Annals of Plant Physiology* **6**(2), 294-296.
- Zade VR, Zode NG, Dighe RS, Changade SP (1993) Seed development and maturation studies in ricebean: (*Vigna umbellata* Thunb). *Annals of Plant Physiology* **7**(1), 68-72.
- Zaki FA, Bhatti DS (1986a) Effect of pigeon-pea cyst nematode, *Heterodera cajani* Koshy, 1967 on macro- and micro-nutrients of pigeon-pea and 'moth'. *Indian Journal of Nematology* **16**(1), 103-105.
- Zaki FA, Bhatti DS (1986b) Pathogenicity of pigeon-pea cyst nematode, *Heterodera cajani* Koshy, 1967 on some pulse crops. *Indian Journal of Nematology* **16**(1), 30-35.
- Zaroug M (1979) Studies on nutritional and symbiotic relations of tropical forage legumes. *Dissertation Abstracts International, B* **40**(2), 522.
- Zaroug MG, Munns DN (1980) Screening strains of Rhizobium for the tropical legumes *Clitoria ternatea* and *Vigna trilobata* in soils of different pH. *Tropical Grasslands* **14**(1), 28-33.
- Zeng Y, Liu F, Liu B, Fan Z (2006) Study on in vitro carbohydrate digestion rates of cereal foods. *Journal of China Agricultural University* **11**(1), 95-99.
- Zhang C, Lu Q, Verma DPS (1995) Removal of feedback inhibition of  $\Delta 1$ -pyrroline-5- carboxylate synthetase, a bifunctional enzyme catalyzing the first two steps of proline biosynthesis in plants. *Journal of Biological Chemistry* **270**(35), 20491-20496.
- Zhang M, Jin W, Wang J, Sun S, Qin X, Zhang Z (2007) Insecticidal efficacy of microwave treatment against *Callosobruchus chinensis* (L.) (Coleoptera: Bruchidae) and its effects on the germination rate of adzuki beans. *Acta Entomologica Sinica* **50**(9), 967-974.
- Zhang NC, Lin CM (1992) Analysis of the ecological characteristics of food legumes. *Crop Genetic Resources*(1), 4-5.
- Zhang Y, Liu F, Jia R, Xie X (2009) Effect of seed-soaking in La (NO<sub>3</sub>)<sub>3</sub> solution on growth and antioxidant enzyme activity of adzuki bean seedlings under saline-alkali stress. *Journal of Ecology and Rural Environment* **25**(4), 12-18.
- Zhang Y, Liu JW, Zheng WJ, Wang L, Zhang HY, Fang GZ, Wang S (2008) Optimization and validation of enzyme-linked immunosorbent assay for the determination of endosulfan residues in food samples. *Journal of Environmental Science and Health. Part B, Pesticides, Food Contaminants, and Agricultural Wastes* **43**(2), 127-133.
- Zhang Y, Yang C, Wang Y (2007) Cloning and analysis of metallothionein gene from *Tamarix androssowii*. *Bulletin of Botanical Research* **27**(3), 293-296.
- Zhang Z, Zhou MD, Rao VR (2002) Plant genetic resources network in East Asia. Proceedings of the meeting for the regional network for conservation and use of plant genetic resources in East Asia, Ulaanbaatar, Mongolia, 13-16 August, 2001. vi + 137 pp.
- Zhao Q, Huang X, Lou Y, Weber N, Proksch P (2007) Effects of ethanol extracts from Adzuki bean (*Phaseolus angularis* Wight.) and Lima bean (*Phaseolus lunatus* L.) on estrogen and progesterone receptor phenotypes of MCF-7/BOS cells. *Phytotherapy Research* **21**(7), 648-652.
- Zhao Q, Li B, Weber N, Lou Y, Proksch P (2005) Estrogen-like effects of ethanol extracts from several Chinese legumes on MCF-7 cell. *European Food Research and Technology* **221**(6), 828-833.
- Zhao T, Feng G, Liu S (2000) A study of the accumulation and partitioning of nitrogen, phosphorus and potassium in adzuki bean (*Phaseolus angularis*). *Journal of Hebei Agricultural University* **23**(3), 16-19.
- Zheng D (1992) Germplasm resources of food crops in Shennongjia and Sanxia regions. *Crop Genetic Resources*(2), 1-3.
- Zheng JY, Irihara K, Miyashita N, Terauchi R, Uchiyama H, Nakata M, Tanaka R, Morikawa H (1995a) Phylogenetic relationships between six species of *Phaseolus* and *Vigna* as determined by RFLP analysis of ribosomal DNA. *Japanese Journal of Genetics* **70**(5), 615-625.

- Zheng JY, Nakata M, Irifune K, Tanaka R, Morikawa H (1993a) [Fluorescent banding pattern analysis of eight taxa of \*Phaseolus\* and \*Vigna\* in relation to their phylogenetic relationships](#). *Theoretical and Applied Genetics* **87**(1/2), 38-43.
- Zheng JY, Nakata M, Uchiyama H, Morikawa H, Tanaka R (1991a) [Giemsa C-banding patterns in several species of \*Phaseolus\* L. and \*Vigna\* Savi, Fabaceae](#). *Cytologia* **56**(3), 459-466.
- Zheng S (1995) Seed quality of summer adzuki bean in northern Kyushu. *Report of the Kyushu Branch of the Crop Science Society of Japan*(61), 75-76.
- Zheng S, Kawagoe Y (1998) [Effect of summer sowing on the seed quality of azuki bean in northern Kyushu, Japan](#). *Japanese Journal of Crop Science* **67**(4), 473-477.
- Zheng SH, Inouye J (1990) [Comparison of elongation force, stem thickness and evolution of ethylene in seedlings of several epigeal leguminous crops](#). *Japanese Journal of Crop Science* **59**(2), 277-282.
- Zheng SH, Inouye J (1991) [The response to vertical load in seedlings of several epigeal leguminous crops](#). *Japanese Journal of Crop Science* **60**(3), 421-426.
- Zheng SH, Jin ID, Inouye J (1993) [Elongation inhibition, stem enlargement, and histological stem changes caused by vertical load in several epigeal leguminous crops](#). *Japanese Journal of Crop Science* **62**(4), 502-508.
- Zheng SH, Kawabata M (2000) [Exudation of amino acids and sugars from imbibing seeds of several leguminous crops](#). *Japanese Journal of Crop Science* **69**(3), 380-384.
- Zhu B, Su J, Chang M, Verma DPS, Fan Y, Wu R (1998) [Overexpression of a  \$\Delta 1\$ -pyrroline-5-carboxylate synthetase gene and analysis of tolerance to water- and salt-stress in transgenic rice](#). *Plant Science (Limerick)* **139**(1), 41-48.
- Zhu W, Huang Q (1998) Studies on embryology of *Vigna angularis* (Willd) Ohiwi & Ohishi. I. The microsporogenesis, the development of male gametophytes and starch granule distribution. *Journal of Hunan Agricultural University* **24**(2), 99-104.
- Zhu W, Sheng X, Huang Q, Zhao B (2000) Studies on embryology of *Vigna angularis*. II. The megasporogenesis and the development of female gametophytes. *Journal of Hunan Agricultural University* **26**(3), 178-181.
- Zhu Z, Wang X (2003) Identification of pathogen causing Phytophthora, stem rot of adzuki bean and screening for resistant germplasm. *Acta Phytophylacica Sinica* **30**(3), 289-294.
- Zink D, Schumann K, Nagl W (1994) Restriction fragment length polymorphisms of the phytohemagglutinin genes in *Phaseolus* and *Vigna* (Leguminosae). *Plant Systematics and Evolution* **191**(3/4), 131-146.
- Zong X, Kaga A, Tomooka N, Wang X, Han O, Vaughan D (2003) [The genetic diversity of the \*Vigna angularis\* complex in Asia](#). *Genome* **46**(4), 647-658.
- Zong X, Vaughan D, Kaga A, Tomooka N, Wang X, Guan J, Wang S (2003a) Genetic diversity in *Vigna angularis* revealed by AFLP analysis. *Acta Agronomica Sinica* **29**(4), 562-568.
- Zong X, Vaughan D, Tomooka N, Kaga A, Wang X, Guan J, Wang S (2003b) Preliminary study on geographical distribution and evolutionary relationships between cultivated and wild adzuki bean (*Vigna angularis* var. *angularis* and var *nipponensis*) by AFLP analysis. *Scientia Agricultura Sinica* **36**(4), 367-374.
- Zong X, Vaughan D, Tomooka N, Kaga A, Wang X, Guan J, Wang S (2003c) [Preliminary study on geographical distribution and evolutionary relationships between cultivated and wild adzuki bean \(\*Vigna angularis\* var. \*angularis\* and var. \*nipponensis\*\) by AFLP analysis](#). *Plant Genetic Resources: Characterization and Utilization* **1**(2/3), 175-183.